

MIHA Data Snapshot, Fresno County by Education, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Fresno Cou	nty	<high schoo<="" th=""><th> </th><th>High School Graduate</th><th>e or GED</th><th>Some College</th><th>2</th><th>College Gradua</th><th>ate</th></high>		High School Graduate	e or GED	Some College	2	College Gradua	ate
	% 95% CI	Ν	% 95% CI	Ν	% 95% CI	N	% 95% CI	N	% 95% CI	Ν
Total	100	15,400	24.9 19.6 - 30.2	3,700	21.8 17.0 - 26.7	3,300	32.9 27.5 - 38.3	5,000	20.3 16.4 - 24.3	3,100
Health Status before Pregnancy										
In good to excellent health	89.1 85.2 - 93.	0 13,700	82.5 * 72.1 - 93.0	3,100	87.0 * 76.7 - 97.3	2,900	91.4 86.7 - 96.1	4,500		
Chronic conditions										
Diabetes ⁺	3.0 * 0.5 - 5.	4 500								
Hypertension ⁺	2.9 * 0.9 - 4.	9 400					1.2 [*] 0.1 - 2.3	<100		
Asthma ⁺	13.3 9.0 - 17.	5 2,000	10.6 * 2.3 - 18.9	400			18.2 10.2 - 26.1	900	15.2 * 3.9 - 26.4	500
Nutrition and Weight										
Daily folic acid use, month before pregnancy	27.1 22.0 - 32.	2 4,200	21.4 11.0 - 31.7	800	23.1 12.7 - 33.5	800	25.4 16.5 - 34.4	1,300	40.6 29.3 - 52.0	1,200
Overweight before pregnancy	28.0 22.6 - 33.	3 4,000	25.4 12.9 - 37.9	800	36.3 23.4 - 49.2	1,100	30.8 21.6 - 39.9	1,500	16.8 9.9 - 23.7	500
Obese before pregnancy	24.2 19.0 - 29.		39.7 25.6 - 53.7	1,300	17.5 8.3 - 26.7	500	25.4 16.8 - 34.0	1,200	13.1 5.7 - 20.5	400
Inadequate weight gain during pregnancy	16.6 11.9 - 21.		17.9 * 6.3 - 29.5	400	24.0 11.1 - 36.9	600	13.9 6.9 - 20.8	600	12.3 6.3 - 18.3	300
Excessive weight gain during pregnancy	43.1 36.9 - 49.		29.5 15.3 - 43.7	700	33.7 20.7 - 46.6	900	48.7 38.2 - 59.2	2,100	57.0 45.3 - 68.8	1,500
Food insecurity during pregnancy	16.1 12.1 - 20.	2 2,500	19.9 9.5 - 30.2	700	20.1 10.1 - 30.0	700	15.4 9.2 - 21.5	800	5.9 * 0.8 - 10.9	200
Intimate Partner Violence (IPV) and Depressive Symptoms										
Physical or psychological IPV during pregnancy	8.0 4.8 - 11.	2 1,200	9.2 * 2.3 - 16.2	300	8.0 * 1.1 - 14.9	300	6.8 2.8 - 10.7	300		
Prenatal depressive symptoms	14.1 9.9 - 18.	,	8.7 * 1.6 - 15.7	300	18.5 8.5 - 28.5	600	14.9 8.1 - 21.8	700	14.2 * 3.8 - 24.6	400
Postpartum depressive symptoms	12.2 8.5 - 16.	0 1,900	10.5 * 3.2 - 17.7	400	10.5 * 3.4 - 17.5	300	13.2 7.0 - 19.3	700	14.9 * 4.5 - 25.3	500
Hardships and Support during Pregnancy										
Experienced two or more hardships during childhood	29.8 24.3 - 35.	2 4,400	29.2 17.2 - 41.1	1,000	33.9 21.3 - 46.4	1,100	35.6 26.1 - 45.1	1,700	15.2 * 5.0 - 25.4	400
Homeless or did not have a regular place to sleep	2.5 * 0.7 - 4.	3 400								
Moved due to problems paying rent or mortgage	6.0 3.1 - 8.	9 900			5.6 * 2.2 - 9.0	200	7.6 * 1.6 - 13.5	400		
Woman or partner lost job	12.1 8.5 - 15.		10.5 * 2.8 - 18.3	400	13.1 5.6 - 20.6	400	16.1 8.9 - 23.2	800	5.1 * 1.2 - 9.0	200
Woman or partner had pay or hours cut back	11.8 7.8 - 15.	,	13.1 * 3.0 - 23.1	500	8.6 * 2.0 - 15.2	300	17.5 9.5 - 25.6	900	5.6 [*] 0.9 - 10.3	200
Became separated or divorced	10.2 6.1 - 14.		7.3 * 0.6 - 14.0	300	9.4 * 2.4 - 16.3	300	14.4 * 5.9 - 22.9	700		
Had no practical or emotional support	3.1 [*] 0.9 - 5.	3 500								
Substance Use										
Any smoking, 3 months before pregnancy	13.7 9.3 - 18.		20.8 8.8 - 32.7	800	10.9 * 2.4 - 19.4	400	17.5 9.8 - 25.2	900	2.9 * 1.0 - 4.8	<100
Any smoking, 3rd trimester [†]	3.7 * 1.5 - 5.		9.4 * 1.6 - 17.1	300			2.6 * 0.1 - 5.0	100		
Any smoking, postpartum	6.0 3.0 - 8.		13.9 * 3.7 - 24.0	500			3.0 * 0.6 - 5.4	100	1.9 [*] 0.4 - 3.3	<100
Any binge drinking, 3 months before pregnancy	15.3 10.9 - 19.				14.4 * 4.8 - 24.1	500	20.5 12.0 - 28.9	1,000	23.7 12.2 - 35.3	700
Any alcohol use, 3rd trimester [†]	3.1 1.7 - 4.	5 500					3.5 * 0.6 - 6.3	200	9.0 3.7 - 14.2	300
Pregnancy Intention and Family Planning										
Mistimed or unwanted pregnancy	27.0 21.9 - 32.		27.3 15.5 - 39.1	1,000	43.9 31.1 - 56.8	1,400	26.2 17.8 - 34.5	1,300	11.4 5.7 - 17.0	300
Unsure of pregnancy intentions	18.9 14.0 - 23.		26.0 14.6 - 37.4	1,000	17.1 * 6.5 - 27.6	600	19.8 10.9 - 28.6	1,000	10.1 * 3.2 - 17.0	300
Postpartum birth control use ⁺	82.1 77.4 - 86.	7 12,400	79.4 69.0 - 89.8	2,900	84.7 * 74.7 - 94.8	2,700	84.5 77.1 - 91.8	4,100	77.7 67.1 - 88.4	2,300
Infant Sleep and Breastfeeding										
Placed infant on back to sleep	80.2 75.6 - 84.		75.0 63.7 - 86.3	2,700	84.5 75.5 - 93.5	2,800	77.4 68.8 - 86.1	3,800	86.3 80.6 - 92.1	2,600
Infant always or often shared bed	41.5 35.8 - 47.		42.4 29.4 - 55.5	1,500	44.0 31.4 - 56.6	1,400	43.7 33.9 - 53.5	2,100	33.4 21.3 - 45.5	1,000
Intended to breastfeed, before birth	89.5 86.0 - 93.		87.1 * 78.4 - 95.8	3,100	87.6 * 79.2 - 96.1	2,900	88.2 82.2 - 94.2	4,300		
Intended to breastfeed exclusively, before birth	52.2 46.4 - 58.	0 7,900	39.9 26.9 - 53.0	1,400	43.3 30.6 - 56.1	1,400	58.6 48.8 - 68.3	2,900	68.1 56.6 - 79.6	2,100
Breastfeeding practices										
Any breastfeeding, 1 month after delivery	76.3 71.5 - 81.		76.6 66.0 - 87.3	2,700	62.1 49.4 - 74.9	2,000	80.5 73.5 - 87.5	3,900	87.3 * 79.2 - 95.3	2,600
Exclusive breastfeeding, 1 month after delivery	36.7 31.1 - 42.		40.2 26.3 - 54.2	1,400	26.8 15.7 - 37.9	900	35.4 26.2 - 44.5	1,700	47.2 35.4 - 59.0	1,400
Any breastfeeding, 3 months after delivery	51.2 44.4 - 57.		40.6 26.1 - 55.2	1,100	38.0 23.0 - 53.0	1,000	54.9 43.5 - 66.3	2,100	72.9 60.8 - 85.0	1,400
Exclusive breastfeeding, 3 months after delivery	18.5 14.1 - 22.	9 2,000			13.2 * 3.2 - 23.2	300	20.9 13.0 - 28.9	800	42.4 30.2 - 54.5	800



MIHA Data Snapshot, Fresno County by Education, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Free	sno Count	ty		<high schoo<="" th=""><th>l</th><th>High School Gradu</th><th>ate or GED</th><th></th><th>Some College</th><th>e</th><th>Co</th><th>llege Gradua</th><th>ate</th></high>	l	High School Gradu	ate or GED		Some College	e	Co	llege Gradua	ate
	%	95% CI	N	%	95% CI	N	% 95% CI	N	%	95% CI	N	%	95% CI	N
Service Utilization and Health Insurance Coverage														
Had a usual source of pre-pregnancy care	66.9 61	5 - 72.4	10,300	57.0	44.0 - 70.0	2,100	70.5 59.4 - 81	7 2,300	65.2	55.7 - 74.7	3,200	78.7	69.7 - 87.7	2,400
Initiated prenatal care in 1st trimester	86.4 82	.3 - 90.6	13,100	84.0	75.0 - 93.0	3,100	83.8 75.0 - 92	6 2,700	84.2	75.9 - 92.5	4,200			
Mom or infant needed but couldn't afford care postpartum	7.7 4	.7 - 10.8	1,200				9.1 * 2.0 - 16	2 300	9.9	* 4.1 - 15.8	500			
Participated in WIC during pregnancy ⁺	69.5 65	.4 - 73.6	10,500	88.7	* 79.7 - 97.7	3,300	94.0 * 90.0 - 97	9 3,100	66.6	57.8 - 75.3	3,300	20.4	9.3 - 31.5	600
Received CalFresh (food stamps) during pregnancy	47.5 41	9 - 53.0	7,200	76.6	65.9 - 87.4	2,800	46.1 33.2 - 59	1 1,500	46.1	36.3 - 56.0	2,300	13.1 *	* 2.9 - 23.3	400
Pre-pregnancy insurance														
Medi-Cal	46.9 41	.4 - 52.4	7,200	78.8	68.6 - 88.9	2,900	52.9 40.0 - 65	8 1,700	40.1	30.2 - 49.9	2,000	11.6 *	* 1.7 - 21.4	400
Uninsured	22.3 17	.5 - 27.0	3,400	17.6	8.1 - 27.1	700	30.3 18.2 - 42	5 1,000	25.8	17.1 - 34.5	1,300	14.2	6.0 - 22.3	400
Private	28.9 24	.8 - 33.0	4,400				15.1 7.5 - 22	8 500	31.2	22.7 - 39.6	1,500	72.3	61.0 - 83.7	2,200
Prenatal insurance														
Medi-Cal	72.2 68	8.5 - 75.9	11,000				89.4 * 83.2 - 95	7 2,900	71.5	63.8 - 79.3	3,500	21.7	10.9 - 32.6	700
Uninsured		0.0 - 1.4	100											
Private	25.0 21	.4 - 28.5	3,800				7.8 * 2.6 - 13	0 300	24.1	17.0 - 31.1	1,200	74.4	63.3 - 85.4	2,300
Maternal postpartum insurance														
Medi-Cal	60.8 55	.8 - 65.8	9,300	83.0	73.4 - 92.5	3,000	75.6 64.9 - 86	3 2,500	60.0	50.5 - 69.5	3,000	19.8	9.1 - 30.5	600
Uninsured	10.6 6	5.9 - 14.3	1,600	12.7	* 4.1 - 21.3	500	16.8 * 6.9 - 26	7 500	9.3	* 3.0 - 15.6	500			
Private	26.6 22	2.8 - 30.4	4,100				5.9 * 1.8 - 10	0 200	28.0	19.8 - 36.3	1,400	75.3	64.3 - 86.2	2,300
Infant health insurance														
Medi-Cal	71.7 67	'.7 - 75.7	10,800				87.6 80.4 - 94	7 2,800	69.8	61.5 - 78.0	3,400	23.9	12.6 - 35.2	700
Uninsured	1.9* 0).6 - 3.2	300											
Private	25.0 21	3 - 28.7	3,800				6.5 * 1.8 - 11	3 200	26.9	19.0 - 34.7	1,300	71.4	59.9 - 82.8	2,200
Maternal Demographics														
Unmarried	51.8 46	5.3 - 57.3	7,900	61.7	49.2 - 74.1	2,300	68.6 56.8 - 80	4 2,300	50.5	40.7 - 60.2	2,500	20.9	9.6 - 32.2	600
Race/Ethnicity														
White	19.9 15	.9 - 23.9	3,000				9.8 * 2.2 - 17	5 300	20.0	13.3 - 26.6	1,000	43.1	32.0 - 54.3	1,300
Hispanic	61.7 56	5.6 - 66.8	9,400	74.5	63.0 - 86.0	2,800	69.9 58.0 - 81	9 2,300	65.3	56.8 - 73.8	3,200	29.2	16.8 - 41.5	900
Asian/Pacific Islander	12.1 8	8.0 - 16.2	1,800				15.3 * 4.7 - 25	8 500	5.9	* 0.8 - 11.0	300	25.5	14.5 - 36.5	800
Black	5.5 4	.7 - 6.3	800	6.1	2.9 - 9.3	200	4.1 2.0 - 6	2 100	7.7	5.4 - 10.1	400	2.0 *	* 0.7 - 3.2	<100
Born outside the U.S.	30.5 25	.2 - 35.7	4,700	50.1	37.1 - 63.1	1,900	45.7 32.7 - 58	7 1,500	12.7	5.9 - 19.4	600	18.3	11.1 - 25.5	600
Speaks non-English language at home	30.4 25	.1 - 35.8	4,600	52.1	39.0 - 65.1	1,900	41.9 28.9 - 55	0 1,400	16.1	7.8 - 24.5	800	14.0	7.2 - 20.8	400
Lives in a high poverty neighborhood	74.0 69	9.4 - 78.6	11,100				87.0 * 79.1 - 95	0 2,800	65.6	56.1 - 75.2	3,200	44.5	32.7 - 56.3	1,300
Income as a percent of the Federal Poverty Guideline														
0-100%	58.7 53	8.4 - 64.0	8,600	89.3	* 79.6 - 98.9	3,100	76.5 65.9 - 87	2 2,300	50.3	40.3 - 60.3	2,500	17.0*	* 6.1 - 27.9	500
101-200%	19.0 14	.0 - 24.0	2,800	9.7	* 0.2 - 19.2	300	21.9 11.3 - 32	5 700	30.0	19.8 - 40.2	1,500	9.8*	* 3.3 - 16.3	300
> 200%	22.2 18	8.7 - 25.8	3,300						19.7	12.9 - 26.4	1,000	73.2	61.7 - 84.7	2,200

-- Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

* Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

+ Measure/definition may not be comparable to previous Snapshots (see Technical Notes).

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Fresno County was 562. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.



MIHA Data Snapshot, Fresno County by Geographical Area, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		Fresno County				Urban			Rural/Frontier	
	%	95% CI	Ν	-	%	95% CI	Ν	%	95% CI	Ν
Total	100		15,400		61.8	56.2 - 67.5	9,300	38.2	32.5 - 43.8	5,700
Health Status before Pregnancy										
In good to excellent health	89.1	85.2 - 93.0	13,700		86.4	80.8 - 92.1	8,000	93.9 *	89.4 - 98.4	5,400
Chronic conditions										
Diabetes ⁺	3.0 *	0.5 - 5.4	500							
Hypertension [†]	2.9 *	0.9 - 4.9	400		4.3 *	1.1 - 7.6	400			
Asthma†	13.3	9.0 - 17.6	2,000		15.0	9.2 - 20.8	1,400	11.2 *	4.3 - 18.1	600
Nutrition and Weight	a= 1									
Daily folic acid use, month before pregnancy	27.1	22.0 - 32.2	4,200		26.9	20.2 - 33.5	2,500	27.3	19.2 - 35.5	1,600
Overweight before pregnancy	28.0	22.6 - 33.3	4,000		25.1	18.6 - 31.7	2,200	32.8	23.5 - 42.1	1,700
Obese before pregnancy	24.2	19.0 - 29.3	3,500		25.3	18.5 - 32.0	2,200	23.1	14.8 - 31.4	1,200
Inadequate weight gain during programou	16.6	11.9 - 21.2	2,000		17.9	11.3 - 24.4	1,300	13.0	7.4 - 18.7	600
Excessive weight gain during pregnancy	43.1	36.9 - 49.4 12.1 - 20.2	5,200 2,500		41.2	32.9 - 49.6 13.8 - 25.6	3,100 1,800	47.5	37.5 - 57.6 4.6 - 14.0	2,100 500
Food insecurity during pregnancy	16.1	12.1 - 20.2	2,500		19.7	13.8 - 25.6	1,800	9.3	4.6 - 14.0	500
Intimate Partner Violence (IPV) and Depressive Symptoms		4.0 44.0	4 200		0.2	4.6 44.0	000	F 0 *	4.0 0.7	200
Physical or psychological IPV during pregnancy Propatal depressive symptoms	8.0	4.8 - 11.2	1,200		9.3	4.6 - 14.0	900	5.8 *	1.9 - 9.7	300
Prenatal depressive symptoms Postpartum depressive symptoms	14.1	9.9 - 18.2	2,200		15.1	9.5 - 20.6	1,400	13.0	6.5 - 19.6	700
	12.2	8.5 - 16.0	1,900		13.6	8.4 - 18.7	1,300	10.7	5.2 - 16.2	600
Hardships and Support during Pregnancy	20.0				24.2			22.0		
Experienced two or more hardships during childhood	29.8	24.3 - 35.2	4,400		34.3	26.9 - 41.8	3,100	22.0	14.0 - 29.9	1,200
Homeless or did not have a regular place to sleep	2.5 *	0.7 - 4.3	400		3.8 *	0.8 - 6.7	300			
Moved due to problems paying rent or mortgage Woman or partner lost job	6.0 12.1	3.1 - 8.9 8.5 - 15.7	900		7.5	3.4 - 11.6 8.1 - 18.2	700		5.0 14.0	c00
Woman or partner had pay or hours cut back	12.1 11.8		1,800 1,800		13.1 13.8	8.1 - 18.2 8.2 - 19.4	1,200 1,300	9.9 9.4 *	5.0 - 14.9 3.6 - 15.2	600 500
Became separated or divorced	11.8	7.8 - 15.9 6.1 - 14.2	1,800		13.8	6.7 - 19.4	1,300	9.4 6.9 *	1.3 - 12.5	400
Had no practical or emotional support	3.1 *	0.9 - 5.3	500		3.7 *	0.7 - 18.2	300	0.9	1.5 - 12.5	400
Substance Use	5.1	0.5 5.5	500		5.7	0.5 0.0	500			
Any smoking, 3 months before pregnancy	13.7	9.3 - 18.2	2,100		17.9	11.6 - 24.3	1,700	7.4 *	1.9 - 12.9	400
Any smoking, 3rd trimester [†]	3.7 *	1.5 - 5.8	600		5.4 *	1.9 - 8.8	500	7.4	1.9 - 12.9	400
Any smoking, postpartum	6.0	3.0 - 8.9	900		8.6	3.9 - 13.4	800			
Any binge drinking, 3 months before pregnancy	15.3	10.9 - 19.7	2,300		17.7	11.5 - 23.9	1,600	12.2	6.0 - 18.3	700
Any alcohol use, 3rd trimester [†]	3.1	1.7 - 4.5	500		2.9	1.2 - 4.5	300	3.6 *	0.9 - 6.4	200
Pregnancy Intention and Family Planning	012				2.0			0.0		
Mistimed or unwanted pregnancy	27.0	21.9 - 32.1	4,100		31.4	24.2 - 38.5	2,900	20.9	13.6 - 28.2	1,200
Unsure of pregnancy intentions	18.9	14.0 - 23.7	2,900		18.2	12.0 - 24.3	1,700	19.5	11.4 - 27.6	1,100
Postpartum birth control use [†]	82.1	77.4 - 86.7	12,400		78.3	71.6 - 85.0	7,100	88.1	82.3 - 93.8	5,000
Infant Sleep and Breastfeeding										
Placed infant on back to sleep	80.2	75.6 - 84.8	12,200		79.9	74.2 - 85.7	7,400	79.8	71.7 - 88.0	4,500
Infant always or often shared bed	41.5	35.8 - 47.2	6,300		42.4	34.8 - 50.0	3,900	41.3	32.0 - 50.5	2,300
Intended to breastfeed, before birth	89.5	86.0 - 93.0	13,500		90.2	86.0 - 94.5	8,300	87.9	81.5 - 94.2	4,900
Intended to breastfeed exclusively, before birth	52.2	46.4 - 58.0	7,900		50.0	42.4 - 57.6	4,600	55.7	46.2 - 65.1	3,100
Breastfeeding practices					-					
Any breastfeeding, 1 month after delivery	76.3	71.5 - 81.1	11,400		76.0	69.7 - 82.2	7,000	76.8	68.9 - 84.7	4,200
Exclusive breastfeeding, 1 month after delivery	36.7	31.1 - 42.3	5,400		35.5	28.2 - 42.9	3,100	38.0	28.8 - 47.1	2,100
Any breastfeeding, 3 months after delivery	51.2	44.4 - 57.9	5,700		47.1	38.3 - 55.9	3,200	55.8	45.0 - 66.6	2,300
Exclusive breastfeeding, 3 months after delivery	18.5	14.1 - 22.9	2,000		15.4	10.0 - 20.7	1,000	22.6	14.7 - 30.4	900



MIHA Data Snapshot, Fresno County by Geographical Area, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		Fresno County			Urban			Rural/Frontier	
	%	95% CI	N	%	95% CI	N	%	95% CI	N
Service Utilization and Health Insurance Coverage									
Had a usual source of pre-pregnancy care	66.9	61.5 - 72.4	10,300	60.2	52.7 - 67.7	5,600	78.0	70.6 - 85.4	4,500
Initiated prenatal care in 1st trimester	86.4	82.3 - 90.6	13,100	86.3	81.1 - 91.5	7,900	86.8	79.7 - 94.0	4,900
Mom or infant needed but couldn't afford care postpartum	7.7	4.7 - 10.8	1,200	7.5	3.9 - 11.1	700	8.3 *	2.6 - 14.1	500
Participated in WIC during pregnancy [†]	69.5	65.4 - 73.6	10,500	71.6	65.7 - 77.5	6,600	64.8	56.8 - 72.8	3,600
Received CalFresh (food stamps) during pregnancy	47.5	41.9 - 53.0	7,200	50.5	43.0 - 58.0	4,600	42.7	33.5 - 51.9	2,400
Pre-pregnancy insurance									
Medi-Cal	46.9	41.4 - 52.4	7,200	50.1	42.7 - 57.5	4,600	39.7	30.7 - 48.8	2,300
Uninsured	22.3	17.5 - 27.0	3,400	21.5	15.4 - 27.6	2,000	24.3	16.0 - 32.5	1,400
Private	28.9	24.8 - 33.0	4,400	26.3	20.8 - 31.9	2,400	34.3	26.5 - 42.0	2,000
Prenatal insurance									
Medi-Cal	72.2	68.5 - 75.9	11,000	75.1	70.1 - 80.1	7,000	66.5	58.8 - 74.3	3,700
Uninsured	0.7 *	0.0 - 1.4	100						
Private	25.0	21.4 - 28.5	3,800	21.3	16.7 - 25.8	2,000	31.9	24.3 - 39.6	1,800
Maternal postpartum insurance									
Medi-Cal	60.8	55.8 - 65.8	9,300	63.8	57.1 - 70.5	5,800	54.3	45.5 - 63.1	3,100
Uninsured	10.6	6.9 - 14.3	1,600	9.9	5.4 - 14.4	900	12.5	5.8 - 19.2	700
Private	26.6	22.8 - 30.4	4,100	23.8	18.6 - 29.0	2,200	32.0	24.5 - 39.6	1,800
Infant health insurance									
Medi-Cal	71.7	67.7 - 75.7	10,800	75.8	70.6 - 81.0	6,900	65.2	57.3 - 73.2	3,700
Uninsured	1.9 *	0.6 - 3.2	300	1.6	0.3 - 2.8	100			
Private	25.0	21.3 - 28.7	3,800	21.2	16.4 - 25.9	1,900	31.0	23.5 - 38.5	1,700
Maternal Demographics									
Age									
15-19	8.3	5.1 - 11.5	1,300	8.3	4.2 - 12.4	800	7.6 *	2.4 - 12.9	400
20-34	78.3	73.3 - 83.3	12,000	79.7	73.6 - 85.8	7,400	76.3	67.7 - 85.0	4,400
35+	13.4	9.2 - 17.6	2,100	12.0	7.1 - 16.9	1,100	16.0	8.2 - 23.9	900
Did not complete high school (or GED)	24.9	19.6 - 30.2	3,700	26.8	19.5 - 34.1	2,400	20.8	12.9 - 28.8	1,200
Unmarried	51.8	46.3 - 57.3	7,900	55.6	48.3 - 63.0	5,100	42.9	33.6 - 52.1	2,500
Race/Ethnicity									
White	19.9	15.9 - 23.9	3,000	20.7	15.0 - 26.4	1,900	19.4	13.3 - 25.5	1,100
Hispanic	61.7	56.6 - 66.8	9,400	55.7	48.5 - 62.9	5,100	71.4	64.0 - 78.9	4,000
Asian/Pacific Islander	12.1	8.0 - 16.2	1,800	14.9	9.1 - 20.7	1,400	7.1 *	2.1 - 12.0	400
Black	5.5	4.7 - 6.3	800	7.7	6.1 - 9.3	700	1.6 *	0.6 - 2.6	<100
Born outside the U.S.	30.5	25.2 - 35.7	4,700	29.5	22.4 - 36.5	2,700	31.0	22.7 - 39.3	1,800
Speaks non-English language at home	30.4	25.1 - 35.8	4,600	28.3	21.1 - 35.6	2,600	33.8	25.1 - 42.5	1,900
Lives in a high poverty neighborhood	74.0	69.4 - 78.6	11,100	78.4	72.8 - 84.1	7,300	66.8	58.4 - 75.1	3,800
Income as a percent of the Federal Poverty Guideline									
0-100%	58.7	53.4 - 64.0	8,600	62.7	55.7 - 69.8	5,800	51.0	41.6 - 60.5	2,600
101-200%	19.0	14.0 - 24.0	2,800	19.3	12.7 - 25.9	1,800	19.5	11.3 - 27.7	1,000
> 200%	22.2	18.7 - 25.8	3,300	18.0	13.7 - 22.3	1,600	29.5	21.6 - 37.3	1,500

-- Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

* Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Fresno County was 562. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. Urban and Rural/Frontier designations are based on the population size or densities of Medical Service Study Areas. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.



MIHA Data Snapshot, Fresno County by Income, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Fresno County	,	0-100% Federal Poverty	Guideline	101-200% Federal Poverty	Guideline	> 200% Federal Poverty	Guideline
	% 95% CI	Ν	% 95% CI	Ν	% 95% CI	Ν	% 95% CI	Ν
Total	100	15,400	58.7 53.4 - 64.0	8,600	19.0 14.0 - 24.0	2,800	22.2 18.7 - 25.8	3,300
Health Status before Pregnancy								
In good to excellent health	89.1 85.2 - 93.0	13,700	86.4 80.8 - 92.1	7,500			96.8 * 94.2 - 99.4	3,200
Chronic conditions								
Diabetes [†]	3.0 [*] 0.5 - 5.4	500						
Hypertension ⁺	2.9 * 0.9 - 4.9	400	4.2 * 0.8 - 7.7	400				
Asthma†	13.3 9.0 - 17.6	2,000	12.1 6.4 - 17.9	1,000	16.1 * 3.4 - 28.7	500	16.0 8.2 - 23.8	500
Nutrition and Weight	274 22 0 22 2	4 200	40 5 40 4 25 6	4 700	20 7 44 7 49 7	000	46 7 26 4 57 2	4 500
Daily folic acid use, month before pregnancy Overweight before pregnancy	27.1 22.0 - 32.2	4,200 4,000	19.5 13.4 - 25.6 26.9 19.6 - 34.3	1,700 2,100	28.7 14.7 - 42.7 33.0 18.8 - 47.2	800 900	46.7 36.4 - 57.0	1,500 900
Obese before pregnancy	28.0 22.6 - 33.3 24.2 19.0 - 29.3	4,000 3,500	20.9 19.6 - 34.3 29.9 22.2 - 37.6 - 37.6 - - - - - - - - -	2,100	$23.0 \ 18.8 \ - \ 47.2 \ 23.9 \ 11.9 \ - \ 36.0 \$	900 700	27.8 19.1 - 36.6 12.8 6.0 - 19.6	900 400
Inadequate weight gain during pregnancy	16.6 11.9 - 21.2	2,000	17.8 11.0 - 24.6	1,200	16.1 * 4.4 - 27.9	300	13.2 6.7 - 19.8	400
Excessive weight gain during pregnancy	43.1 36.9 - 49.4	5,200	37.7 28.9 - 46.5	2,600	44.2 28.8 - 59.6	1,000	53.9 42.9 - 64.8	1,500
Food insecurity during pregnancy	16.1 12.1 - 20.2	2,500	21.9 15.9 - 27.9	1,900	16.5 * 4.3 - 28.7	400	2.4 * 0.4 - 4.4	<100
Intimate Partner Violence (IPV) and Depressive Symptoms	10.1	,		,	2010			
Physical or psychological IPV during pregnancy	8.0 4.8 - 11.2	1,200	10.0 5.0 - 15.0	900			4.8 * 0.5 - 9.0	200
Prenatal depressive symptoms	14.1 9.9 - 18.2	2,200	18.4 12.0 - 24.9	1,600	7.2 * 0.5 - 13.9	200	8.6 * 3.1 - 14.1	300
Postpartum depressive symptoms	12.2 8.5 - 16.0	1,900	13.8 8.3 - 19.3	1,200	11.3 * 3.0 - 19.7	300	11.3 * 4.4 - 18.2	400
Hardships and Support during Pregnancy								
Experienced two or more hardships during childhood	29.8 24.3 - 35.2	4,400	36.2 28.4 - 44.0	3,000	29.7 15.3 - 44.1	800	17.1 9.3 - 24.9	600
Homeless or did not have a regular place to sleep	2.5 * 0.7 - 4.3	400	2.2 * 0.9 - 3.5	200				
Moved due to problems paying rent or mortgage	6.0 3.1 - 8.9	900	10.0 5.1 - 15.0	900				
Woman or partner lost job	12.1 8.5 - 15.7	1,800	15.3 10.1 - 20.4	1,300	12.8 * 2.1 - 23.5	300	3.2 [*] 0.9 - 5.5	100
Woman or partner had pay or hours cut back	11.8 7.8 - 15.9	1,800	12.3 6.9 - 17.7	1,100	18.3 * 5.3 - 31.3	500	5.6 * 1.7 - 9.6	200
Became separated or divorced	10.2 6.1 - 14.2	1,500	11.1 5.8 - 16.5	900	18.6 [*] 4.9 - 32.3	500		
Had no practical or emotional support	3.1 * 0.9 - 5.3	500	4.7 * 1.1 - 8.3	400				
Substance Use								
Any smoking, 3 months before pregnancy	13.7 9.3 - 18.2	2,100	15.9 9.8 - 21.9	1,400	19.3 * 5.1 - 33.5	500	5.6 * 2.2 - 8.9	200
Any smoking, 3rd trimester ⁺	3.7 * 1.5 - 5.8	600	5.4 * 1.8 - 8.9	500				100
Any smoking, postpartum Any binge drinking, 3 months before pregnancy	6.0 3.0 - 8.9 15.3 10.9 - 19.7	900	7.1 3.1 - 11.2 11.9 6.3 - 17.5	600	 25.6 11.7 - 39.6	700	1.6 * 0.2 - 3.0 18.7 11.4 - 26.1	<100 600
Any alcohol use, 3rd trimester [†]	3.1 1.7 - 4.5	2,300 500	11.9 0.3 - 17.3 $1.9^* 0.6 - 3.3$	1,000 200	25.0 II.7 - 39.0 	700	18.7 11.4 - 26.1 8.9* 3.4 - 14.3	300
Pregnancy Intention and Family Planning	5.1 1.7 4.5	500	1.9 0.0 5.5	200			0.5 5.4 14.5	500
Mistimed or unwanted pregnancy	27 .0 21.9 - 32.1	4,100	35.1 27.5 - 42.8	3,000	24.0 11.4 - 36.7	700	10.0 5.1 - 14.9	300
Unsure of pregnancy intentions	18.9 14.0 - 23.7	2,900	21.5 14.6 - 28.5	3,000 1,800	24.0 11.4 - 30.7 21.4 * 8.1 - 34.7	600	5.0^{*} 1.4 - 8.7	200
Postpartum birth control use [†]	82.1 77.4 - 86.7	12,400	80.5 73.8 - 87.1	6,800	80.4 * 67.5 - 93.3	2,200	85.7 79.6 - 91.7	2,800
Infant Sleep and Breastfeeding	02.11	,	0010 1010 0112	-,		_,	0017 1010 011	_,
Placed infant on back to sleep	80.2 75.6 - 84.8	12,200	78.8 72.1 - 85.5	6,800	84.8 * 73.8 - 95.9	2,400	83.2 77.0 - 89.4	2,700
Infant always or often shared bed	41.5 35.8 - 47.2	6,300	43.9 36.0 - 51.8	3,800	45.1 29.9 - 60.3	1,300	30.1 20.4 - 39.8	1,000
Intended to breastfeed, before birth	89.5 86.0 - 93.0	13,500	85.9 80.4 - 91.4	7,300	94.7 * 90.5 - 98.9	2,600		
Intended to breastfeed exclusively, before birth	52.2 46.4 - 58.0	7,900	44.0 35.9 - 52.0	3,700	53.3 38.5 - 68.1	1,500	74.5 64.3 - 84.7	2,400
Breastfeeding practices								
Any breastfeeding, 1 month after delivery	76.3 71.5 - 81.1	11,400	71.1 64.0 - 78.2	6,000	77.3 66.1 - 88.4	2,100	91.3 * 84.9 - 97.7	2,900
Exclusive breastfeeding, 1 month after delivery	36.7 31.1 - 42.3	5,400	34.3 26.3 - 42.2	2,800	32.8 19.3 - 46.4	900	46.1 35.8 - 56.4	1,500
Any breastfeeding, 3 months after delivery	51.2 44.4 - 57.9	5,700	44.0 34.7 - 53.4	2,700	45.7 28.2 - 63.3	900	73.5 62.3 - 84.7	1,800
Exclusive breastfeeding, 3 months after delivery	18.5 14.1 - 22.9	2,000	12.4 6.8 - 18.1	700	12.1 * 4.7 - 19.5	300	37.9 27.0 - 48.7	900



MIHA Data Snapshot, Fresno County by Income, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Fresno County	/	0-100% Federal Poverty	Guideline	101-200% Federal Povert	y Guideline	> 200% Federal Poverty (Guideline
	% 95% CI	Ν	% 95% CI	Ν	% 95% CI	N	% 95% CI	Ν
Service Utilization and Health Insurance Coverage								
Had a usual source of pre-pregnancy care	66.9 61.5 - 72.4	10,300	61.5 53.8 - 69.2	5,300	63.6 48.8 - 78.4	1,800	81.1 73.2 - 89.0	2,700
Initiated prenatal care in 1st trimester	86.4 82.3 - 90.6	13,100	81.7 75.6 - 87.8	7,000	85.4 * 73.9 - 96.8	2,400		
Mom or infant needed but couldn't afford care postpartum	7.7 4.7 - 10.8	1,200	9.4 4.7 - 14.1	800	10.2 * 2.9 - 17.6	300		
Participated in WIC during pregnancy ⁺	69.5 65.4 - 73.6	10,500	91.5 86.6 - 96.4	7,900	67.5 54.1 - 80.9	1,800	8.1 * 2.2 - 14.1	300
Received CalFresh (food stamps) during pregnancy	47.5 41.9 - 53.0	7,200	68.8 61.3 - 76.2	5,900	33.5 18.0 - 49.1	900		
Pre-pregnancy insurance								
Medi-Cal	46.9 41.4 - 52.4	7,200	64.7 57.2 - 72.2	5,600	36.7 21.5 - 51.8	1,000		
Uninsured	22.3 17.5 - 27.0	3,400	28.8 21.6 - 35.9	2,500	26.1 13.7 - 38.6	700	5.1 * 0.9 - 9.2	200
Private	28.9 24.8 - 33.0	4,400	5.9 2.9 - 8.9	500	33.7 20.8 - 46.7	900	87.2 80.3 - 94.1	2,900
Prenatal insurance								
Medi-Cal	72.2 68.5 - 75.9	11,000	96.4 * 94.1 - 98.7	8,300	72.0 60.5 - 83.6	2,000	7.8 * 1.5 - 14.1	300
Uninsured	0.7 * 0.0 - 1.4	100						
Private	25.0 21.4 - 28.5	3,800	2.4 * 0.5 - 4.2	200	24.6 14.1 - 35.1	700	86.7 79.4 - 94.0	2,800
Maternal postpartum insurance								
Medi-Cal	60.8 55.8 - 65.8	9,300	85.5 79.6 - 91.4	7,300	47.0 32.0 - 62.0	1,300	7.3 * 1.4 - 13.1	200
Uninsured	10.6 6.9 - 14.3	1,600	11.8 6.1 - 17.5	1,000	18.1 8.2 - 28.1	500		
Private	26.6 22.8 - 30.4	4,100	2.0 * 0.4 - 3.6	200	31.5 18.5 - 44.5	900	88.3 81.5 - 95.1	2,900
Infant health insurance								
Medi-Cal	71.7 67.7 - 75.7	10,800	97.5 * 95.7 - 99.3	8,200	68.7 56.4 - 81.0	1,900	7.6 * 2.7 - 12.5	200
Uninsured	1.9 [*] 0.6 - 3.2	300			8.0 * 1.5 - 14.5	200		
Private	25.0 21.3 - 28.7	3,800	2.2 * 0.4 - 4.0	200	21.2 10.9 - 31.5	600	88.5 82.4 - 94.6	2,900
Maternal Demographics								
Did not complete high school (or GED)	24.9 19.6 - 30.2	3,700	37.2 29.1 - 45.3	3,100	12.3 * 0.5 - 24.1	300		
Unmarried	51.8 46.3 - 57.3	7,900	69.0 61.5 - 76.4	5,900	45.1 30.0 - 60.1	1,300	13.3 6.4 - 20.1	400
Race/Ethnicity								
White	19.9 15.9 - 23.9	3,000	8.9 4.0 - 13.7	800	21.5 9.9 - 33.1	600	50.2 39.7 - 60.6	1,600
Hispanic	61.7 56.6 - 66.8	9,400	72.3 65.5 - 79.1	6,200	66.5 52.8 - 80.1	1,800	25.1 15.4 - 34.8	800
Asian/Pacific Islander	12.1 8.0 - 16.2	1,800	10.3 5.1 - 15.5	900			21.7 11.3 - 32.0	700
Black	5.5 4.7 - 6.3	800	7.7 6.1 - 9.2	700	1.8 [*] 0.3 - 3.3	<100	2.6 * 1.0 - 4.1	<100
Born outside the U.S.	30.5 25.2 - 35.7	4,700	36.6 28.8 - 44.3	3,200	27.1 14.5 - 39.7	800	16.9 9.8 - 23.9	600
Speaks non-English language at home	30.4 25.1 - 35.8	4,600	37.9 30.1 - 45.7	3,300	28.1 13.7 - 42.4	800	10.7 5.1 - 16.2	300
Lives in a high poverty neighborhood	74.0 69.4 - 78.6	11,100	89.8 85.8 - 93.8	7,500	71.6 57.2 - 86.1	2,000	36.6 26.7 - 46.5	1,200

-- Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

* Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

⁺ Measure/definition may not be comparable to previous Snapshots (see Technical Notes).

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Fresno County was 562. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.



MIHA Data Snapshot, Fresno County by Maternal Age, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Fresno Count	ty .	15-19 Years		20-34 Years		35 Years or Olde	er
	% 95% CI	Ν	% 95% CI	Ν	% 95% CI	N	% 95% CI	Ν
Total	100	15,400	8.3 5.1 - 11.5	1,300	78.3 73.3 - 83.3	12,000	13.4 9.2 - 17.6	2,100
Health Status before Pregnancy								
In good to excellent health	89.1 85.2 - 93.0	13,700			88.8 84.3 - 93.3	10,700		
Chronic conditions								
Diabetes [†]	3.0 * 0.5 - 5.4	500			2.3 [*] 0.1 - 4.4	300		
Hypertension ⁺	2.9 * 0.9 - 4.9	400			2.4 * 0.4 - 4.5	300		
Asthma ⁺	13.3 9.0 - 17.6	2,000	6.7 * 1.5 - 12.0	<100	13.7 8.8 - 18.6	1,600	15.1 [*] 0.8 - 29.5	300
Nutrition and Weight								
Daily folic acid use, month before pregnancy	27.1 22.0 - 32.2	4,200	17.4 * 3.7 - 31.0	200	25.6 20.0 - 31.1	3,100	42.0 25.4 - 58.5	900
Overweight before pregnancy	28.0 22.6 - 33.3	4,000	20.1 * 3.4 - 36.8	200	28.4 22.4 - 34.3	3,200	31.1 14.6 - 47.6	500
Obese before pregnancy	24.2 19.0 - 29.3	3,500	17.8 * 2.6 - 33.0	200	24.5 18.7 - 30.3	2,800	26.4 * 10.7 - 42.1	500
Inadequate weight gain during pregnancy	16.6 11.9 - 21.2	2,000			16.8 11.5 - 22.1	1,700	17.1 * 6.2 - 27.9	200
Excessive weight gain during pregnancy	43.1 36.9 - 49.4	5,200	42.7 21.3 - 64.1	400	40.8 33.9 - 47.7	4,100	61.2 43.7 - 78.7	800
Food insecurity during pregnancy	16.1 12.1 - 20.2	2,500			17.7 12.9 - 22.5	2,100	9.8 * 2.2 - 17.4	200
Intimate Partner Violence (IPV) and Depressive Symptoms								
Physical or psychological IPV during pregnancy	8.0 4.8 - 11.2	1,200			8.1 4.4 - 11.9	1,000		
Prenatal depressive symptoms	14.1 9.9 - 18.2	2,200			15.8 10.9 - 20.8	1,900	9.4 * 1.8 - 16.9	200
Postpartum depressive symptoms	12.2 8.5 - 16.0	1,900			14.1 9.6 - 18.5	1,700	9.1 * 0.4 - 17.7	200
Hardships and Support during Pregnancy								
Experienced two or more hardships during childhood	29.8 24.3 - 35.2	4,400	19.0 * 3.9 - 34.0	200	32 .5 26.2 - 38.8	3,800	21.1 8.8 - 33.5	400
Homeless or did not have a regular place to sleep	2.5 * 0.7 - 4.3	400			2.2 * 0.4 - 4.0	300		
Moved due to problems paying rent or mortgage	6.0 3.1 - 8.9	900			7.3 3.6 - 11.0	900		
Woman or partner lost job	12.1 8.5 - 15.7	1,800			12.9 8.8 - 16.9	1,500	12.2 * 0.5 - 24.0	300
Woman or partner had pay or hours cut back	11.8 7.8 - 15.9	1,800			12.4 7.6 - 17.1	1,500	11.8 [*] 2.2 - 21.5	200
Became separated or divorced	10.2 6.1 - 14.2	1,500			9.4 5.1 - 13.7	1,100		
Had no practical or emotional support	3.1 * 0.9 - 5.3	500			2.8 * 0.5 - 5.2	300		
Substance Use								
Any smoking, 3 months before pregnancy	13.7 9.3 - 18.2	2,100			14.1 9.1 - 19.1	1,700	17.6 * 3.2 - 32.0	400
Any smoking, 3rd trimester [†]	3.7 * 1.5 - 5.8	600			4.2 * 1.5 - 6.9	500		
Any smoking, postpartum	6.0 3.0 - 8.9	900			7.0 3.3 - 10.7	800		
Any binge drinking, 3 months before pregnancy	15.3 10.9 - 19.7	2,300			14.2 9.4 - 18.9	1,700	26.9 11.4 - 42.4	600
Any alcohol use, 3rd trimester ⁺	3.1 1.7 - 4.5	500			2.8 1.3 - 4.4	300	6.5 * 0.6 - 12.3	100
Pregnancy Intention and Family Planning								
Mistimed or unwanted pregnancy	27.0 21.9 - 32.1	4,100	49.1 28.8 - 69.3	600	27.0 21.2 - 32.9	3,200	13.1 * 3.6 - 22.6	300
Unsure of pregnancy intentions	18.9 14.0 - 23.7	2,900	34.5 * 14.2 - 54.8	400	16.2 11.2 - 21.2	1,900	24.8 * 9.1 - 40.5	500
Postpartum birth control use ⁺	82.1 77.4 - 86.7	12,400	84.1 * 69.2 - 99.0	1,100	81.7 76.3 - 87.1	9,700	82.7 * 72.0 - 93.4	1,700
Infant Sleep and Breastfeeding								
Placed infant on back to sleep	80.2 75.6 - 84.8	12,200	66.1 46.6 - 85.6	800	81.4 76.4 - 86.3	9,700	81.6 * 67.7 - 95.6	1,700
Infant always or often shared bed	41.5 35.8 - 47.2	6,300	40.2 19.6 - 60.8	500	41.6 35.3 - 48.0	5,000	41.6 24.1 - 59.0	900
Intended to breastfeed, before birth	89.5 86.0 - 93.0	13,500	79.8 * 61.0 - 98.5	900	90.2 86.6 - 93.7	10,800		
Intended to breastfeed exclusively, before birth	52.2 46.4 - 58.0	7,900	33.2 * 12.7 - 53.7	400	53.9 47.4 - 60.3	6,400	52.8 35.5 - 70.1	1,100
Breastfeeding practices		11 100		700	76 2 70 0 01 0	0.000	04.0*72.0	4.000
Any breastfeeding, 1 month after delivery	76.3 71.5 - 81.1	11,400	62.5 42.8 - 82.3	700	76.2 70.9 - 81.6	8,900	84.8 * 72.9 - 96.8	1,800
Exclusive breastfeeding, 1 month after delivery	36.7 31.1 - 42.3	5,400	27.7 * 8.8 - 46.6	300	41.1 34.6 - 47.5	4,700	16.3 [*] 6.6 - 26.0	300
Any breastfeeding, 3 months after delivery Exclusive breastfeeding, 3 months after delivery	51.2 44.4 - 57.9	5,700	20.9 * 5.0 - 36.7	200	51.9 44.3 - 59.5	4,400	66.4 48.9 - 83.8	1,100
LACIUSIVE DIEASLIEEUING, 5 MONUNS AILER UEIIVELY	18.5 14.1 - 22.9	2,000			20.9 15.5 - 26.3	1,700	13.3 * 4.1 - 22.6	200



MIHA Data Snapshot, Fresno County by Maternal Age, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Fresno Coun	ty	15-19 Years		20-34 Years		35 Years or Old	ler
	% 95% CI	Ν	% 95% CI	Ν	% 95% CI	N	% 95% CI	Ν
Service Utilization and Health Insurance Coverage								
Had a usual source of pre-pregnancy care	66.9 61.5 - 72.4	10,300	68.1 * 48.5 - 87.7	900	65.2 59.0 - 71.4	7,900	76.4 * 62.3 - 90.4	1,600
Initiated prenatal care in 1st trimester	86.4 82.3 - 90.6	13,100	86.4 * 74.3 - 98.6	1,100	86.2 81.6 - 90.7	10,300		
Mom or infant needed but couldn't afford care postpartum	7.7 4.7 - 10.8	1,200			8.4 4.7 - 12.2	1,000		
Participated in WIC during pregnancy ⁺	69.5 65.4 - 73.6	10,500			70.1 65.4 - 74.8	8,300	48.6 31.8 - 65.4	1,000
Received CalFresh (food stamps) during pregnancy	47.5 41.9 - 53.0	7,200	48.6 28.1 - 69.1	600	49.0 42.8 - 55.3	5,800	37.6 20.8 - 54.4	800
Pre-pregnancy insurance								
Medi-Cal	46.9 41.4 - 52.4	7,200	76.7 * 61.7 - 91.8	1,000	47.2 41.0 - 53.4	5,700	26.8 * 10.3 - 43.2	600
Uninsured	22.3 17.5 - 27.0	3,400			23.6 18.2 - 29.0	2,800	21.7 * 6.9 - 36.5	400
Private	28.9 24.8 - 33.0	4,400	12.7 * 2.2 - 23.1	200	27.2 22.6 - 31.7	3,300	49.2 32.6 - 65.7	1,000
Prenatal insurance								
Medi-Cal	72.2 68.5 - 75.9	11,000			72.6 68.3 - 77.0	8,700	55.7 39.8 - 71.7	1,200
Uninsured	0.7 * 0.0 - 1.4	100						
Private	25.0 21.4 - 28.5	3,800			24.3 20.2 - 28.3	2,900	41.7 26.0 - 57.3	900
Maternal postpartum insurance								
Medi-Cal	60.8 55.8 - 65.8	9,300			61.8 56.2 - 67.4	7,400	38.1 21.0 - 55.2	800
Uninsured	10.6 6.9 - 14.3	1,600			9.8 5.8 - 13.7	1,200	16.3 * 2.8 - 29.9	300
Private	26.6 22.8 - 30.4	4,100			25.9 21.5 - 30.4	3,100	45.6 29.4 - 61.7	900
Infant health insurance								
Medi-Cal	71.7 67.7 - 75.7	10,800			72.1 67.5 - 76.7	8,600	53.3 36.8 - 69.9	1,100
Uninsured	1.9 [*] 0.6 - 3.2	300			2.2 * 0.6 - 3.7	300		
Private	25.0 21.3 - 28.7	3,800			24.0 19.8 - 28.3	2,900	45.0 28.7 - 61.3	900
Maternal Demographics								
Did not complete high school (or GED)	24.9 19.6 - 30.2	3,700	50.8 30.3 - 71.4	600	22.7 16.8 - 28.6	2,700	21.6 * 7.7 - 35.5	400
Unmarried	51.8 46.3 - 57.3	7,900	84.5 * 70.5 - 98.5	1,100	52.0 45.8 - 58.2	6,300	30.3 14.6 - 46.0	600
Race/Ethnicity								
White	19.9 15.9 - 23.9	3,000			20.8 16.0 - 25.5	2,500	21.9 10.1 - 33.7	400
Hispanic	61.7 56.6 - 66.8	9,400	77.3 64.1 - 90.5	1,000	60.4 54.4 - 66.3	7,200	59.7 44.0 - 75.4	1,200
Asian/Pacific Islander	12.1 8.0 - 16.2	1,800			12.7 7.9 - 17.5	1,500	13.3 * 2.4 - 24.1	300
Black	5.5 4.7 - 6.3	800	8.3 * 2.2 - 14.3	100	5.4 4.4 - 6.5	600	4.3 * 1.2 - 7.4	<100
Born outside the U.S.	30.5 25.2 - 35.7	4,700			29.1 23.3 - 34.9	3,500	50.5 33.8 - 67.3	1,000
Speaks non-English language at home	30.4 25.1 - 35.8	4,600			31.0 25.0 - 37.0	3,700	37.3 20.6 - 54.0	800
Lives in a high poverty neighborhood	74.0 69.4 - 78.6	11,100	92.0 * 85.1 - 98.9	1,100	74.9 69.9 - 79.8	8,800	58.1 40.8 - 75.3	1,200

- - Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

* Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Fresno County was 562. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.



MIHA Data Snapshot, Fresno County by Prenatal Health Insurance, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		Fresno County			Medi-Cal				Private	e	
	%	95% CI	Ν	%	95% CI	Ν	%		95% CI		Ν
Total	100		15,400	72.2	68.5 - 75.9	11,000	25.0	0 21	l.4 - 2	28.5	3,800
Health Status before Pregnancy											
In good to excellent health Chronic conditions	89.1	85.2 - 93.0	13,700	86.6	81.4 - 91.8	9,600	97.3	2*94	1.9 - 9	99.5	3,700
Diabetes [†]	3.0 *	0.5 - 5.4	500	3.7 *	0.3 - 7.0	400					
Hypertension [†]	2.9 *	0.9 - 4.9	400	3.5 *	0.8 - 6.3	400	1.			2.3	<100
Asthma ⁺	13.3	9.0 - 17.6	2,000	13.0	7.6 - 18.4	1,400	15.	7 8	3.3 - 2	23.1	600
Nutrition and Weight	27.4			24.4							1 699
Daily folic acid use, month before pregnancy	27.1	22.0 - 32.2	4,200	21.1	15.2 - 27.0	2,300	41.			51.0	1,600
Overweight before pregnancy	28.0	22.6 - 33.3	4,000	29.8	22.8 - 36.8	3,000	22.0			29.5 23.7	900
Obese before pregnancy Inadequate weight gain during pregnancy	24.2 16.6	19.0 - 29.3 11.9 - 21.2	3,500 2,000	27.5 18.8	20.8 - 34.3 12.5 - 25.1	2,800 1,600	16. 12.			23.7 18.0	600 400
Excessive weight gain during pregnancy	43.1	36.9 - 49.4	2,000 5,200	18.8 37.1	29.3 - 45.0	3,100	12. 56.4			56.0	400 1,900
Food insecurity during pregnancy	45.1 16.1	12.1 - 20.2	2,500	21.3	15.7 - 26.8	2,300	2.3			4.4	<100
	10.1	12.1 - 20.2	2,300	21.5	15.7 - 20.8	2,300	۷.,	5 (.2 -	4.4	<100
Intimate Partner Violence (IPV) and Depressive Symptoms Physical or psychological IPV during pregnancy	8.0	4.8 - 11.2	1,200	8.8	4.7 - 13.0	1,000	5.	5 * 1	L.O - 1	LO.0	200
Prenatal depressive symptoms	8.0 14.1	9.9 - 18.2	2,200	0.0 16.1	4.7 - 13.0 10.7 - 21.5	1,000	9.0			LU.U L4.4	300
Postpartum depressive symptoms	14.1	8.5 - 16.0	1,900	10.1	8.0 - 17.4	1,800	11.			14.4	400
Hardships and Support during Pregnancy	12.2	0.5 10.0	1,500	12.7	0.0 17.4	1,400	11.	/ -		10.1	400
Experienced two or more hardships during childhood	29.8	24.3 - 35.2	4,400	34.7	27.7 - 41.8	3,700	18.	7 11	1.2 - 2	26.1	700
Homeless or did not have a regular place to sleep	29.8	0.7 - 4.3	4,400	3.4 *	0.9 - 5.9	400	10.		1.2 - 2	.0.1	700
Moved due to problems paying rent or mortgage	6.0	3.1 - 8.9	900	3.4 7.4	3.5 - 11.3	800					
Woman or partner lost job	12.1	8.5 - 15.7	1,800	14.6	9.8 - 19.4	1,600	3.3		L.O -	5.6	100
Woman or partner had pay or hours cut back	11.8	7.8 - 15.9	1,800	13.3	8.1 - 18.5	1,500	4.0			7.5	200
Became separated or divorced	10.2	6.1 - 14.2	1,500	13.2	7.7 - 18.7	1,400					
Had no practical or emotional support	3.1 *	0.9 - 5.3	500	4.3 *	1.2 - 7.3	500		-			
Substance Use											
Any smoking, 3 months before pregnancy	13.7	9.3 - 18.2	2,100	16.6	10.7 - 22.5	1,800	6.8	8 3	3.3 - 1	10.2	300
Any smoking, 3rd trimester ⁺	3.7 *	1.5 - 5.8	600	4.5 *	1.6 - 7.4	500		-			
Any smoking, postpartum	6.0	3.0 - 8.9	900	7.5	3.4 - 11.5	800	1.9	9* 0).1 -	3.7	<100
Any binge drinking, 3 months before pregnancy	15.3	10.9 - 19.7	2,300	13.2	7.8 - 18.5	1,400	21.4	4 13	3.4 - 2	29.4	800
Any alcohol use, 3rd trimester [†]	3.1	1.7 - 4.5	500	1.3 *	0.3 - 2.3	100	8.3	3 3	3.5 - 1	L3.1	300
Pregnancy Intention and Family Planning											
Mistimed or unwanted pregnancy	27.0	21.9 - 32.1	4,100	32.1	25.3 - 38.9	3,500	14.	28	3.6 - 1	19.8	500
Unsure of pregnancy intentions	18.9	14.0 - 23.7	2,900	22.3	16.0 - 28.6	2,400	10.	5* 3	3.9 - 1	17.1	400
Postpartum birth control use [†]	82.1	77.4 - 86.7	12,400	80.7	74.7 - 86.7	8,700	86.4	4 80).9 - 9	91.9	3,300
Infant Sleep and Breastfeeding											
Placed infant on back to sleep	80.2	75.6 - 84.8	12,200	79.4	73.4 - 85.4	8,700	85.	1 79	9.8 - 9	90.5	3,200
Infant always or often shared bed	41.5	35.8 - 47.2	6,300	46.4	39.2 - 53.6	5,100	29.9			38.8	1,100
Intended to breastfeed, before birth	89.5	86.0 - 93.0	13,500	87.0	82.5 - 91.6	9,500	95.3			99.9	3,600
Intended to breastfeed exclusively, before birth	52.2	46.4 - 58.0	7,900	45.0	37.7 - 52.3	4,900	71.	3 61	L.6 - 8	31.0	2,700
Breastfeeding practices	76.0			74.0				• •	-		
Any breastfeeding, 1 month after delivery	76.3	71.5 - 81.1	11,400	71.9	65.6 - 78.2	7,700	88.			94.5	3,300
Exclusive breastfeeding, 1 month after delivery	36.7	31.1 - 42.3	5,400	32.6	25.6 - 39.6	3,400	47.:			56.6	1,700
Any breastfeeding, 3 months after delivery	51.2	44.4 - 57.9	5,700	45.7	37.2 - 54.2	3,600	67.4			78.1	1,900
Exclusive breastfeeding, 3 months after delivery	18.5	14.1 - 22.9	2,000	12.4	7.5 - 17.4	1,000	34.3	Z 24	1.8 - 4	13.6	900



MIHA Data Snapshot, Fresno County by Prenatal Health Insurance, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		Fresno County			Medi-Cal			Private	
	%	95% CI	N	%	95% CI	N	%	95% CI	Ν
Service Utilization and Health Insurance Coverage									
Had a usual source of pre-pregnancy care	66.9	61.5 - 72.4	10,300	61.4	54.4 - 68.3	6,800	85.2	78.5 - 91.9	3,300
Initiated prenatal care in 1st trimester	86.4	82.3 - 90.6	13,100	83.0	77.6 - 88.5	9,000			
Mom or infant needed but couldn't afford care postpartum	7.7	4.7 - 10.8	1,200	9.5	5.3 - 13.7	1,000	3.3 *	0.8 - 5.9	100
Participated in WIC during pregnancy [†]	69.5	65.4 - 73.6	10,500	89.8	85.3 - 94.3	9,900	11.8	6.1 - 17.4	400
Received CalFresh (food stamps) during pregnancy	47.5	41.9 - 53.0	7,200	64.3	57.4 - 71.3	7,100			
Pre-pregnancy insurance									
Medi-Cal	46.9	41.4 - 52.4	7,200	63.8	57.0 - 70.6	7,000			
Uninsured	22.3	17.5 - 27.0	3,400	28.4	22.1 - 34.7	3,100			
Private	28.9	24.8 - 33.0	4,400	7.5	3.9 - 11.1	800	95.0 *	90.0 - 99.9	3,600
Maternal postpartum insurance									
Medi-Cal	60.8	55.8 - 65.8	9,300	82.3	76.6 - 87.9	9,000			
Uninsured	10.6	6.9 - 14.3	1,600	13.9	8.8 - 18.9	1,500			
Private	26.6	22.8 - 30.4	4,100	3.4 *	0.5 - 6.3	400	95.4 *	91.8 - 99.0	3,600
Infant health insurance									
Medi-Cal	71.7	67.7 - 75.7	10,800	95.0	92.3 - 97.7	10,300	9.4 *	3.8 - 14.9	400
Uninsured	1.9 *	0.6 - 3.2	300	1.7 *	0.1 - 3.4	200			
Private	25.0	21.3 - 28.7	3,800	2.7 *	0.7 - 4.7	300	88.5	82.6 - 94.3	3,300
Maternal Demographics									
Age									
15-19	8.3	5.1 - 11.5	1,300	10.6	6.3 - 14.9	1,200			
20-34	78.3	73.3 - 83.3	12,000	79.0	72.9 - 85.1	8,700	76.4	67.5 - 85.2	2,900
35+	13.4	9.2 - 17.6	2,100	10.4	5.5 - 15.3	1,200	22.5	13.8 - 31.2	900
Did not complete high school (or GED)	24.9	19.6 - 30.2	3,700	33.9	26.7 - 41.0	3,600			
Unmarried	51.8	46.3 - 57.3	7,900	65.5	58.7 - 72.3	7,200	16.4	9.6 - 23.3	600
Race/Ethnicity									
White	19.9	15.9 - 23.9	3,000	10.6	5.8 - 15.3	1,200	48.9	39.4 - 58.4	1,800
Hispanic	61.7	56.6 - 66.8	9,400	71.9	65.7 - 78.1	7,900	32.9	23.2 - 42.6	1,200
Asian/Pacific Islander	12.1	8.0 - 16.2	1,800	10.4	5.8 - 14.9	1,100	14.7	6.4 - 23.0	500
Black	5.5	4.7 - 6.3	800	6.5	5.3 - 7.6	700	2.5	1.3 - 3.8	<100
Born outside the U.S.	30.5	25.2 - 35.7	4,700	36.3	29.4 - 43.2	4,000	14.2	8.6 - 19.9	500
Speaks non-English language at home	30.4	25.1 - 35.8	4,600	37.4	30.3 - 44.4	4,100	8.3	3.9 - 12.6	300
Lives in a high poverty neighborhood	74.0	69.4 - 78.6	11,100	87.5	82.8 - 92.3	9,400	41.7	32.4 - 51.1	1,600
Income as a percent of the Federal Poverty Guideline									
0-100%	58.7	53.4 - 64.0	8,600	78.6	72.1 - 85.1	8,300	5.5 *	1.3 - 9.7	200
101-200%	19.0	14.0 - 24.0	2,800	19.0	12.7 - 25.3	2,000	18.6	11.1 - 26.1	700
> 200%	22.2	18.7 - 25.8	3,300	2.4 *	0.4 - 4.4	300	75.9	67.8 - 84.0	2,800

-- Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

* Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%)

⁺ Measure/definition may not be comparable to previous Snapshots (see Technical Notes).

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Fresno County was 562. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.



MIHA Data Snapshot, Fresno County by Neighborhood Poverty, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Fr	esno Coun	ty		9% of F elow P				9% of Reside elow Povert			9.9% of Resid elow Povert			0% of Reside	
	%	95% CI	Ν	%	95%	CI	Ν	%	95% CI	N	%	95% CI	N	%	95% CI	N
Total	100		15,400	3.5	1.7 -	5.3	500	11.3	8.1 - 14.5	1,700	11.2	7.5 - 14.8	1,700	74.0	69.4 - 78.6	11,10
Health Status before Pregnancy																
In good to excellent health Chronic conditions	89.1	85.2 - 93.0	13,700											86.3	81.1 - 91.5	9,60
Diabetes ⁺	3.0 *	0.5 - 5.4	500											2.5	* 0.1 - 4.8	30
Hypertension ⁺	2.9 *	0.9 - 4.9	400											3.5	* 0.8 - 6.2	40
Asthma ⁺	13.3	9.0 - 17.6	2,000								20.7 *	3.8 - 37.6	300	11.9	7.0 - 16.8	1,30
Nutrition and Weight																
Daily folic acid use, month before pregnancy	27.1	22.0 - 32.2	4,200	43.8 *	17.8 -	69.8	200	46.0	30.3 - 61.6	800	34.2	17.7 - 50.7	600	22.2	16.6 - 27.9	2,50
Overweight before pregnancy	28.0	22.6 - 33.3	4,000					21.0	10.0 - 32.0	400	28.4 *	10.6 - 46.3	400	30.0	23.4 - 36.6	3,10
Obese before pregnancy	24.2	19.0 - 29.3	3,500					5.2 *	1.3 - 9.1	<100	15.6 *	4.8 - 26.4	200	29.5	22.8 - 36.2	3,10
Inadequate weight gain during pregnancy	16.6	11.9 - 21.2	2,000	29.1 *	5.0 -	53.2	100	11.4 *	4.3 - 18.6	200	11.6 *	0.4 - 22.9	200	17.0	11.1 - 22.8	1,50
Excessive weight gain during pregnancy	-	36.9 - 49.4	5,200	50.8	25.0 -	76.6	200		33.4 - 66.3	700	54.3	36.1 - 72.6	700		33.0 - 48.2	3,50
Food insecurity during pregnancy	16.1	12.1 - 20.2	2,500					6.4 *	0.7 - 12.1	100				19.0	13.7 - 24.3	2,10
Intimate Partner Violence (IPV) and Depressive Symptoms																
Physical or psychological IPV during pregnancy		4.8 - 11.2	1,200									0.2 - 4.9	<100	9.0	4.8 - 13.2	1,00
Prenatal depressive symptoms		9.9 - 18.2	2,200									0.5 - 16.5	100	-	10.4 - 21.0	1,70
Postpartum depressive symptoms	12.2	8.5 - 16.0	1,900								13.4 *	2.4 - 24.3	200	13.5	8.7 - 18.3	1,50
Hardships and Support during Pregnancy																
Experienced two or more hardships during childhood	29.8	24.3 - 35.2	4,400	38.8 *	12.7 -	64.9	200	12.5 *	3.6 - 21.4	200	30.2	15.0 - 45.4	500	31.8	25.0 - 38.6	3,40
Homeless or did not have a regular place to sleep	2.5 *	0.7 - 4.3	400											-	* 0.9 - 5.9	40
Moved due to problems paying rent or mortgage		3.1 - 8.9	900											7.5	3.6 - 11.5	80
Woman or partner lost job		8.5 - 15.7	1,800								16.8 *	1.8 - 31.9	300	12.5	8.2 - 16.7	1,40
Woman or partner had pay or hours cut back	-	7.8 - 15.9	1,800											13.9	8.7 - 19.1	1,50
Became separated or divorced	-	6.1 - 14.2	1,500											11.6	6.6 - 16.7	1,30
Had no practical or emotional support	3.1 *	0.9 - 5.3	500											4.2	* 1.2 - 7.2	50
Substance Use																
Any smoking, 3 months before pregnancy	-	9.3 - 18.2	2,100					-	0.2 - 8.5	<100					10.2 - 21.2	1,70
Any smoking, 3rd trimester ⁺	-	1.5 - 5.8	600											-	* 1.8 - 7.7	50
Any smoking, postpartum		3.0 - 8.9	900											7.9	3.9 - 12.0	90
Any binge drinking, 3 months before pregnancy		10.9 - 19.7	2,300	26.2 *	1.7 -	50.7	100		3.3 - 22.7	200	-	2.3 - 33.5	300	15.2	9.9 - 20.4	1,70
Any alcohol use, 3rd trimester [†]	3.1	1.7 - 4.5	500					6.8*	0.6 - 13.0	100	6.1*	1.1 - 11.0	100	1.4	* 0.4 - 2.4	20
Pregnancy Intention and Family Planning																
Mistimed or unwanted pregnancy	-	21.9 - 32.1	4,100					-	2.2 - 12.4	100	-	4.0 - 24.0	200		26.3 - 39.6	3,60
Unsure of pregnancy intentions		14.0 - 23.7	2,900						2.1 - 18.4	200		15.0 - 51.0	500	-	12.7 - 24.2	2,00
Postpartum birth control use [†]	82.1	77.4 - 86.7	12,400					82.5*	70.5 - 94.6	1,400	89.5*	80.4 - 98.6	1,500	80.7	74.8 - 86.6	8,80
Infant Sleep and Breastfeeding																
Placed infant on back to sleep		75.6 - 84.8	12,200		33.4 -		300		81.8 - 95.9	1,500		59.2 - 90.8	1,300		74.7 - 85.9	8,80
Infant always or often shared bed	-	35.8 - 47.2	6,300	-	8.3 -	55.9	200		12.6 - 42.7	500		44.3 - 76.6	1,000	-	34.8 - 48.7	4,60
Intended to breastfeed, before birth		86.0 - 93.0	13,500		44.0	02.0	400		CE 4 00 5	1 200		20.6 74.5	000		82.9 - 91.8	9,50
Intended to breastfeed exclusively, before birth Breastfeeding practices		46.4 - 58.0	7,900		44.0 -	93.8	400	/9.4*	65.4 - 93.4	1,300		38.6 - 74.2	900		39.5 - 53.6	5,10
Any breastfeeding, 1 month after delivery		71.5 - 81.1	11,400									72.9 - 96.6	1,400		66.2 - 78.4	7,80
Exclusive breastfeeding, 1 month after delivery		31.1 - 42.3	5,400		34.6 -	84.4	300		36.9 - 69.1	800		21.1 - 51.5	600		26.1 - 39.7	3,50
Any breastfeeding, 3 months after delivery	-	44.4 - 57.9	5,700						51.9 - 85.7	900		42.7 - 83.1	700		36.0 - 52.2	3,50
Exclusive breastfeeding, 3 months after delivery	18.5	14.1 - 22.9	2,000	23.7 *	0.7 -	46.7	<100	38.3	22.5 - 54.0	500	33.9	16.0 - 51.7	400	12.3	7.8 - 16.9	1,000



MIHA Data Snapshot, Fresno County by Neighborhood Poverty, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		esno Coun		0-4.9% of Resid		5-9.9% of Reside below Povert		10-19.9% of Resi below Pover			0% of Reside elow Povert	
	%	95% CI	N	% 95% CI	Ν	% 95% CI	Ν	% 95% CI	Ν	%	95% CI	Ν
Service Utilization and Health Insurance Coverage												
Had a usual source of pre-pregnancy care	66.9	61.5 - 72.4	10,300			69.3 54.7 - 84.0	1,200	64.7 48.8 - 80.5	1,100	66.5	59.9 - 73.1	7,400
Initiated prenatal care in 1st trimester	86.4	82.3 - 90.6	13,100					79.0 * 61.3 - 96.7	1,300	85.5	80.6 - 90.3	9,400
Mom or infant needed but couldn't afford care postpartum	7.7	4.7 - 10.8	1,200							9.6	5.4 - 13.7	1,000
Participated in WIC during pregnancy [†]	69.5	65.4 - 73.6	10,500			16.3 * 6.6 - 26.1	300	56.7 39.3 - 74.2	900	80.9	76.2 - 85.6	9,000
Received CalFresh (food stamps) during pregnancy	47.5	41.9 - 53.0	7,200			7.6 * 1.6 - 13.6	100	39.4 21.1 - 57.7	600	56.4	49.5 - 63.3	6,200
Pre-pregnancy insurance												
Medi-Cal	46.9	41.4 - 52.4	7,200			9.3 * 1.5 - 17.0	200	37.5 19.2 - 55.8	600	55.0	48.2 - 61.8	6,100
Uninsured	22.3	17.5 - 27.0	3,400			8.0 * 1.0 - 15.0	100	24.3 * 9.8 - 38.8	400	25.0	18.9 - 31.1	2,800
Private	28.9	24.8 - 33.0	4,400	78.6 * 60.3 - 97.0	400	72.5 58.7 - 86.4	1,200	35.1 20.8 - 49.3	600	19.5	14.8 - 24.2	2,200
Prenatal insurance												
Medi-Cal	72.2	68.5 - 75.9	11,000			15.3 * 5.7 - 24.8	300	57.7 41.5 - 73.8	1,000	84.9	81.3 - 88.6	9,400
Uninsured	0.7 *	0.0 - 1.4	100									
Private	25.0	21.4 - 28.5	3,800	66.0 * 41.0 - 91.0	300	73.8 59.9 - 87.6	1,200	36.9 21.7 - 52.2	600	14.3	10.7 - 17.9	1,600
Maternal postpartum insurance												
Medi-Cal	60.8	55.8 - 65.8	9,300			16.4 * 6.7 - 26.1	300	54.7 37.8 - 71.6	900	69.7	63.6 - 75.8	7,700
Uninsured	10.6	6.9 - 14.3	1,600							13.6	8.7 - 18.6	1,500
Private	26.6	22.8 - 30.4	4,100	70.5 * 48.4 - 92.6	400	72.9 59.2 - 86.5	1,200	37.7 22.1 - 53.3	600	16.2	11.9 - 20.5	1,800
Infant health insurance												
Medi-Cal		67.7 - 75.7	10,800	32.3 * 6.6 - 58.1	200	17.1 7.4 - 26.8	300	62.6 47.5 - 77.7	1,000		79.1 - 87.4	9,100
Uninsured	1.9 *	0.6 - 3.2	300							2.4 *	0.7 - 4.1	300
Private	25.0	21.3 - 28.7	3,800	59.5 * 33.6 - 85.4	300	78.5 67.1 - 89.8	1,300	34.2 19.9 - 48.4	600	13.8	10.1 - 17.6	1,500
Maternal Demographics												
Age												
15-19	8.3	5.1 - 11.5	1,300					5.2 * 0.4 - 9.9	<100	10.0	5.7 - 14.3	1,100
20-34	78.3	73.3 - 83.3	12,000			78.7 67.2 - 90.1	1,300	71.7 * 52.9 - 90.5	1,200	79.4	73.6 - 85.2	8,800
35+	13.4	9.2 - 17.6	2,100			20.7 9.3 - 32.1	400	23.2 * 3.9 - 42.4	400	10.6	6.2 - 15.0	1,200
Did not complete high school (or GED)	-	19.6 - 30.2	3,700							32.4	25.5 - 39.3	3,500
Unmarried	51.8	46.3 - 57.3	7,900	18.2 * 0.6 - 35.9	<100	14.9 * 5.5 - 24.3	300	33.0 15.7 - 50.3	600	60.5	53.8 - 67.2	6,700
Race/Ethnicity												
White		15.9 - 23.9	3,000	53.9 27.3 - 80.5		44.2 29.4 - 59.0	800	24.4 13.2 - 35.6	400	14.5	9.6 - 19.4	1,600
Hispanic	-	56.6 - 66.8	9,400	25.0 * 2.8 - 47.3	100	18.8 7.9 - 29.6	300	58.1 41.8 - 74.3	900		64.3 - 76.3	7,800
Asian/Pacific Islander	12.1	8.0 - 16.2	1,800			31.9 14.8 - 48.9	500	13.3 * 2.6 - 24.0	200	8.3	4.2 - 12.4	900
Black	5.5	4.7 - 6.3	800			3.6 * 0.4 - 6.8	<100	4.3 * 1.5 - 7.1	<100	6.0	4.8 - 7.3	700
Born outside the U.S.		25.2 - 35.7	4,700			11.0 [*] 3.1 - 18.9	200	22.6 * 7.4 - 37.9	400		28.0 - 41.2	3,800
Speaks non-English language at home	30.4	25.1 - 35.8	4,600			12.4 * 1.1 - 23.7	200	26.0 * 9.7 - 42.3	400	34.6	27.9 - 41.3	3,800
Income as a percent of the Federal Poverty Guideline	-											
0-100%		53.4 - 64.0	8,600			16.7 * 6.7 - 26.7	300	32.0 16.2 - 47.9	500		64.3 - 76.7	7,500
101-200%		14.0 - 24.0	2,800					33.2 * 12.7 - 53.6	500		13.0 - 24.3	2,000
> 200%		18.7 - 25.8	3,300	60.3 * 33.6 - 86.9		72.2 58.3 - 86.0	1,200	34.8 19.5 - 50.1	500	10.9	7.5 - 14.2	1,200

-- Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

* Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

⁺ Measure/definition may not be comparable to previous Snapshots (see Technical Notes).

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Fresno County was 562. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. Neighborhood poverty is the percentage of residents with incomes below poverty in a woman's census tract. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.



MIHA Data Snapshot, Fresno County by Race/Ethnicity, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Fresno Cou	nty	Hispanic		Black		White		Asian/Pacific Islander	
	% 95% CI	Ν	% 95% CI	Ν	% 95% CI	Ν	% 95% CI	N	% 95% CI	Ν
Total	100	15,400	61.7 56.6 - 66.8	9,400	5.5 4.7 - 6.3	800	19.9 15.9 - 23.9	3,000	12.1 8.0 - 16.2	1,800
Health Status before Pregnancy										
In good to excellent health	89.1 85.2 - 93.0	13,700	89.8 85.2 - 94.4	8,400	84.7 76.7 - 92.6	700			82.9 * 67.1 - 98.6	1,500
Chronic conditions										
Diabetes [†]	3.0 * 0.5 - 5.4	500	4.4 * 0.5 - 8.3	400						
Hypertension ⁺	2.9 * 0.9 - 4.9	400			6.4 * 1.2 - 11.6	<100	1.5 * 0.1 - 2.8	<100		
Asthma ⁺	13.3 9.0 - 17.6	2,000	10.8 5.0 - 16.5	1,000	33.3 21.4 - 45.3	300	19.5 9.0 - 29.9	600		
Nutrition and Weight										
Daily folic acid use, month before pregnancy	27.1 22.0 - 32.2	4,200	22.0 15.4 - 28.6	2,100	24.3 14.4 - 34.1	200	38.2 27.6 - 48.8	1,200	35.4 17.9 - 53.0	700
Overweight before pregnancy	28.0 22.6 - 33.3	4,000	29.2 21.8 - 36.6	2,500	27.0 15.9 - 38.2	200	34.3 22.3 - 46.2	1,000	13.8 [*] 3.2 - 24.4	200
Obese before pregnancy	24.2 19.0 - 29.3	3,500	25.7 18.4 - 33.0	2,200	34.1 22.7 - 45.5	300	18.7 9.6 - 27.9	600	18.7 * 4.3 - 33.1	300
Inadequate weight gain during pregnancy	16.6 11.9 - 21.2	2,000	15.4 9.4 - 21.4	1,100	11.8 * 4.0 - 19.5	<100	11.9 6.0 - 17.9	300	35.6 15.3 - 55.9	500
Excessive weight gain during pregnancy	43.1 36.9 - 49.4	5,200	41.8 33.1 - 50.6	3,000	51.8 39.0 - 64.6	300	47.1 35.5 - 58.6	1,200	37.0 16.9 - 57.1	500
Food insecurity during pregnancy	16.1 12.1 - 20.2	2,500	16.8 11.5 - 22.0	1,600	22.5 13.1 - 31.9	200	16.2 * 5.1 - 27.3	500	12.1 * 3.0 - 21.1	200
Intimate Partner Violence (IPV) and Depressive Symptoms										
Physical or psychological IPV during pregnancy	8.0 4.8 - 11.2	1,200	8.9 4.1 - 13.7	800	15.2 7.4 - 23.0	100				
Prenatal depressive symptoms	14.1 9.9 - 18.2		15.2 9.1 - 21.3	1,400	25.5 15.5 - 35.6	200	8.6 * 2.3 - 14.9	300	10.0 * 1.9 - 18.1	200
Postpartum depressive symptoms	12.2 8.5 - 16.0		13.2 7.8 - 18.7	1,200	12.1 * 4.9 - 19.2	100	10.2 * 3.5 - 17.0	300	10.2 * 2.0 - 18.5	200
Hardships and Support during Pregnancy		,	1012	,			1012		2012	
Experienced two or more hardships during childhood	29.8 24.3 - 35.2	4,400	32.5 25.0 - 40.1	3,000	36.6 25.3 - 48.0	300	25.9 13.7 - 38.0	700	16.0 * 5.0 - 27.0	300
Homeless or did not have a regular place to sleep	2.5 * 0.7 - 4.3			3,000	9.9 * 3.6 - 16.2	<100	50.0	700		500
Moved due to problems paying rent or mortgage	6.0 3.1 - 8.9		7.4 * 2.8 - 11.9	700	13.9 6.0 - 21.8	100				
Woman or partner lost job	12.1 8.5 - 15.7		14.1 8.7 - 19.4	1,300	22.3 12.8 - 31.9	200	6.2 * 1.1 - 11.2	200		
Woman or partner had pay or hours cut back	11.8 7.8 - 15.9		9.0 4.5 - 13.5	800	14.0 6.6 - 21.4	100	15.8 * 4.6 - 27.0	500	19.8 * 4.7 - 34.8	40
Became separated or divorced	10.2 6.1 - 14.2	,	13.7 7.3 - 20.0	1,300	24.1 13.6 - 34.6	200		500		400
Had no practical or emotional support	3.1 * 0.9 - 5.3		3.1 * 0.2 - 6.1	300		200				
Substance Use	5.1 0.5 5.5	500	J.1 0.2 0.1	500						
Any smoking, 3 months before pregnancy	13.7 9.3 - 18.2	2,100	9.8 4.8 - 14.8	900	24.1 14.4 - 33.8	200	26.3 13.9 - 38.8	800		
Any smoking, 3rd trimester ⁺	3.7 * 1.5 - 5.8	,	1.9 * 0.3 - 3.6	900 200	9.6 * 3.1 - 16.0	<100	10.0 * 1.0 - 19.1	300		
Any smoking, postpartum	6.0 3.0 - 8.9		1.9 0.3 - 3.8 2.1^* 0.3 - 3.8	200	17.3 8.8 - 25.8	100	15.3 * 3.6 - 26.9	500		
Any binge drinking, 3 months before pregnancy	15.3 10.9 - 19.7		16.4 10.1 - 22.7	1,500	7.5 * 2.0 - 13.0	<100	15.0 8.2 - 21.9	400		300
Any alcohol use, 3rd trimester [†]	3.1 1.7 - 4.5		$10.4 \ 10.1 - 22.7$ $1.2^* \ 0.1 - 2.4$	1,500	7.5 2.0 - 13.0	<100	6.8 * 2.8 - 10.9	400 200	16.4 * 2.8 - 30.0 	30
•	5.1 1.7 - 4.5	500	1.2 0.1 - 2.4	100			0.0 2.8 - 10.9	200		
Pregnancy Intention and Family Planning	27.0				F2 F 1 1 1 1 1				C 0 * 00 /05	
Mistimed or unwanted pregnancy	27.0 21.9 - 32.1		27.5 20.6 - 34.5	2,600	53.5 41.7 - 65.3	400	31.5 19.3 - 43.7	900	6.9 * 0.2 - 13.5	10
Unsure of pregnancy intentions	18.9 14.0 - 23.7		21.5 14.8 - 28.2	2,000	13.2 6.2 - 20.1	100	6.6 * 1.4 - 11.9	200	29.7 * 12.1 - 47.4	50
Postpartum birth control use [†]	82.1 77.4 - 86.7	12,400	86.2 80.4 - 92.0	8,000	69.7 58.8 - 80.5	600	85.5 78.6 - 92.4	2,500	62.5 43.8 - 81.2	1,10
Infant Sleep and Breastfeeding										
Placed infant on back to sleep	80.2 75.6 - 84.8		83.4 77.3 - 89.5	7,800	61.5 50.3 - 72.6	500	74.1 63.7 - 84.5	2,200	84.4 * 72.9 - 96.0	1,50
Infant always or often shared bed	41.5 35.8 - 47.2		45.0 37.1 - 52.8	4,200	66.2 54.2 - 78.1	500	27.3 15.9 - 38.7	800	34.2 17.1 - 51.4	60
Intended to breastfeed, before birth	89.5 86.0 - 93.0		86.9 81.6 - 92.3	8,000	75.9 64.3 - 87.4	600	95.8 * 92.6 - 99.0	2,900		
Intended to breastfeed exclusively, before birth	52.2 46.4 - 58.0	7,900	45.8 37.9 - 53.7	4,200	43.1 31.4 - 54.7	300	81.4 73.0 - 89.7	2,400	35.3 17.7 - 52.9	60
Breastfeeding practices	76.0		764 655		65 4 5 5 5		07.0		60 4 4 5 -	
Any breastfeeding, 1 month after delivery	76.3 71.5 - 81.1		76.1 69.6 - 82.7	6,900	65.4 53.7 - 77.0	500	87.6 81.1 - 94.2	2,600	62.1 44.4 - 79.9	1,10
Exclusive breastfeeding, 1 month after delivery	36.7 31.1 - 42.3		32.3 25.0 - 39.7	2,900	29.6 18.3 - 40.9	200	55.4 44.3 - 66.5	1,600	28.5 12.9 - 44.1	50
Any breastfeeding, 3 months after delivery	51.2 44.4 - 57.9		50.3 40.9 - 59.6	3,200	34.0 22.3 - 45.8	200	63.4 49.8 - 76.9	1,500	36.6 * 15.0 - 58.1	500
Exclusive breastfeeding, 3 months after delivery	18.5 14.1 - 22.9	2,000	11.9 7.1 - 16.8	800	17.0 7.6 - 26.3	100	38.0 25.8 - 50.2	900	14.0 * 0.9 - 27.1	200



MIHA Data Snapshot, Fresno County by Race/Ethnicity, 2013-2015

DPH Maternal and Infant Health Assessment (MIHA) Survey

	Fresno County			Hispanic		Black	White			Asian/Pacific Islander				
	% 9	95% CI	N	%	95% CI	N	% 95% CI	N	%	95% CI	N	%	95% CI	Ν
Service Utilization and Health Insurance Coverage														
Had a usual source of pre-pregnancy care	66.9 61.	5 - 72.4	10,300	65.5	58.2 - 72.8	6,100	70.3 59.4 - 81.	2 600	74.8	63.9 - 85.7	2,300	56.9	39.1 - 74.7	1,000
Initiated prenatal care in 1st trimester	86.4 82.	3 - 90.6	13,100	83.1	77.1 - 89.1	7,700	81.2 70.8 - 91.	5 700						
Mom or infant needed but couldn't afford care postpartum	7.7 4.	7 - 10.8	1,200	8.7	4.4 - 12.9	800			8.1	* 0.8 - 15.3	200			
Participated in WIC during pregnancy ⁺	69.5 65.	4 - 73.6	10,500	84.2	78.5 - 89.9	7,800	80.3 71.1 - 89.	5 700	31.9	19.5 - 44.2	900	56.7	39.2 - 74.1	1,000
Received CalFresh (food stamps) during pregnancy	47.5 41.	9 - 53.0	7,200	55.7	47.9 - 63.5	5,100	74.4 64.9 - 83.	9 600	24.7	12.5 - 37.0	700	34.7	17.0 - 52.3	600
Pre-pregnancy insurance														
Medi-Cal	46.9 41.	4 - 52.4	7,200	54.3	46.7 - 62.0	5,100	67.0 57.2 - 76.	8 600	23.8	11.5 - 36.1	700	42.0	24.4 - 59.7	800
Uninsured	22.3 17.	5 - 27.0	3,400	28.1	21.2 - 35.0	2,600	15.4 7.3 - 23.	5 100	12.5	* 4.0 - 21.0	400	12.9*	* 0.4 - 25.4	200
Private	28.9 24.	8 - 33.0	4,400	16.7	11.5 - 22.0	1,600	15.1 8.9 - 21.	3 100	62.4	50.2 - 74.6	1,900	37.0	20.2 - 53.8	700
Prenatal insurance														
Medi-Cal	72.2 68.	5 - 75.9	11,000	85.5	80.8 - 90.2	7,900	85.2 78.4 - 92.	0 700	38.6	26.6 - 50.7	1,200	62.0	45.0 - 79.1	1,100
Uninsured	0.7 * 0.	0 - 1.4	100											
Private	25.0 21.	4 - 28.5	3,800	12.9	8.4 - 17.4	1,200	10.9 5.8 - 16.	0 <100	59.0	47.1 - 70.8	1,800	29.1	13.5 - 44.8	50
Maternal postpartum insurance														
Medi-Cal	60.8 55.	8 - 65.8	9,300	69.8	62.7 - 76.9	6,500	86.6 80.5 - 92.	8 700	32.1	19.5 - 44.6	1,000	56.8	39.5 - 74.1	1,00
Uninsured	10.6 6.	9 - 14.3	1,600	14.6	8.9 - 20.4	1,400			7.9	* 2.3 - 13.6	200			
Private	26.6 22.	8 - 30.4	4,100	14.5	9.4 - 19.6	1,300	11.0 5.7 - 16.	3 <100	58.8	46.9 - 70.7	1,800	34.4	18.0 - 50.7	60
Infant health insurance														
Medi-Cal	71.7 67.	7 - 75.7	10,800	86.1	81.4 - 90.7	7,900	89.6 84.2 - 95.	0 700	35.3	22.8 - 47.7	1,100	55.5	37.5 - 73.5	1,00
Uninsured	1.9 [*] 0.	6 - 3.2	300						3.9	* 0.4 - 7.4	100			
Private	25.0 21.	3 - 28.7	3,800	11.6	7.4 - 15.8	1,100	8.3 3.9 - 12.	7 <100	57.9	46.0 - 69.7	1,700	41.4	23.6 - 59.3	70
Maternal Demographics														
Age														
15-19	8.3 5.	1 - 11.5	1,300	10.5	5.5 - 15.4	1,000	12.5 * 4.3 - 20.	7 100						
20-34	78.3 73.	3 - 83.3	12,000	76.5	69.4 - 83.6	7,200	76.9 66.9 - 86.	9 600	81.7	73.2 - 90.2	2,500	81.9*	* 69.2 - 94.6	1,50
35+	13.4 9.	2 - 17.6	2,100	13.0	7.1 - 19.0	1,200	10.6 * 3.8 - 17.	3 <100	14.9	7.0 - 22.8	400	14.8*	* 2.8 - 26.7	30
Did not complete high school (or GED)	24.9 19.	6 - 30.2	3,700	30.5	23.1 - 37.9	2,800	28.5 16.3 - 40.	5 200						
Unmarried	51.8 46.	3 - 57.3	7,900	59.2	51.5 - 66.9	5,500	84.2 77.1 - 91.	3 700	30.9	18.4 - 43.4	900	37.9	20.1 - 55.7	70
Born outside the U.S.	30.5 25.	2 - 35.7	4,700	37.1	29.6 - 44.5	3,500			4.9	* 0.3 - 9.5	100	55.1	37.0 - 73.2	1,00
Speaks non-English language at home	30.4 25.	1 - 35.8	4,600	39.1	31.5 - 46.7	3,700						51.5	32.8 - 70.3	90
Lives in a high poverty neighborhood	74.0 69.	4 - 78.6	11,100	85.1	79.4 - 90.8	7,800	83.4 75.4 - 91.	5 700	53.5	42.4 - 64.6	1,600		33.7 - 70.4	90
Income as a percent of the Federal Poverty Guideline														
0-100%	58.7 53.	4 - 64.0	8,600	70.4	63.0 - 77.8	6,200	83.5 76.1 - 90.	9 700	26.1	14.2 - 38.1	800	48.9	30.9 - 66.8	90
101-200%	19.0 14.	0 - 24.0	2,800	20.8	13.8 - 27.8	1,800	6.4 * 1.4 - 11.	4 <100	-	9.9 - 30.8	600	14.1 *	* 0.6 - 27.6	30
> 200%	22.2 18.		3,300		5.0 - 12.6	800	10.1 4.2 - 16.	0 <100	-	41.8 - 65.2	1,600		19.9 - 54.1	700

-- Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

* Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

⁺ Measure/definition may not be comparable to previous Snapshots (see Technical Notes).

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Fresno County was 562. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.



MIHA Data Snapshot, Fresno County by Total Live Births, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		Fresno County			First Live Birth		Second Live Birth or More		
	%	95% CI	Ν	%	95% CI	Ν	%	95% CI	Ν
Total	100		15,400	32.4	27.1 - 37.7	5,000	67.6	62.3 - 72.9	10,300
Health Status before Pregnancy									
In good to excellent health Chronic conditions	89.1	85.2 - 93.0	13,700	90.8	84.3 - 97.3	4,500	88.2	83.4 - 93.1	9,100
Diabetes ⁺	3.0 *	0.5 - 5.4	500				4.1 *	0.5 - 7.7	400
Hypertension ⁺	2.9 *	0.9 - 4.9	400				4.0 *	1.1 - 6.9	400
Asthma ⁺	13.3	9.0 - 17.6	2,000	12.7	6.6 - 18.8	600	13.7	8.0 - 19.4	1,400
Nutrition and Weight									
Daily folic acid use, month before pregnancy	27.1	22.0 - 32.2	4,200	35.4	26.3 - 44.6	1,800	23.1	17.1 - 29.0	2,400
Overweight before pregnancy	28.0	22.6 - 33.3	4,000	27.7	18.9 - 36.6	1,300	28.3	21.6 - 35.0	2,700
Obese before pregnancy	24.2	19.0 - 29.3	3,500	12.8	7.3 - 18.2	600	29.6	22.6 - 36.6	2,800
Inadequate weight gain during pregnancy	16.6	11.9 - 21.2	2,000	16.7	8.6 - 24.7	700	16.6	10.9 - 22.4	1,300
Excessive weight gain during pregnancy	43.1	36.9 - 49.4	5,200	45.5	35.6 - 55.5	1,800	41.6	33.6 - 49.7	3,400
Food insecurity during pregnancy	16.1	12.1 - 20.2	2,500	20.1	11.7 - 28.5	1,000	14.3	10.0 - 18.7	1,500
Intimate Partner Violence (IPV) and Depressive Symptoms									
Physical or psychological IPV during pregnancy	8.0	4.8 - 11.2	1,200	6.5	2.0 - 11.0	300	8.8	4.5 - 13.1	900
Prenatal depressive symptoms	14.1	9.9 - 18.2	2,200	12.8	6.6 - 19.0	600	14.8	9.4 - 20.2	1,500
Postpartum depressive symptoms	12.2	8.5 - 16.0	1,900	12.9	7.2 - 18.7	600	12.0	7.1 - 16.8	1,200
Hardships and Support during Pregnancy									
Experienced two or more hardships during childhood	29.8	24.3 - 35.2	4,400	28.1	18.9 - 37.2	1,400	30.8	24.0 - 37.7	3,100
Homeless or did not have a regular place to sleep	2.5 *	0.7 - 4.3	400				1.4 *	0.5 - 2.4	100
Moved due to problems paying rent or mortgage	6.0	3.1 - 8.9	900	6.5	2.7 - 10.3	300	5.8 *	1.8 - 9.7	600
Woman or partner lost job	12.1	8.5 - 15.7	1,800	10.3	5.1 - 15.5	500	13.0	8.3 - 17.7	1,300
Woman or partner had pay or hours cut back	11.8	7.8 - 15.9	1,800	14.6	6.6 - 22.7	700	10.6	6.0 - 15.1	1,100
Became separated or divorced	10.2	6.1 - 14.2	1,500	9.5	3.4 - 15.6	500	10.5	5.2 - 15.9	1,100
Had no practical or emotional support	3.1 *	0.9 - 5.3	500				2.9 *	0.3 - 5.5	300
Substance Use									
Any smoking, 3 months before pregnancy	13.7	9.3 - 18.2	2,100	13.5	5.9 - 21.0	700	14.0	8.5 - 19.4	1,400
Any smoking, 3rd trimester [†]	3.7 *	1.5 - 5.8	600	0.9	0.1 - 1.6	<100	5.0 *	1.8 - 8.2	500
Any smoking, postpartum	6.0	3.0 - 8.9	900	7.0	0.6 - 13.3	300	5.5	2.3 - 8.7	600
Any binge drinking, 3 months before pregnancy	15.3	10.9 - 19.7	2,300	21.5	13.8 - 29.2	1,000	12.5	7.1 - 17.9	1,300
Any alcohol use, 3rd trimester [†]	3.1	1.7 - 4.5	500	3.5	0.7 - 6.3	200	2.9	1.3 - 4.5	300
Pregnancy Intention and Family Planning									
Mistimed or unwanted pregnancy	27.0	21.9 - 32.1	4,100	28.9	19.7 - 38.0	1,400	26.3	20.1 - 32.5	2,700
Unsure of pregnancy intentions	18.9	14.0 - 23.7	2,900	13.9	7.7 - 20.1	700	21.4	14.9 - 27.9	2,200
Postpartum birth control use [†]	82.1	77.4 - 86.7	12,400	78.1	70.2 - 86.1	3,900	84.1	78.3 - 89.8	8,500
Infant Sleep and Breastfeeding									
Placed infant on back to sleep	80.2	75.6 - 84.8	12,200	80.7	74.3 - 87.1	3,900	79.9	73.8 - 86.0	8,200
Infant always or often shared bed	41.5	35.8 - 47.2	6,300	44.2	34.6 - 53.9	2,100	40.4	33.3 - 47.6	4,200
Intended to breastfeed, before birth	89.5	86.0 - 93.0	13,500	92.4	87.1 - 97.8	4,400	88.1	83.6 - 92.6	9,100
Intended to breastfeed exclusively, before birth	52.2	46.4 - 58.0	7,900	65.0	55.9 - 74.1	3,100	46.0	38.8 - 53.2	4,700
Breastfeeding practices	76.0	71 5 01 4	11 400	744	65.2 02.0	3 500	77 0	71.0 02.0	7 000
Any breastfeeding, 1 month after delivery Exclusive breastfeeding, 1 month after delivery	76.3	71.5 - 81.1	11,400	74.1	65.2 - 83.0 29.9 - 48.5	3,500	77.3	71.6 - 83.0 28.1 - 42.2	7,900
Any breastfeeding, 3 months after delivery	36.7	31.1 - 42.3 44.4 - 57.9	5,400 5,700	39.2	29.9 - 48.5 36.0 - 57.8	1,900 1,800	35.2 53.8	28.1 - 42.2 45.2 - 62.4	3,500 3,800
Exclusive breastfeeding, 3 months after delivery	51.2 18.5		2,000	46.9 19.9		1,800	53.8 17.9		
LACIUSIVE DIEASCIEEUING, S MONICIS AILEI UEIIVELY	18.5	14.1 - 22.9	2,000	19.9	12.5 - 27.2	800	17.9	12.2 - 23.6	1,200



MIHA Data Snapshot, Fresno County by Total Live Births, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		Fresno County			First Live Birth		Second Live Birth or More		
	%	95% CI	N	%	95% CI	N	%	95% CI	N
Service Utilization and Health Insurance Coverage									
Had a usual source of pre-pregnancy care	66.9	61.5 - 72.4	10,300	57.0	47.3 - 66.6	2,800	71.5	65.0 - 77.9	7,400
Initiated prenatal care in 1st trimester	86.4	82.3 - 90.6	13,100	92.7	* 88.3 - 97.2	4,500	83.4	77.8 - 89.1	8,600
Mom or infant needed but couldn't afford care postpartum	7.7	4.7 - 10.8	1,200	6.4	* 2.3 - 10.5	300	8.4	4.3 - 12.5	900
Participated in WIC during pregnancy [†]	69.5	65.4 - 73.6	10,500	70.8	63.8 - 77.8	3,500	69.3	63.4 - 75.3	7,000
Received CalFresh (food stamps) during pregnancy	47.5	41.9 - 53.0	7,200	32.0	23.2 - 40.7	1,600	55.3		5,600
Pre-pregnancy insurance									
Medi-Cal	46.9	41.4 - 52.4	7,200	33.8	24.4 - 43.2	1,700	53.0	46.0 - 59.9	5,500
Uninsured	22.3	17.5 - 27.0	3,400	26.0	18.0 - 34.0	1,300	20.7	14.6 - 26.7	2,100
Private	28.9	24.8 - 33.0	4,400	36.6	28.2 - 45.0	1,800	25.2	20.1 - 30.3	2,600
Prenatal insurance									
Medi-Cal	72.2	68.5 - 75.9	11,000	68.3	60.8 - 75.8	3,300	74.6	69.6 - 79.5	7,700
Uninsured	0.7 *	0.0 - 1.4	100						
Private	25.0	21.4 - 28.5	3,800	26.7	19.9 - 33.4	1,300	23.6	18.8 - 28.5	2,400
Maternal postpartum insurance									
Medi-Cal	60.8	55.8 - 65.8	9,300	61.3	52.7 - 70.0	3,000	61.0	54.4 - 67.6	6,200
Uninsured	10.6	6.9 - 14.3	1,600	6.3	* 2.3 - 10.4	300	12.8	7.7 - 17.9	1,300
Private	26.6	22.8 - 30.4	4,100	27.7	20.2 - 35.3	1,400	25.5	20.5 - 30.5	2,600
Infant health insurance									
Medi-Cal	71.7	67.7 - 75.7	10,800	69.4	61.7 - 77.0	3,400	73.1	67.8 - 78.4	7,500
Uninsured	1.9 *	0.6 - 3.2	300				2.1	* 0.3 - 3.9	200
Private	25.0	21.3 - 28.7	3,800	27.7	20.3 - 35.1	1,300	23.4	18.6 - 28.3	2,400
Maternal Demographics									
Age									
15-19	8.3	5.1 - 11.5	1,300	18.1	10.8 - 25.4	900	3.6	* 0.6 - 6.7	400
20-34	78.3	73.3 - 83.3	12,000	76.4	68.6 - 84.2	3,800	79.1	72.8 - 85.5	8,200
35+	13.4	9.2 - 17.6	2,100	5.6	* 2.2 - 8.9	300	17.3	11.4 - 23.2	1,800
Did not complete high school (or GED)	24.9	19.6 - 30.2	3,700	13.2	5.5 - 20.9	600	30.8	23.8 - 37.7	3,100
Unmarried	51.8	46.3 - 57.3	7,900	57.5	48.3 - 66.6	2,900	49.5	42.3 - 56.6	5,100
Race/Ethnicity									
White	19.9	15.9 - 23.9	3,000	20.7	13.7 - 27.7	1,000	19.4	14.3 - 24.5	2,000
Hispanic	61.7	56.6 - 66.8	9,400	55.8	46.7 - 64.9	2,800	64.5	58.0 - 71.0	6,600
Asian/Pacific Islander	12.1	8.0 - 16.2	1,800	17.1	9.4 - 24.8	800	9.8	5.0 - 14.6	1,000
Black	5.5	4.7 - 6.3	800	6.0	3.9 - 8.2	300	5.3	4.0 - 6.6	500
Born outside the U.S.	30.5	25.2 - 35.7	4,700	20.3	12.7 - 27.8	1,000	35.1	28.3 - 42.0	3,600
Speaks non-English language at home	30.4	25.1 - 35.8	4,600	28.2	19.1 - 37.3	1,400	31.2	24.5 - 38.0	3,200
Lives in a high poverty neighborhood	74.0	69.4 - 78.6	11,100	69.7	61.6 - 77.9	3,300	75.8	69.9 - 81.7	7,700
Income as a percent of the Federal Poverty Guideline									
0-100%	58.7	53.4 - 64.0	8,600	48.7	39.1 - 58.2	2,300	63.4	56.9 - 69.9	6,300
101-200%	19.0	14.0 - 24.0	2,800	22.0	12.2 - 31.7	1,000	17.8	12.1 - 23.5	1,800
> 200%	22.2	18.7 - 25.8	3,300	29.4	21.8 - 36.9	1,400	18.8	14.2 - 23.5	1,900

-- Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

* Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%)

⁺ Measure/definition may not be comparable to previous Snapshots (see Technical Notes).

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Fresno County was 562. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.