

MIHA Data Snapshot, Riverside County by Education, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Riv	erside Cou	nty			High School G	raduate o	or GED	,	Some Colleg	e	Co	llege Gradu	ate	
	%	95% CI	N	%	95% CI	N	% 95%	% CI	N	%	95% CI	N	%	95% CI	N
Total	100		29,700	16.1	12.2 - 20.0	4,700	21.8 17.5 -	- 26.0	6,300	40.7	35.7 - 45.7	11,900	21.5	17.6 - 25.4	6,200
Health Status before Pregnancy															
In good to excellent health	92.3	89.6 - 94.9	27,300	84.0	* 73.6 - 94.4	3,800	93.0 * 88.6 -	97.5	5,900	92.6	88.4 - 96.8	11,000			
Chronic conditions															
Diabetes†	2.2 *	0.7 - 3.8	700												
Hypertension†	2.6	1.3 - 3.9	800				3.0 * 0.3 -	5.8	200	2.8	* 0.6 - 5.0	300			
Asthma†	8.2	5.6 - 10.8	2,400				7.3 * 2.0 -	- 12.6	500	8.0	4.3 - 11.7	900	9.1 *	3.3 - 14.9	600
Nutrition and Weight															
Daily folic acid use, month before pregnancy	33.4	28.6 - 38.2	9,900	15.1	* 6.2 - 24.0	700	26.2 16.1 -	36.4	1,700	36.0	28.5 - 43.4	4,300	47.9	36.8 - 59.1	3,000
Overweight before pregnancy	25.9	21.2 - 30.6	7,400	12.5	* 3.1 - 22.0	500	39.0 27.0 -	- 50.9	2,300	22.8	16.3 - 29.3	2,700	27.0	16.8 - 37.2	1,700
Obese before pregnancy	25.4	20.9 - 29.9	7,300	36.0	22.2 - 49.8	1,500	25.3 14.5 -	36.1	1,500	27.9	21.0 - 34.8	3,300	15.3	7.5 - 23.2	900
Inadequate weight gain during pregnancy	18.8	14.3 - 23.3	4,700	24.4	10.5 - 38.4	900	21.3 11.0 -	31.5	1,100	15.5	9.5 - 21.6	1,600	18.0	8.6 - 27.5	900
Excessive weight gain during pregnancy	41.0	35.6 - 46.4	10,200	27.7	14.2 - 41.3	1,100	33.5 21.7 -	45.3	1,800	47.1	38.8 - 55.5	4,700	44.2	33.1 - 55.3	2,300
Food insecurity during pregnancy	21.7	17.5 - 26.0	6,400	32.7	19.7 - 45.6	1,500	25.3 14.9 -	- 35.7	1,600	21.3	14.9 - 27.7	2,500	6.6 *	1.3 - 11.9	400
Intimate Partner Violence (IPV) and Depressive Symptoms															
Physical or psychological IPV during pregnancy	5.9	3.7 - 8.2	1,800	10.9	* 3.2 - 18.5	500				5.6	* 2.2 - 9.1	700			
Prenatal depressive symptoms	13.1	9.8 - 16.4	3,900	14.7	* 5.7 - 23.8	700	14.9 7.4 -	- 22.3	900	10.9	6.4 - 15.5	1,300	10.8 *	3.5 - 18.0	700
Postpartum depressive symptoms	16.0	12.3 - 19.7	4,700	16.7	* 6.4 - 27.0	800	16.7 8.7 -	24.6	1,000	18.4	12.3 - 24.5	2,200	8.9	4.0 - 13.8	600
Hardships and Support during Pregnancy															
Experienced two or more hardships during childhood	25.6	21.2 - 30.0	7,400	26.2	13.8 - 38.5	1,200	34.9 23.6 -	46.2	2,100	26.5	19.8 - 33.1	3,100	14.1	7.3 - 20.9	900
Homeless or did not have a regular place to sleep		1.3 - 4.5	900		* 1.0 - 14.1	400			,		* 0.2 - 2.1	100			
Moved due to problems paying rent or mortgage	7.9	5.2 - 10.6	2,300	13.5	* 4.5 - 22.6	600	8.7 * 2.3 -	- 15.1	500	7.4	3.5 - 11.3	900			
Woman or partner lost job	17.6	13.7 - 21.6	5,200	17.9	* 6.9 - 28.9	800	19.7 11.1 -	- 28.3	1,200	19.6	13.1 - 26.0	2,300	12.5 *	4.9 - 20.1	800
Woman or partner had pay or hours cut back	10.5	7.6 - 13.4	3,100	12.1	* 2.7 - 21.5	600	7.0 * 2.8 -	- 11.2	400	13.0	8.1 - 17.8	1,500	9.2 *	3.7 - 14.8	600
Became separated or divorced	9.4	6.5 - 12.2	2,800	13.2	* 4.6 - 21.9	600	8.2 * 3.3 -	- 13.1	500	10.3	5.4 - 15.1	1,200	5.3 *	0.8 - 9.7	300
Had no practical or emotional support	3.5	1.7 - 5.2	1,000				4.3 * 0.2 -	8.3	300				6.8 *	1.0 - 12.6	400
Substance Use															
Any smoking, 3 months before pregnancy	10.1	7.1 - 13.0	3,000	14.2	* 5.1 - 23.3	700	11.4 * 4.1 -	18.8	700	9.8	5.4 - 14.2	1,100			
Any smoking, 3rd trimester†	1.9 *	0.6 - 3.1	600							2.3	* 0.3 - 4.3	300			
Any smoking, postpartum	4.4	2.3 - 6.4	1,300	9.1	* 1.0 - 17.1	400	3.3 * 0.5 -	- 6.1	200	4.7	* 1.3 - 8.0	600			
Any binge drinking, 3 months before pregnancy	15.4	11.6 - 19.1	4,500	9.6	* 1.9 - 17.3	400	17.1 7.9 -	- 26.3	1,000	17.4	11.1 - 23.7	2,000	15.2	8.2 - 22.2	900
Any alcohol use, 3rd trimester†	3.0	1.5 - 4.5	900							3.9	* 1.3 - 6.6	500	3.6 *	0.6 - 6.5	200
Pregnancy Intention and Family Planning															
Mistimed or unwanted pregnancy	29.2	24.6 - 33.8	8,600	31.8	18.4 - 45.1	1,500	39.5 28.1 -	- 51.0	2,500	32.1	24.9 - 39.2	3,800	11.7	6.3 - 17.1	700
Unsure of pregnancy intentions	16.1	12.2 - 20.0	4,700	25.4	13.2 - 37.5	1,200	19.2 10.0 -	- 28.4	1,200	12.9	8.0 - 17.8	1,500	13.5 *	4.8 - 22.2	800
Postpartum birth control use†	80.5	76.5 - 84.6	23,500	73.6	61.2 - 86.1	3,200	87.2 * 79.6 -	94.7	5,400	84.4	78.9 - 89.9	10,000	71.1	60.8 - 81.4	4,400
Infant Sleep and Breastfeeding															
Placed infant on back to sleep	79.5	75.4 - 83.6	23,000	80.8	69.6 - 91.9	3,700	82.5 74.3 -	90.8	5,200	76.0	69.3 - 82.8	8,800	81.5	72.5 - 90.4	4,800
Infant always or often shared bed	34.8	29.9 - 39.6	10,000		25.3 - 52.2	1,800	34.3 23.3 -		2,200		28.6 - 43.5	4,200		17.0 - 35.6	1,500
Intended to breastfeed, before birth		92.1 - 96.3	27,500			•	92.2 * 86.6 -		5,800		90.5 - 97.1	10,900			
Intended to breastfeed exclusively, before birth		61.3 - 70.9	19,300	47.5	33.5 - 61.5	2,200	56.7 45.2 -		3,600		64.4 - 78.3	8,300	81.3	73.9 - 88.8	4,800
Breastfeeding practices			•			•						•			•
Any breastfeeding, 1 month after delivery	83.1	79.3 - 86.9	23,900	77.4	66.0 - 88.8	3,600	82.8 74.6 -	91.0	5,100	80.5	74.1 - 86.8	9,400	94.3 *	90.6 - 98.0	5,500
Exclusive breastfeeding, 1 month after delivery	41.8	36.7 - 46.9	11,700	37.4	24.1 - 50.8	1,700	29.0 19.2 -	- 38.9	1,700	46.5	38.7 - 54.3	5,300	51.6	40.3 - 62.9	2,900
Any breastfeeding, 3 months after delivery	63.2	57.4 - 69.0	14,000	50.9	35.4 - 66.3	2,000	58.9 45.2 -	72.6	2,600	58.4	49.6 - 67.2	5,500	89.7 *	83.5 - 95.9	3,800
Exclusive breastfeeding, 3 months after delivery	25.9	20.8 - 31.0	5,700	14.6	* 3.2 - 26.0	600	16.2 7.2 -	- 25.2	700	28.9	20.9 - 36.9	2,700	41.9	28.8 - 55.0	1,700



MIHA Data Snapshot, Riverside County by Education, 2013-2015

Maternal and Infant Health Assessment (MIHA) Survey

	Riv	erside Cou	nty	_ -		High School Graduate or GED					College Graduate		ate		
	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N
Service Utilization and Health Insurance Coverage															
Had a usual source of pre-pregnancy care	67.4	62.7 - 72.1	19,900	53.9	40.1 - 67.7	2,500	59.1	47.6 - 70.6	3,700	70.5	63.6 - 77.4	8,400	81.5	73.4 - 89.6	5,100
Initiated prenatal care in 1st trimester	85.7	82.2 - 89.1	24,700	76.3	64.8 - 87.8	3,400	84.8	77.4 - 92.1	5,200	86.1	80.8 - 91.5	10,100	93.2	* 87.5 - 98.8	5,600
Mom or infant needed but couldn't afford care postpartum	17.5	13.6 - 21.5	5,100	19.8	8.6 - 31.1	900	14.2	6.1 - 22.3	900	19.3	13.1 - 25.5	2,300	13.6	6.0 - 21.1	800
Participated in WIC during pregnancy†	58.3	54.7 - 61.9	17,000	87.5	* 78.9 - 96.2	4,100	78.8	69.0 - 88.6	4,900	53.8	46.9 - 60.7	6,300	19.4	11.3 - 27.5	1,200
Received CalFresh (food stamps) during pregnancy	27.1	22.8 - 31.4	8,000	45.7	32.0 - 59.4	2,100	39.3	27.8 - 50.9	2,500	23.9	17.5 - 30.3	2,800	6.1	* 1.2 - 10.9	400
Pre-pregnancy insurance															
Medi-Cal	24.9	20.8 - 29.1	7,400	40.5	27.0 - 54.0	1,800	41.0	29.6 - 52.5	2,600	20.2	14.1 - 26.3	2,400	6.9	* 1.6 - 12.2	400
Uninsured	28.0	23.5 - 32.5	8,300	39.2	25.6 - 52.8	1,800	36.0	24.6 - 47.3	2,300	26.9	20.2 - 33.6	3,200	12.1	* 4.6 - 19.5	800
Private	41.9	37.7 - 46.1	12,400	15.4	* 4.8 - 25.9	700	20.8	11.6 - 30.0	1,300	46.8	39.5 - 54.1	5,500	75.6	66.6 - 84.7	4,700
Prenatal insurance															
Medi-Cal	54.1	50.3 - 58.0	16,100	84.9	* 75.1 - 94.7	4,000	76.7	67.4 - 86.0	4,900	47.5	40.3 - 54.6	5,600	17.5	9.4 - 25.6	1,100
Uninsured	1.7 *	0.4 - 3.0	500												
Private	38.2	34.4 - 42.0	11,300				18.0	9.4 - 26.5	1,100	43.2	36.0 - 50.3	5,100	77.3	68.5 - 86.0	4,800
Maternal postpartum insurance															
Medi-Cal	38.8	34.4 - 43.3	11,400	60.9	46.9 - 74.9	2,800	59.6	48.0 - 71.1	3,700	34.5	27.4 - 41.5	4,000	11.4	5.4 - 17.4	700
Uninsured	16.1	12.2 - 20.0	4,700	24.5	11.9 - 37.2	1,100	20.4	10.7 - 30.1	1,300	13.7	8.3 - 19.1	1,600	7.1	* 0.4 - 13.8	400
Private	40.4	36.3 - 44.5	11,900	8.9	* 0.2 - 17.5	400	19.5	10.1 - 28.8	1,200	45.4	38.0 - 52.7	5,300	77.7	68.9 - 86.5	4,800
Infant health insurance															
Medi-Cal	58.9	54.8 - 63.0	16,900	90.8	* 82.0 - 99.6	4,200	78.2	68.7 - 87.7	4,900	54.0	46.7 - 61.4	6,300	20.4	11.4 - 29.4	1,200
Uninsured	2.8	1.2 - 4.3	800				4.8	* 0.1 - 9.4	300	3.2	* 0.8 - 5.6	400			
Private	32.8	28.9 - 36.6	9,400				15.3	6.5 - 24.0	1,000	34.4	27.6 - 41.3	4,000	72.3	62.5 - 82.2	4,200
Maternal Demographics															
Unmarried	44.1	39.4 - 48.9	13,000	68.1	55.1 - 81.1	3,200	57.6	46.0 - 69.2	3,600	38.3	30.9 - 45.6	4,500	20.7	12.0 - 29.4	1,300
Race/Ethnicity															
White	27.5	23.2 - 31.8	8,100	12.9	* 4.2 - 21.5	600	18.0	9.0 - 27.0	1,100	31.2	24.2 - 38.2	3,600	41.7	30.9 - 52.5	2,600
Hispanic	59.7	55.1 - 64.2	17,500	78.3	67.9 - 88.7	3,700	74.4	65.0 - 83.9	4,700	56.9	49.7 - 64.2	6,600	34.1	23.8 - 44.3	2,100
Asian/Pacific Islander	6.5	3.6 - 9.4	1,900							3.5	* 1.1 - 6.0	400	19.8	9.0 - 30.5	1,200
Black	5.9	5.1 - 6.7	1,700	4.7	* 1.5 - 8.0	200	5.3	2.8 - 7.8	300	7.8	5.8 - 9.8	900	4.4	2.4 - 6.4	300
Born outside the U.S.	28.4	23.6 - 33.3	8,500	47.8	34.0 - 61.7	2,200	38.1	26.7 - 49.5	2,400	18.0	12.0 - 24.1	2,100	21.7	11.0 - 32.5	1,400
Speaks non-English language at home	19.2	15.1 - 23.3	5,700	36.6	23.2 - 49.9	1,700	28.8	18.6 - 39.1	1,800	10.1	5.6 - 14.6	1,200	12.5	* 4.6 - 20.4	800
Lives in a high poverty neighborhood	37.7	32.6 - 42.7	10,600	50.0	35.5 - 64.5	2,100	43.5	31.7 - 55.4	2,700	33.2	25.9 - 40.5	3,800	28.4	17.0 - 39.8	1,700
Income as a percent of the Federal Poverty Guideline															
0-100%	44.0	39.3 - 48.7	12,200	71.1	57.7 - 84.6	3,100	71.6	61.2 - 81.9	4,300	34.8	27.5 - 42.2	3,800	9.9	* 3.1 - 16.7	600
101-200%	25.5	20.9 - 30.1	7,100	28.9	15.4 - 42.3	1,300	23.1	13.3 - 32.9	1,400	32.0	24.5 - 39.4	3,500	14.4	7.8 - 21.1	800
> 200%	30.5	26.8 - 34.2	8,400				5.3	* 0.6 - 10.0	300	33.2	26.4 - 40.0	3,700	75.7	66.8 - 84.6	4,400

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Riverside County was 610. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

^{*} Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



MIHA Data Snapshot, Riverside County by Geographical Area, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		Riverside County	/		Urb	an			Rural/Frontier	
	%	95% CI	N	%	95%	CI	N	%	95% CI	N
Total	100		29,700	98.4 *	97.2 -	99.6	27,700	1.6 *	0.4 - 2.8	400
Health Status before Pregnancy										
In good to excellent health	92.3	89.6 - 94.9	27,300	92.8	90.2 -	95.5	25,600			
Chronic conditions										
Diabetes†	2.2 *	0.7 - 3.8	700	2.0 *	0.5 -	3.5	600			
Hypertension†	2.6	1.3 - 3.9	800	2.7	1.3 -	4.1	800			
Asthma†	8.2	5.6 - 10.8	2,400	8.3	5.6 -	10.9	2,300			
Nutrition and Weight										
Daily folic acid use, month before pregnancy	33.4	28.6 - 38.2	9,900	32.5	27.6 -	37.5	9,000	66.0 *	35.7 - 96.2	300
Overweight before pregnancy	25.9	21.2 - 30.6	7,400	25.3	20.5 -	30.1	6,800			
Obese before pregnancy	25.4	20.9 - 29.9	7,300	25.8	21.1 -	30.5	6,900			
Inadequate weight gain during pregnancy	18.8	14.3 - 23.3	4,700	18.5	13.9 -	23.1	4,300			
Excessive weight gain during pregnancy	41.0	35.6 - 46.4	10,200	40.5	34.9 -	46.1	9,500			
Food insecurity during pregnancy	21.7	17.5 - 26.0	6,400	20.8	16.5 -	25.2	5,800			
Intimate Partner Violence (IPV) and Depressive Symptoms										
Physical or psychological IPV during pregnancy	5.9	3.7 - 8.2	1,800	5.2	3.0 -	7.5	1,400			
Prenatal depressive symptoms	13.1	9.8 - 16.4	3,900	13.4	9.9 -	16.8	3,700			
Postpartum depressive symptoms	16.0	12.3 - 19.7	4,700	15.8	12.0 -	19.6	4,300			
Hardships and Support during Pregnancy	10.0	12.5	4,700	13.0	12.0	15.0	4,500			
Experienced two or more hardships during childhood	25.0	24.2 20.0	7.400	25.0	20.5	20.6	6.700			
Homeless or did not have a regular place to sleep	25.6	21.2 - 30.0	7,400	25.0	20.5 -	29.6	6,700			
• • • • • • • • • • • • • • • • • • • •	2.9	1.3 - 4.5	900	2.8	1.2 -	4.5	800			
Moved due to problems paying rent or mortgage	7.9	5.2 - 10.6	2,300	8.0	5.1 -	10.8	2,200			
Woman or partner lost job	17.6	13.7 - 21.6	5,200	17.1	13.0 -	21.1	4,700			
Woman or partner had pay or hours cut back	10.5	7.6 - 13.4	3,100	10.1	7.2 -	13.1	2,800			
Became separated or divorced	9.4	6.5 - 12.2	2,800	9.0	6.1 -	11.8	2,500			
Had no practical or emotional support	3.5	1.7 - 5.2	1,000	3.4	1.6 -	5.2	900			
Substance Use										
Any smoking, 3 months before pregnancy	10.1	7.1 - 13.0	3,000	10.4	7.3 -	13.6	2,900			
Any smoking, 3rd trimester†	1.9 *	0.6 - 3.1	600	1.9 *	0.6 -	3.2	500			
Any smoking, postpartum	4.4	2.3 - 6.4	1,300	4.6	2.4 -	6.8	1,300			
Any binge drinking, 3 months before pregnancy	15.4	11.6 - 19.1	4,500	15.2	11.3 -	19.1	4,100			
Any alcohol use, 3rd trimester†	3.0	1.5 - 4.5	900	3.2	1.6 -	4.8	900			
Pregnancy Intention and Family Planning										
Mistimed or unwanted pregnancy	29.2	24.6 - 33.8	8,600	28.9	24.2 -	33.7	8,000	47.3 *	9.7 - 84.8	200
Unsure of pregnancy intentions	16.1	12.2 - 20.0	4,700	15.8	11.8 -	19.8	4,300			
Postpartum birth control use†	80.5	76.5 - 84.6	23,500	80.4	76.1 -	84.6	21,800			
Infant Sleep and Breastfeeding										
Placed infant on back to sleep	79.5	75.4 - 83.6	23,000	79.5	75.2 -	83.8	21,500			
Infant always or often shared bed	34.8	29.9 - 39.6	10,000	33.1	28.1 -	38.1	9,000			
Intended to breastfeed, before birth	94.2	92.1 - 96.3	27,500	94.7	92.6 -	96.8	25,900			
Intended to breastfeed exclusively, before birth	66.1	61.3 - 70.9	19,300	66.9	61.9 -	71.9	18,300	54.4 *	14.7 - 94.0	200
Breastfeeding practices	00.1	52.5 , 0.5	15,550	00.5	01.3	. 1.3	20,000	54.4	50	200
Any breastfeeding, 1 month after delivery	83.1	79.3 - 86.9	23,900	83.4	79.5 -	87.2	22,400			
Exclusive breastfeeding, 1 month after delivery	41.8	36.7 - 46.9	11,700	42.8	37.5 -	48.1	11,200			
Any breastfeeding, 3 months after delivery	63.2	57.4 - 69.0	14,000	63.5	57.5 -	69.5	13,200			
Exclusive breastfeeding, 3 months after delivery	25.9	20.8 - 31.0	5,700	26.1	20.8 -	31.5	5,400			



MIHA Data Snapshot, Riverside County by Geographical Area, 2013-2015

Maternal and Infant Health Assessment (MIHA) Survey

	Riverside County				Urban		Rural/Frontier			
	%	95% CI	N	%	95% CI	N	%	95% CI	N	
Service Utilization and Health Insurance Coverage										
Had a usual source of pre-pregnancy care	67.4	62.7 - 72.1	19,900	66.7	61.8 - 71.7	18,400				
Initiated prenatal care in 1st trimester	85.7	82.2 - 89.1	24,700	86.1	82.6 - 89.7	23,300				
Mom or infant needed but couldn't afford care postpartum	17.5	13.6 - 21.5	5,100	17.6	13.5 - 21.8	4,800				
Participated in WIC during pregnancy†	58.3	54.7 - 61.9	17,000	57.5	53.6 - 61.4	15,600				
Received CalFresh (food stamps) during pregnancy	27.1	22.8 - 31.4	8,000	26.6	22.1 - 31.1	7,400	57.4 *	20.2 - 94.6	200	
Pre-pregnancy insurance										
Medi-Cal	24.9	20.8 - 29.1	7,400	24.1	19.7 - 28.4	6,600				
Uninsured	28.0	23.5 - 32.5	8,300	27.9	23.2 - 32.6	7,700				
Private	41.9	37.7 - 46.1	12,400	42.6	38.2 - 47.1	11,700				
Prenatal insurance										
Medi-Cal	54.1	50.3 - 58.0	16,100	53.2	49.0 - 57.3	14,700				
Uninsured	1.7 *	0.4 - 3.0	500	1.7 *	0.4 - 3.1	500				
Private	38.2	34.4 - 42.0	11,300	38.8	34.7 - 42.8	10,700				
Maternal postpartum insurance										
Medi-Cal	38.8	34.4 - 43.3	11,400	38.2	33.6 - 42.9	10,500				
Uninsured	16.1	12.2 - 20.0	4,700	15.9	11.9 - 20.0	4,400				
Private	40.4	36.3 - 44.5	11,900	40.9	36.6 - 45.3	11,300				
Infant health insurance										
Medi-Cal	58.9	54.8 - 63.0	16,900	58.1	53.7 - 62.4	15,700				
Uninsured	2.8	1.2 - 4.3	800	2.9	1.3 - 4.6	800				
Private	32.8	28.9 - 36.6	9,400	33.1	29.1 - 37.2	8,900				
Maternal Demographics										
Age										
15-19	6.7	4.2 - 9.2	2,000	7.1	4.4 - 9.8	2,000				
20-34	78.3	73.9 - 82.7	23,300	77.1	72.4 - 81.7	21,400				
35+	15.0	11.2 - 18.9	4,500	15.8	11.7 - 19.9	4,400				
Did not complete high school (or GED)	16.1	12.2 - 20.0	4,700	15.6	11.6 - 19.7	4,300				
Unmarried	44.1	39.4 - 48.9	13,000	43.6	38.6 - 48.6	12,000	55.5 *	17.7 - 93.3	200	
Race/Ethnicity										
White	27.5	23.2 - 31.8	8,100	28.4	23.9 - 32.9	7,800				
Hispanic	59.7	55.1 - 64.2	17,500	58.3	53.5 - 63.1	16,000				
Asian/Pacific Islander	6.5	3.6 - 9.4	1,900	7.0	3.9 - 10.0	1,900				
Black	5.9	5.1 - 6.7	1,700	6.0	5.1 - 6.9	1,600				
Born outside the U.S.	28.4	23.6 - 33.3	8,500	27.0	21.9 - 32.0	7,500				
Speaks non-English language at home	19.2	15.1 - 23.3	5,700	18.2	14.0 - 22.3	5,000				
Lives in a high poverty neighborhood	37.7	32.6 - 42.7	10,600	36.7	31.6 - 41.8	10,200				
Income as a percent of the Federal Poverty Guideline			•			*				
0-100%	44.0	39.3 - 48.7	12,200	42.7	37.8 - 47.7	11,100	51.7 *	9.5 - 93.8	200	
101-200%	25.5	20.9 - 30.1	7,100	26.4	21.6 - 31.2	6,900				
> 200%	30.5	26.8 - 34.2	8,400	30.9	26.9 - 34.8	8,000				

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Riverside County was 610. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. Urban and Rural/Frontier designations are based on the population size or densities of Medical Service Study Areas. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

^{*} Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



MIHA Data Snapshot, Riverside County by Income, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Riverside Coun	ty	0-100% Federal Poverty	Guideline	101-200% Federal Poverty	Guideline	ne > 200% Federal Poverty Guideline		
	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N	
Total	100	29,700	44.0 39.3 - 48.7	12,200	25.5 20.9 - 30.1	7,100	30.5 26.8 - 34.2	8,400	
Health Status before Pregnancy									
In good to excellent health	92.3 89.6 - 94.9	27,300	87.5 82.3 - 92.7	10,700	97.4 * 95.1 - 99.6	6,800	93.8 * 88.8 - 98.7	7,900	
Chronic conditions									
Diabetes†	2.2 * 0.7 - 3.8	700	4.8 * 1.2 - 8.4	600					
Hypertension†	2.6 1.3 - 3.9	800	3.9 * 1.4 - 6.5	500			2.6 * 0.3 - 5.0	200	
Asthma†	8.2 5.6 - 10.8	2,400	10.9 6.2 - 15.6	1,300	5.4 * 1.1 - 9.7	400	7.9 * 3.2 - 12.6	700	
Nutrition and Weight									
Daily folic acid use, month before pregnancy	33.4 28.6 - 38.2	9,900	21.5 14.8 - 28.1	2,600	33.9 23.7 - 44.1	2,400	49.0 39.5 - 58.4	4,100	
Overweight before pregnancy	25.9 21.2 - 30.6	7,400	28.3 20.2 - 36.3	3,300	22.3 14.1 - 30.5	1,500	26.2 17.7 - 34.7	2,200	
Obese before pregnancy	25.4 20.9 - 29.9	7,300	25.7 18.5 - 33.0	3,000	28.2 18.5 - 37.9	1,900	25.0 17.1 - 33.0	2,100	
Inadequate weight gain during pregnancy	18.8 14.3 - 23.3	4,700	20.9 13.4 - 28.3	2,100	19.3 10.0 - 28.6	1,200	11.8 5.9 - 17.7	800	
Excessive weight gain during pregnancy	41.0 35.6 - 46.4	10,200	36.8 28.0 - 45.5	3,700	36.7 26.2 - 47.2	2,300	52.8 43.1 - 62.5	3,800	
Food insecurity during pregnancy	21.7 17.5 - 26.0	6,400	36.5 28.4 - 44.6	4,400	18.4 10.1 - 26.8	1,300	6.3 * 1.7 - 11.0	500	
Intimate Partner Violence (IPV) and Depressive Symptoms									
Physical or psychological IPV during pregnancy	5.9 3.7 - 8.2	1,800	11.0 6.1 - 15.8	1,300	4.5 * 0.7 - 8.3	300			
Prenatal depressive symptoms	13.1 9.8 - 16.4	3,900	17.6 11.6 - 23.7	2,100	13.5 7.3 - 19.6	1,000	8.2 * 2.6 - 13.8	700	
Postpartum depressive symptoms	16.0 12.3 - 19.7	4,700	19.3 13.0 - 25.7	2,300	14.0 6.6 - 21.5	1,000	16.4 9.8 - 23.0	1,400	
Hardships and Support during Pregnancy									
Experienced two or more hardships during childhood	25.6 21.2 - 30.0	7,400	34.9 27.0 - 42.7	4,100	21. 5 12.3 - 30.7	1,500	20.2 13.2 - 27.2	1,700	
Homeless or did not have a regular place to sleep	2.9 1.3 - 4.5	900	5.6 2.4 - 8.8	700					
Moved due to problems paying rent or mortgage	7.9 5.2 - 10.6	2,300	14.7 9.0 - 20.4	1,800	5.4 * 0.6 - 10.1	400			
Woman or partner lost job	17.6 13.7 - 21.6	5,200	27. 5 20.1 - 34.8	3,300	12.5 5.7 - 19.4	900	8.5 * 2.9 - 14.0	700	
Woman or partner had pay or hours cut back	10.5 7.6 - 13.4	3,100	13.8 8.5 - 19.1	1,700	10.5 5.2 - 15.8	700	6.7 3.0 - 10.4	600	
Became separated or divorced	9.4 6.5 - 12.2	2,800	16.5 10.8 - 22.3	2,000	5.6 * 1.1 - 10.1	400			
Had no practical or emotional support	3.5 1.7 - 5.2	1,000	4.6 * 1.4 - 7.7	600					
Substance Use									
Any smoking, 3 months before pregnancy	10.1 7.1 - 13.0	3,000	16.3 10.2 - 22.4	1,900	4.8 * 1.0 - 8.7	300	6.4 * 2.2 - 10.5	500	
Any smoking, 3rd trimester†	1.9 * 0.6 - 3.1	600	2.6 * 0.4 - 4.7	300					
Any smoking, postpartum	4.4 2.3 - 6.4	1,300	6.8 2.9 - 10.8	800					
Any binge drinking, 3 months before pregnancy	15.4 11.6 - 19.1	4,500	15.0 8.5 - 21.4	1,800	19.4 11.2 - 27.7	1,400	14.9 9.1 - 20.8	1,200	
Any alcohol use, 3rd trimester†	3.0 1.5 - 4.5	900	1.9 * 0.3 - 3.5	200	4.9 * 0.4 - 9.4	300	3.7 * 1.1 - 6.3	300	
Pregnancy Intention and Family Planning									
Mistimed or unwanted pregnancy	29.2 24.6 - 33.8	8,600	40.4 32.2 - 48.6	4,900	25.6 16.8 - 34.4	1,800	18.3 11.5 - 25.1	1,500	
Unsure of pregnancy intentions	16.1 12.2 - 20.0	4,700	18.2 12.0 - 24.4	2,200	19.1 10.6 - 27.5	1,300	10.2 * 3.8 - 16.7	900	
Postpartum birth control use†	80.5 76.5 - 84.6	23,500	80.9 74.5 - 87.2	9,400	88.6 82.2 - 95.1	6,200	76.8 68.7 - 84.8	6,500	
Infant Sleep and Breastfeeding									
Placed infant on back to sleep	79.5 75.4 - 83.6	23,000	74.1 66.8 - 81.3	8,600	85.0 77.4 - 92.6	6,000	83.2 76.2 - 90.2	6,800	
Infant always or often shared bed	34.8 29.9 - 39.6	10,000	41.4 33.1 - 49.6	4,800	29.7 20.5 - 38.9	2,100	30.1 21.6 - 38.7	2,500	
Intended to breastfeed, before birth	94.2 92.1 - 96.3	27,500	94.0 90.7 - 97.3	11,100	90.8 * 85.1 - 96.5	6,400	96.8 * 94.3 - 99.3	8,000	
Intended to breastfeed exclusively, before birth	66.1 61.3 - 70.9	19,300	55.3 46.8 - 63.7	6,500	67.2 57.5 - 76.9	4,700	80.1 73.4 - 86.9	6,600	
Breastfeeding practices			00.0	_			0=0 =		
Any breastfeeding, 1 month after delivery	83.1 79.3 - 86.9	23,900	80.9 74.6 - 87.2	9,300	79.7 71.1 - 88.3	5,600	85.9 79.7 - 92.0	7,000	
Exclusive breastfeeding, 1 month after delivery	41.8 36.7 - 46.9	11,700	37.3 29.2 - 45.4	4,200	41.0 30.9 - 51.1	2,900	44.2 35.0 - 53.5	3,500	
Any breastfeeding, 3 months after delivery	63.2 57.4 - 69.0	14,000	57.2 47.5 - 66.9	5,100	61.0 49.2 - 72.9	3,300	73.5 64.1 - 82.8	4,400	
Exclusive breastfeeding, 3 months after delivery	25.9 20.8 - 31.0	5,700	18.7 11.3 - 26.1	1,700	27.6 17.2 - 38.1	1,500	31.1 21.3 - 41.0	1,800	



MIHA Data Snapshot, Riverside County by Income, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

				0-100% F	ederal Poverty	Guideline	<u> </u>				> 200% Federal Poverty Guideline		
	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	
Service Utilization and Health Insurance Coverage													
Had a usual source of pre-pregnancy care	67.4	62.7 - 72.1	19,900	61.5	53.5 - 69.5	7,500	59.4	49.1 - 69.6	4,100	82.1	74.8 - 89.3	6,900	
Initiated prenatal care in 1st trimester	85.7	82.2 - 89.1	24,700	80.8	74.9 - 86.8	9,500	83.8	75.8 - 91.8	5,800	94.0 *	89.6 - 98.3	7,800	
Mom or infant needed but couldn't afford care postpartum	17.5	13.6 - 21.5	5,100	21.6	14.6 - 28.5	2,600	22.3	13.6 - 31.0	1,600	7.6 *	2.4 - 12.9	600	
Participated in WIC during pregnancy†	58.3	54.7 - 61.9	17,000	85.9	80.2 - 91.6	10,500	64.6	55.3 - 74.0	4,600	8.2 *	3.2 - 13.1	700	
Received CalFresh (food stamps) during pregnancy	27.1	22.8 - 31.4	8,000	54.1	45.9 - 62.3	6,600	14.8	7.7 - 21.9	1,000				
Pre-pregnancy insurance													
Medi-Cal	24.9	20.8 - 29.1	7,400	49.9	41.8 - 58.1	6,100	9.3	4.3 - 14.3	600				
Uninsured	28.0	23.5 - 32.5	8,300	38.1	30.1 - 46.1	4,600	37.7	27.4 - 48.1	2,600	6.1 *	1.9 - 10.3	500	
Private	41.9	37.7 - 46.1	12,400	7.5	3.2 - 11.7	900	47.2	36.8 - 57.5	3,300	87.3	81.4 - 93.2	7,400	
Prenatal insurance													
Medi-Cal	54.1	50.3 - 58.0	16,100	90.2	85.6 - 94.7	11,000	50.1	40.0 - 60.3	3,500	4.9 *	0.4 - 9.4	400	
Uninsured	1.7	* 0.4 - 3.0	500	2.2 *	0.1 - 4.3	300							
Private	38.2	34.4 - 42.0	11,300	5.4 *	1.9 - 8.8	700	39.1	29.2 - 48.9	2,800	86.4	79.8 - 92.9	7,300	
Maternal postpartum insurance													
Medi-Cal	38.8	34.4 - 43.3	11,400	66.4	58.3 - 74.4	8,000	34.9	25.0 - 44.8	2,400	2.3 *	0.4 - 4.1	200	
Uninsured	16.1	12.2 - 20.0	4,700	24.3	16.8 - 31.8	2,900	15.3	7.6 - 23.1	1,100	5.9 *	0.7 - 11.1	500	
Private	40.4	36.3 - 44.5	11,900	6.9 *	2.3 - 11.5	800	44.3	34.0 - 54.6	3,100	85.4	78.7 - 92.2	7,200	
Infant health insurance													
Medi-Cal	58.9	54.8 - 63.0	16,900	92.9 *	88.6 - 97.1	10,700	60.3	50.3 - 70.4	4,300	8.1 *	3.3 - 12.8	700	
Uninsured	2.8	1.2 - 4.3	800							3.8 *	0.5 - 7.1	300	
Private	32.8	28.9 - 36.6	9,400	4.4 *	0.6 - 8.2	500	30.0	20.5 - 39.5	2,100	79.1	71.5 - 86.8	6,500	
Maternal Demographics													
Did not complete high school (or GED)	16.1	12.2 - 20.0	4,700	26.4	19.1 - 33.8	3,100	18.0	9.1 - 27.0	1,300				
Unmarried	44.1	39.4 - 48.9	13,000	71.3	63.8 - 78.8	8,600	33.4	23.9 - 42.9	2,400	13.6	7.7 - 19.5	1,100	
Race/Ethnicity													
White	27.5	23.2 - 31.8	8,100	16.5	10.4 - 22.6	2,000	25.1	16.5 - 33.7	1,800	50.2	40.8 - 59.6	4,100	
Hispanic	59.7	55.1 - 64.2	17,500	72.5	65.9 - 79.2	8,800	66.3	57.2 - 75.4	4,700	31.1	22.5 - 39.6	2,600	
Asian/Pacific Islander	6.5	3.6 - 9.4	1,900	2.6 *	0.1 - 5.0	300	3.8	* 0.7 - 6.9	300	13.7 *	5.6 - 21.9	1,100	
Black	5.9	5.1 - 6.7	1,700	7.9	5.9 - 9.9	1,000	4.8	2.6 - 7.0	300	4.5	3.0 - 6.1	400	
Born outside the U.S.		23.6 - 33.3	8,500	_	27.3 - 43.6	4,300		10.4 - 28.0	1,400	21.1	12.1 - 30.0	1,800	
Speaks non-English language at home		15.1 - 23.3	5,700		21.8 - 37.0	3,500	14.2		1,000	6.6 *		600	
Lives in a high poverty neighborhood	37.7	32.6 - 42.7	10,600	50.2	41.6 - 58.8	5,700	29.9	20.7 - 39.1	2,100	24.1	14.7 - 33.4	2,000	

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Riverside County was 610. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

^{*} Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



MIHA Data Snapshot, Riverside County by Maternal Age, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Riverside County 15-19 Years				20-34 Years		35 Years or Older	
	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N
Total	100	29,700	6.7 4.2 - 9.2	2,000	78.3 73.9 - 82.7	23,300	15.0 11.2 - 18.9	4,500
Health Status before Pregnancy								
In good to excellent health	92.3 89.6 - 94.9	27,300	86.8 * 74.7 - 98.8	1,700	92.8 90.0 - 95.7	21,600	91.8 * 84.1 - 99.6	4,000
Chronic conditions								
Diabetes†	2.2 * 0.7 - 3.8	700			1.2 * 0.0 - 2.3	300		
Hypertension†	2.6 1.3 - 3.9	800			2. 5 1.1 - 4.0	600		
Asthma†	8.2 5.6 - 10.8	2,400			8.1 5.2 - 11.0	1,900	7.8 * 0.6 - 15.0	300
Nutrition and Weight								
Daily folic acid use, month before pregnancy	33.4 28.6 - 38.2	9,900	17.8 * 4.1 - 31.5	400	32.2 26.8 - 37.5	7,500	46.7 32.5 - 60.9	2,100
Overweight before pregnancy	25 .9 21.2 - 30.6	7,400	23.4 * 7.4 - 39.4	500	24.6 19.4 - 29.8	5,500	33.8 20.4 - 47.3	1,500
Obese before pregnancy	25.4 20.9 - 29.9	7,300	14.8 * 0.3 - 29.2	300	26.4 21.4 - 31.3	5,900	25.0 11.4 - 38.6	1,100
Inadequate weight gain during pregnancy	18.8 14.3 - 23.3	4,700	23.2 * 5.2 - 41.2	400	18.0 13.3 - 22.8	3,500	20.5 * 6.8 - 34.2	800
Excessive weight gain during pregnancy	41.0 35.6 - 46.4	10,200	44.5 23.5 - 65.5	800	43.1 37.0 - 49.3	8,300	28.7 15.8 - 41.5	1,100
Food insecurity during pregnancy	21.7 17.5 - 26.0	6,400	31.9 13.9 - 50.0	600	21.8 16.9 - 26.6	5,000	17.0 * 5.8 - 28.1	800
Intimate Partner Violence (IPV) and Depressive Symptoms								
Physical or psychological IPV during pregnancy	5.9 3.7 - 8.2	1,800			6.8 4.1 - 9.4	1,600		
Prenatal depressive symptoms	13.1 9.8 - 16.4	3,900			13.0 9.5 - 16.6	3,000	14.3 * 4.2 - 24.5	600
Postpartum depressive symptoms	16.0 12.3 - 19.7	4,700			16.6 12.4 - 20.7	3,800	15.2 * 5.1 - 25.4	700
Hardships and Support during Pregnancy								
Experienced two or more hardships during childhood	25 .6 21.2 - 30.0	7,400	27.3 * 11.0 - 43.7	500	25.5 20.6 - 30.5	5,700	24.9 12.3 - 37.5	1,100
Homeless or did not have a regular place to sleep	2.9 1.3 - 4.5	900			2.8 * 1.0 - 4.6	700		
Moved due to problems paying rent or mortgage	7.9 5.2 - 10.6	2,300			7.4 4.5 - 10.3	1,700	9.4 * 1.4 - 17.4	400
Woman or partner lost job	17.6 13.7 - 21.6	5,200	24.2 * 7.2 - 41.3	500	16.7 12.4 - 21.0	3,800	19.6 8.2 - 30.9	900
Woman or partner had pay or hours cut back	10.5 7.6 - 13.4	3,100			11.7 8.3 - 15.0	2,700	8.6 * 1.1 - 16.1	400
Became separated or divorced	9.4 6.5 - 12.2	2,800	13.4 * 1.8 - 25.1	300	9.5 6.2 - 12.7	2,200	7.0 * 1.5 - 12.6	300
Had no practical or emotional support	3.5 1.7 - 5.2	1,000			3.5 1.6 - 5.4	800		
Substance Use								
Any smoking, 3 months before pregnancy	10.1 7.1 - 13.0	3,000			10.7 7.2 - 14.2	2,400	5.9 * 0.7 - 11.0	300
Any smoking, 3rd trimester†	1.9 * 0.6 - 3.1	600			2.0 * 0.6 - 3.3	400		
Any smoking, postpartum	4.4 2.3 - 6.4	1,300			3.5 * 1.4 - 5.6	800	7.8 * 0.9 - 14.6	300
Any binge drinking, 3 months before pregnancy	15.4 11.6 - 19.1	4,500			17.4 12.9 - 21.9	3,900	7.7 * 1.1 - 14.4	300
Any alcohol use, 3rd trimester†	3.0 1.5 - 4.5	900			3.3 1.5 - 5.1	800		
Pregnancy Intention and Family Planning								
Mistimed or unwanted pregnancy	29.2 24.6 - 33.8	8,600	60.0 40.9 - 79.1	1,200	28.6 23.5 - 33.8	6,600	17.9 7.6 - 28.3	800
Unsure of pregnancy intentions	16.1 12.2 - 20.0	4,700	22.0 * 5.6 - 38.3	400	13.2 9.5 - 16.9	3,100	28.8 14.4 - 43.1	1,300
Postpartum birth control use†	80.5 76.5 - 84.6	23,500	79.7 * 65.2 - 94.1	1,500	82.9 78.7 - 87.0	18,900	69.0 55.2 - 82.8	3,100
Infant Sleep and Breastfeeding	-							
Placed infant on back to sleep	79.5 75.4 - 83.6	23,000	77.7 * 62.5 - 92.9	1,500	80.9 76.5 - 85.3	18,100	73.3 59.7 - 87.0	3,300
Infant always or often shared bed	34.8 29.9 - 39.6	10,000	47.3 27.6 - 66.9	900	33.9 28.5 - 39.3	7,600	33.4 20.0 - 46.8	1,500
Intended to breastfeed, before birth	94.2 92.1 - 96.3	27,500			93.0 90.3 - 95.6	21,100	 	
Intended to breastfeed exclusively, before birth	66.1 61.3 - 70.9	19,300	53.5 33.8 - 73.2	1,100	66.2 60.9 - 71.6	15,000	70.9 57.9 - 84.0	3,200
Breastfeeding practices Any breastfeeding 1 month after delivery	001 700 000	22.000	77.0 * 64.2 04.4	1 500	000 700 075	10.000	046*752 040	2.700
Any breastfeeding, 1 month after delivery Exclusive breastfeeding, 1 month after delivery	83.1 79.3 - 86.9	23,900	77.8 * 61.3 - 94.4	1,500	83.2 79.0 - 87.5	18,600	84.6 * 75.2 - 94.0	3,700
Any breastfeeding, 3 months after delivery	41.8 36.7 - 46.9 63.2 57.4 - 69.0	11,700	36.0 18.1 - 53.8	700 600	45.2 39.5 - 50.9	9,900	26.6 14.5 - 38.7 77.4 64.8 - 90.0	1,100 2,600
Exclusive breastfeeding, 3 months after delivery	25.9 20.8 - 31.0	14,000 5,700	45.8 21.9 - 69.7	200	61.7 55.1 - 68.2 28.6 22.7 - 34.6	10,800 4,900		2,600 500
Exclusive preastreeuring, 3 months after delivery	25.9 20.8 - 31.0	5,700	13.2 * 2.0 - 24.5	200	Zð.D ZZ./ - 34.b	4,900	16.5 * 5.1 - 27.8	500



MIHA Data Snapshot, Riverside County by Maternal Age, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Riv	erside Coun	ty	15-19 Years		20-34 Years		35 Years or Older		
	%	95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N	
Service Utilization and Health Insurance Coverage										
Had a usual source of pre-pregnancy care	67.4	52.7 - 72.1	19,900	60.9 42.1 - 79.6	1,200	66.6 61.3 - 71.9	15,500	75.1 62.6 - 87.5	3,300	
Initiated prenatal care in 1st trimester	85.7	32.2 - 89.1	24,700	62.9 44.2 - 81.7	1,200	86.4 82.6 - 90.2	19,700	92.5 * 86.1 - 98.8	3,800	
Mom or infant needed but couldn't afford care postpartum	17.5	13.6 - 21.5	5,100	18.1 * 5.0 - 31.3	400	17.5 13.0 - 22.0	4,000	17.4 * 6.5 - 28.4	800	
Participated in WIC during pregnancy†	58.3	54.7 - 61.9	17,000			58.9 54.6 - 63.1	13,600	40.9 26.3 - 55.4	1,700	
Received CalFresh (food stamps) during pregnancy	27.1	22.8 - 31.4	8,000	34.5 16.1 - 52.9	700	28.7 23.7 - 33.7	6,700	15.6 * 4.9 - 26.3	700	
Pre-pregnancy insurance										
Medi-Cal	24.9	20.8 - 29.1	7,400	34.9 17.4 - 52.5	700	27.4 22.5 - 32.4	6,300	7.1 * 1.0 - 13.2	300	
Uninsured	28.0	23.5 - 32.5	8,300	28.0 * 10.7 - 45.4	600	28.4 23.3 - 33.5	6,600	26.1 13.0 - 39.2	1,100	
Private	41.9	37.7 - 46.1	12,400	26.6 * 8.5 - 44.7	500	39.3 34.5 - 44.1	9,100	62.8 49.3 - 76.4	2,700	
Prenatal insurance										
Medi-Cal	54.1	50.3 - 58.0	16,100	87.0 * 74.4 - 99.5	1,700	55.5 51.0 - 60.1	12,900	31.9 18.6 - 45.2	1,400	
Uninsured	1.7 *	0.4 - 3.0	500			1.5 * 0.1 - 2.8	300			
Private	38.2	34.4 - 42.0	11,300	6.1 * 0.6 - 11.6	100	36.8 32.3 - 41.2	8,600	60.2 46.5 - 73.8	2,700	
Maternal postpartum insurance										
Medi-Cal	38.8	34.4 - 43.3	11,400	72.0 * 54.6 - 89.5	1,400	40.3 35.3 - 45.3	9,300	16.3 * 6.7 - 26.0	700	
Uninsured	16.1	12.2 - 20.0	4,700	16.1 * 3.0 - 29.2	300	15.5 11.1 - 19.9	3,600	19.3 * 7.5 - 31.1	900	
Private	40.4	36.3 - 44.5	11,900			39.1 34.4 - 43.8	9,000	61.8 48.2 - 75.3	2,700	
Infant health insurance										
Medi-Cal	58.9	54.8 - 63.0	16,900			58.5 53.6 - 63.3	13,100	48.3 34.3 - 62.4	2,100	
Uninsured	2.8	1.2 - 4.3	800			3.2 * 1.3 - 5.1	700			
Private	32.8	28.9 - 36.6	9,400			32.0 27.6 - 36.5	7,200	48.1 34.0 - 62.2	2,100	
Maternal Demographics										
Did not complete high school (or GED)	16.1	12.2 - 20.0	4,700	46.2 26.0 - 66.4	900	14.4 10.2 - 18.5	3,300	11.9 * 1.4 - 22.4	500	
Unmarried	44.1	39.4 - 48.9	13,000			42.7 37.2 - 48.1	9,800	27.8 16.0 - 39.5	1,200	
Race/Ethnicity										
White	27.5	23.2 - 31.8	8,100			29.3 24.4 - 34.2	6,700	27.9 15.0 - 40.7	1,200	
Hispanic	_	55.1 - 64.2	17,500	85.7 * 73.7 - 97.8	1,700	58.9 53.6 - 64.2	13,500	52.0 38.0 - 65.9	2,300	
Asian/Pacific Islander	6.5	3.6 - 9.4	1,900			5.4 2.3 - 8.5	1,200	13.8 * 4.3 - 23.2	600	
Black		5.1 - 6.7	1,700	5.2 * 0.3 - 10.1	100	5.9 4.9 - 6.9	1,400	6.4 3.1 - 9.7	300	
Born outside the U.S.		23.6 - 33.3	8,500	14.5 * 4.8 - 24.3	300	26.0 20.7 - 31.3	6,000	47.4 33.1 - 61.7	2,100	
Speaks non-English language at home	_	15.1 - 23.3	5,700	13.9 * 1.0 - 26.8	300	17.2 12.9 - 21.4	4,000	32.1 18.3 - 45.8	1,400	
Lives in a high poverty neighborhood	_	32.6 - 42.7	10,600	53.3 33.7 - 72.9	1,100	38.5 32.8 - 44.2	8,400	27.0 14.1 - 39.8	1,200	

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Riverside County was 610. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

^{*} Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



MIHA Data Snapshot, Riverside County by Neighborhood Poverty, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Riv	erside Cou	nty		9% of Reside elow Povert			of Reside			9.9% of Resid			0% of Reside	
	%	95% CI	N	%	95% CI	N	% 9	95% CI	N	%	95% CI	N	%	95% CI	N
Total	100		29,700	12.3	9.1 - 15.5	3,500	22.4 18.	.1 - 26.7	6,300	27.7	23.0 - 32.3	7,800	37.7	32.6 - 42.7	10,600
Health Status before Pregnancy															
In good to excellent health Chronic conditions	92.3	89.6 - 94.9	27,300				98.7 * 97.	.6 - 99.9	6,200	93.9	* 89.1 - 98.7	7,300	87.6	82.0 - 93.2	9,200
Diabetes†	2.2 *	0.7 - 3.8	700										2.6 *	* 0.4 - 4.9	300
Hypertension† Asthma†	2.6 8.2	1.3 - 3.9 5.6 - 10.8	800 2,400	 9.9 *	2.0 - 17.7	300	 8.0 * 3.	.0 - 13.1	500	3.1 ° 6.2	0.4 - 5.9 2.6 - 9.9	200 500		* 1.1 - 6.6 4.6 - 15.4	400 1,000
Nutrition and Weight															
Daily folic acid use, month before pregnancy	33.4	28.6 - 38.2	9,900	50.2	36.0 - 64.4	1,700	41.0 30.	.2 - 51.7	2,600	21.5	13.0 - 30.1	1,700	31.2	22.8 - 39.7	3,300
Overweight before pregnancy		21.2 - 30.6	7,400		10.1 - 33.1	700	24.0 14.		1,500		14.9 - 32.6	1,800		21.6 - 38.9	3,100
Obese before pregnancy		20.9 - 29.9	7,300	_	6.1 - 24.6	500	22.1 13.		1,300	_	25.4 - 45.0	2,700		16.6 - 31.8	2,400
Inadequate weight gain during pregnancy		14.3 - 23.3	4,700		2.6 - 26.0	400	15.5 * 6.		900		13.8 - 31.8	1,500		11.3 - 26.2	1,600
Excessive weight gain during pregnancy		35.6 - 46.4	10,200		36.5 - 66.8	1,600	44.6 33.		2,500	_	29.7 - 50.9	2,600		26.3 - 44.6	3,100
Food insecurity during pregnancy		17.5 - 26.0	6,400		13.7 - 40.3	900	10.9 * 3.		700		14.9 - 32.7	1,800		14.7 - 29.2	2,300
Intimate Partner Violence (IPV) and Depressive Symptoms			,	_,			20.5					,			,
Physical or psychological IPV during pregnancy	5.9	3.7 - 8.2	1,800							5.4	1.0 - 9.8	400	6.8*	* 2.8 - 10.9	700
Prenatal depressive symptoms		9.8 - 16.4	3,900	6.6 *	0.9 - 12.2	200	13.6 6.	.4 - 20.7	900	13.3	7.3 - 19.2	1,000		8.7 - 21.6	1,600
Postpartum depressive symptoms		12.3 - 19.7	4,700		2.4 - 18.8	400		.5 - 21.8	900		12.7 - 27.9	1,600	_	9.0 - 22.1	1,600
Hardships and Support during Pregnancy	10.0		.,	10.0			11.2			20.5		_,	15.0		_,
Experienced two or more hardships during childhood	25.6	21.2 - 30.0	7,400	22.2	10.7 - 35.7	800	19.2 10.	.5 - 28.0	1,200	3/15	24.6 - 44.4	2,600	21.3	14.7 - 28.0	2,200
Homeless or did not have a regular place to sleep	2.9	1.3 - 4.5	900	25.2	10.7 33.7	800	15.2 10.	.5 20.0	1,200		0.2 - 7.2	300	_	* 0.7 - 5.3	300
Moved due to problems paying rent or mortgage	7.9	5.2 - 10.6	2,300				9.7 * 2.	6 - 168	600		3.4 - 15.2	700		* 2.7 - 10.9	700
Woman or partner lost job		13.7 - 21.6	5,200				_	.5 - 25.8	1,100		10.0 - 25.4	1,400		13.7 - 28.6	2,200
Woman or partner had pay or hours cut back		7.6 - 13.4	3,100	4.4 *	0.1 - 8.8	200		.0 - 17.5	700	15.4	8.4 - 22.4	1,200		3.2 - 10.9	700
Became separated or divorced	9.4	6.5 - 12.2	2,800				_	.2 - 3.7	100	11.6	5.7 - 17.4	900	11.3	5.9 - 16.7	1,200
Had no practical or emotional support		1.7 - 5.2	1,000										_	* 0.7 - 5.9	300
Substance Use			•												
Any smoking, 3 months before pregnancy	10.1	7.1 - 13.0	3,000				8.9 * 3.	.2 - 14.5	600	17.0	8.9 - 25.2	1,300	7.9	4.1 - 11.7	800
Any smoking, 3rd trimester†	1.9 *		600							_	0.5 - 7.4	300			
Any smoking, postpartum	4.4	2.3 - 6.4	1,300								2.4 - 14.5	700	2.8*	* 0.8 - 4.8	300
Any binge drinking, 3 months before pregnancy		11.6 - 19.1	4,500	10.6 *	2.4 - 18.8	400	15.2 7.	.5 - 22.9	900		10.2 - 26.9	1,400	_	7.7 - 20.5	1,500
Any alcohol use, 3rd trimester†		1.5 - 4.5	900								0.2 - 5.1	200		* 0.6 - 5.1	300
Pregnancy Intention and Family Planning															
Mistimed or unwanted pregnancy	29.2	24.6 - 33.8	8,600	22.7	11.4 - 34.0	800	23.2 14.	.4 - 32.0	1,500	29.1	20.1 - 38.1	2,200	34.9	26.3 - 43.5	3,700
Unsure of pregnancy intentions		12.2 - 20.0	4,700		0.4 - 17.3	300		.6 - 25.1	1,000		10.2 - 26.3	1,400		9.0 - 21.6	1,600
Postpartum birth control use†		76.5 - 84.6	23,500		73.8 - 93.2	2,800	79.8 71.		5,000		74.5 - 89.7	6,300		71.6 - 86.2	8,200
Infant Sleep and Breastfeeding															
Placed infant on back to sleep	79.5	75.4 - 83.6	23,000	77.7	65.8 - 89.6	2,700	77.7 68.	.2 - 87.2	4,700	83.8	76.5 - 91.2	6,400	77.6	70.2 - 85.0	8,000
Infant always or often shared bed		29.9 - 39.6	10,000		19.5 - 47.3	1,100	32.3 21.		2,000		23.7 - 42.8	2,500	_	26.6 - 42.9	3,600
Intended to breastfeed, before birth		92.1 - 96.3	27,500		88.1 - 99.3	3,200			•		89.8 - 98.3	7,200		* 91.2 - 98.0	10,000
Intended to breastfeed exclusively, before birth Breastfeeding practices		61.3 - 70.9	19,300		56.4 - 82.7	2,400	73.8 64.	.5 - 83.1	4,500		57.2 - 77.1	5,100		52.9 - 69.9	6,500
Any breastfeeding, 1 month after delivery	83.1	79.3 - 86.9	23,900	88.0 *	80.0 - 96.0	3,000	81.5 72.	.7 - 90.2	5,000	84.4	77.4 - 91.5	6,400	81.3	74.7 - 88.0	8,300
Exclusive breastfeeding, 1 month after delivery		36.7 - 46.9	11,700	41.5	26.8 - 56.1	1,300	43.8 33.	.3 - 54.3	2,700	41.9	32.3 - 51.6	3,100	41.7	32.7 - 50.6	4,100
Any breastfeeding, 3 months after delivery		57.4 - 69.0	14,000	_	66.8 - 92.5	1,900	60.5 47.		2,600	_	53.9 - 75.6	4,000		48.1 - 68.6	4,800
Exclusive breastfeeding, 3 months after delivery		20.8 - 31.0	5,700	227	16.2 - 49.2	800	34.2 22.	2 - 462	1,500		17.0 - 35.5	1,600		10.7 - 26.8	1,500



MIHA Data Snapshot, Riverside County by Neighborhood Poverty, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Riverside Cou	nty	0-4.9% of Reside		5-9.9% of Reside		10-19.9% of Residual		≥ 20% of Reside	
	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N
Service Utilization and Health Insurance Coverage										
Had a usual source of pre-pregnancy care	67.4 62.7 - 72.1	19,900	71.5 58.7 - 84.3	2,500	73.4 64.1 - 82.7	4,600	65.8 56.2 - 75.3	5,100	62.6 54.0 - 71.3	6,600
Initiated prenatal care in 1st trimester	85.7 82.2 - 89.1	24,700	84.3 * 72.8 - 95.7	2,800	91.1 * 85.4 - 96.8	5,700	89.9 84.6 - 95.2	6,900	80.7 73.9 - 87.6	8,200
Mom or infant needed but couldn't afford care postpartun	17.5 13.6 - 21.5	5,100	15.4 * 3.9 - 26.9	500	13.2 * 5.4 - 21.0	800	15.1 8.7 - 21.5	1,200	22.2 14.5 - 29.8	2,300
Participated in WIC during pregnancy†	58.3 54.7 - 61.9	17,000	33.9 20.2 - 47.5	1,100	43.5 33.0 - 54.1	2,700	59.9 50.8 - 69.0	4,700	71.8 63.9 - 79.7	7,500
Received CalFresh (food stamps) during pregnancy	27.1 22.8 - 31.4	8,000	13.2 * 2.2 - 24.2	500	9.2 * 3.2 - 15.3	600	29.8 20.4 - 39.2	2,300	40.2 31.7 - 48.7	4,300
Pre-pregnancy insurance										
Medi-Cal	24.9 20.8 - 29.1	7,400	9.2 * 1.7 - 16.8	300	9.9 * 3.8 - 16.0	600	27.5 18.6 - 36.3	2,100	36.8 28.2 - 45.4	3,800
Uninsured	28.0 23.5 - 32.5	8,300	14.4 * 4.5 - 24.2	500	21.3 11.8 - 30.8	1,300	31.3 22.0 - 40.7	2,400	32.9 24.5 - 41.2	3,400
Private	41.9 37.7 - 46.1	12,400	63.7 50.4 - 76.9	2,200	67.8 57.7 - 77.9	4,300	34.8 26.1 - 43.5	2,700	25.7 17.7 - 33.6	2,700
Prenatal insurance										
Medi-Cal	54.1 50.3 - 58.0	16,100	23.6 11.5 - 35.7	800	37.2 26.9 - 47.5	2,300	56.5 47.4 - 65.6	4,400	70.6 62.6 - 78.6	7,500
Uninsured	1.7 * 0.4 - 3.0	500								
Private	38.2 34.4 - 42.0	11,300	56.6 42.7 - 70.6	1,900	57.4 47.0 - 67.9	3,600	36.1 27.4 - 44.8	2,800	23.3 15.8 - 30.8	2,500
Maternal postpartum insurance										
Medi-Cal	38.8 34.4 - 43.3	11,400	16.6 * 5.8 - 27.3	600	22.7 13.6 - 31.7	1,400	42.4 32.9 - 51.8	3,300	52.7 44.0 - 61.4	5,600
Uninsured	16.1 12.2 - 20.0	4,700	11.7 * 2.0 - 21.3	400	11.2 * 3.3 - 19.2	700	12.8 6.0 - 19.5	1,000	21.8 14.1 - 29.5	2,300
Private	40.4 36.3 - 44.5	11,900	58.2 43.9 - 72.4	2,000	64.9 54.6 - 75.3	4,000	38.4 29.3 - 47.6	3,000	22.5 15.0 - 30.1	2,400
Infant health insurance										
Medi-Cal	58.9 54.8 - 63.0	16,900	37.3 22.8 - 51.7	1,200	41.0 30.5 - 51.5	2,500	59.7 50.5 - 69.0	4,500	74.1 66.0 - 82.3	7,700
Uninsured	2.8 1.2 - 4.3	800					4.0 * 0.5 - 7.4	300	2.6 * 0.1 - 5.1	300
Private	32.8 28.9 - 36.6	9,400	47.3 32.9 - 61.6	1,600	52.6 42.1 - 63.0	3,200	30.3 21.7 - 38.9	2,300	19.0 11.7 - 26.4	2,000
Maternal Demographics										
Age										
15-19	6.7 4.2 - 9.2	2,000			7.5 * 1.0 - 13.9	500	4.4 * 1.6 - 7.3	300	9.9 4.8 - 15.1	1,100
20-34	78.3 73.9 - 82.7	23,300	69.0 55.5 - 82.4	2,400	74.8 64.6 - 84.9	4,700	80.4 72.4 - 88.5	6,300	78.7 71.3 - 86.1	8,400
35+	15.0 11.2 - 18.9	4,500	27.9 14.7 - 41.1	1,000	17.8 8.9 - 26.6	1,100	15.2 7.4 - 22.9	1,200	11.3 5.3 - 17.4	1,200
Did not complete high school (or GED)	16.1 12.2 - 20.0	4,700	11.8 * 0.9 - 22.7	400	13.9 * 5.6 - 22.2	900	11.2 4.9 - 17.5	900	20.9 13.3 - 28.5	2,100
Unmarried	44.1 39.4 - 48.9	13,000	32.7 19.1 - 46.2	1,100	27.1 17.2 - 37.0	1,700	44.3 34.6 - 54.0	3,500	57.0 48.2 - 65.8	6,000
Race/Ethnicity										
White	27.5 23.2 - 31.8	8,100	52.3 38.1 - 66.5	1,800	33.0 23.2 - 42.9	2,000	27.3 19.0 - 35.7	2,100	17.7 10.6 - 24.8	1,900
Hispanic	59.7 55.1 - 64.2	17,500	24.4 12.1 - 36.8	800	53.5 42.6 - 64.4	3,300	60.8 51.7 - 69.9	4,700	71.4 63.5 - 79.3	7,600
Asian/Pacific Islander	6.5 3.6 - 9.4	1,900	17.4 * 5.8 - 29.0	600	9.3 * 1.8 - 16.9	600	3.9 * 0.9 - 7.0	300		
Black	5.9 5.1 - 6.7	1,700	5.9 * 2.2 - 9.7	200	3.9 1.7 - 6.1	200	7.7 4.8 - 10.6	600	6.1 4.3 - 8.0	700
Born outside the U.S.	28.4 23.6 - 33.3	8,500	22.8 10.2 - 35.3	800	24.2 13.9 - 34.4	1,500	26.1 16.7 - 35.5	2,000	31.2 22.6 - 39.7	3,300
Speaks non-English language at home	19.2 15.1 - 23.3	5,700	16.5 * 4.9 - 28.2	600	13.9 * 5.4 - 22.4	900	16.4 8.4 - 24.4	1,300	21.9 15.0 - 28.7	2,300
Income as a percent of the Federal Poverty Guideline										
0-100%	44.0 39.3 - 48.7	12,200	25.4 12.0 - 38.8	800	18.6 9.8 - 27.4	1,100	49.4 39.6 - 59.2	3,700	58.5 49.4 - 67.5	5,700
101-200%	25.5 20.9 - 30.1	7,100	15.0 * 4.5 - 25.4	500	34.5 23.6 - 45.4	2,100	30.9 21.5 - 40.4	2,300	21.3 14.3 - 28.4	2,100
> 200%	30.5 26.8 - 34.2	8,400	59.6 45.3 - 74.0	1,900	46.9 36.1 - 57.7	2,800	19.7 13.1 - 26.3	1,500	20.2 12.2 - 28.2	2,000

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Riverside County was 610. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. Neighborhood poverty is the percentage of residents with incomes below poverty in a woman's census tract. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

^{*} Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



MIHA Data Snapshot, Riverside County by Prenatal Health Insurance, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		Riverside Count	у		Medi-Cal			Private	
	%	95% CI	N	%	95% CI	N	%	95% CI	N
Total	100		29,700	54.1	50.3 - 58.0	16,100	38.	2 34.4 - 42.0	11,300
Health Status before Pregnancy									
In good to excellent health	92.3	89.6 - 94.9	27,300	88.9	84.5 - 93.4	14,200	96.	4 * 93.7 - 99.1	10,900
Chronic conditions									
Diabetes†	2.2 *	0.7 - 3.8	700	3.6	* 0.9 - 6.3	600	-		
Hypertension†	2.6	1.3 - 3.9	800	3.4	* 1.4 - 5.5	500	1.		200
Asthma†	8.2	5.6 - 10.8	2,400	9.2	5.4 - 13.0	1,500	7.	3 3.6 - 11.0	800
Nutrition and Weight									
Daily folic acid use, month before pregnancy	33.4	28.6 - 38.2	9,900	22.5	16.5 - 28.5	3,600	49.	-	5,500
Overweight before pregnancy	25.9	21.2 - 30.6	7,400	27.1	20.2 - 33.9	4,100	24.	2 17.2 - 31.2	2,700
Obese before pregnancy	25.4	20.9 - 29.9	7,300	27.7	21.1 - 34.3	4,200	24.	5 17.6 - 31.4	2,800
Inadequate weight gain during pregnancy	18.8	14.3 - 23.3	4,700	19.2	12.8 - 25.6	2,500	17.	0 10.5 - 23.6	1,600
Excessive weight gain during pregnancy	41.0	35.6 - 46.4	10,200	37.1	29.3 - 44.8	4,900	45.	9 37.5 - 54.2	4,400
Food insecurity during pregnancy	21.7	17.5 - 26.0	6,400	32.0	25.3 - 38.8	5,100	7.	9 3.6 - 12.3	900
Intimate Partner Violence (IPV) and Depressive Symptoms									
Physical or psychological IPV during pregnancy	5.9	3.7 - 8.2	1,800	9.2	5.3 - 13.1	1,500	-	-	
Prenatal depressive symptoms	13.1	9.8 - 16.4	3,900	16.1	11.2 - 21.0	2,600	10.	1 5.2 - 15.0	1,100
Postpartum depressive symptoms	16.0	12.3 - 19.7	4,700	19.9	14.1 - 25.7	3,200	13.	3 8.5 - 18.1	1,500
Hardships and Support during Pregnancy									
Experienced two or more hardships during childhood	25.6	21.2 - 30.0	7,400	31.3	24.6 - 38.1	4,900	19.	1 13.4 - 24.9	2,100
Homeless or did not have a regular place to sleep	2.9	1.3 - 4.5	900	4.0		600			
Moved due to problems paying rent or mortgage	7.9	5.2 - 10.6	2,300	11.4	7.0 - 15.8	1,800	4.	2 * 0.9 - 7.5	500
Woman or partner lost job	17.6	13.7 - 21.6	5,200	22.9	16.9 - 28.9	3,600	9.	6 4.8 - 14.4	1,100
Woman or partner had pay or hours cut back	10.5	7.6 - 13.4	3,100	13.2	8.7 - 17.8	2,100	7.	1 3.6 - 10.7	800
Became separated or divorced	9.4	6.5 - 12.2	2,800	14.6	9.7 - 19.4	2,300	3.	1 * 0.7 - 5.5	300
Had no practical or emotional support	3.5	1.7 - 5.2	1,000	3.5	* 1.2 - 5.8	600	3.	4 * 0.3 - 6.5	400
Substance Use									
Any smoking, 3 months before pregnancy	10.1	7.1 - 13.0	3,000	13.6	8.7 - 18.5	2,100	6.	4 2.9 - 10.0	700
Any smoking, 3rd trimester†	1.9 *	0.6 - 3.1	600	2.6	* 0.6 - 4.7	400	-		
Any smoking, postpartum	4.4	2.3 - 6.4	1,300	5.8	2.5 - 9.0	900	3.	0 * 0.2 - 5.7	300
Any binge drinking, 3 months before pregnancy	15.4	11.6 - 19.1	4,500	14.4	9.1 - 19.8	2,300	16.		1,800
Any alcohol use, 3rd trimester†	3.0	1.5 - 4.5	900	1.6	* 0.3 - 2.8	200	4.	2 * 1.5 - 6.9	500
Pregnancy Intention and Family Planning									
Mistimed or unwanted pregnancy	29.2	24.6 - 33.8	8,600	36.7	29.8 - 43.6	5,800	20.	8 14.8 - 26.9	2,400
Unsure of pregnancy intentions	16.1	12.2 - 20.0	4,700	17.2	11.8 - 22.5	2,700	12.		1,400
Postpartum birth control use†	80.5	76.5 - 84.6	23,500	80.0	74.3 - 85.7	12,400	82.		9,300
Infant Sleep and Breastfeeding									
Placed infant on back to sleep	79.5	75.4 - 83.6	23,000	75.6	69.4 - 81.8	11,700	82.	7 76.6 - 88.8	9,200
Infant always or often shared bed	34.8	29.9 - 39.6	10,000	40.7	33.5 - 47.9	6,300	28.		3,200
Intended to breastfeed, before birth	94.2	92.1 - 96.3	27,500	95.0	92.5 - 97.6	14,900	95.		10,600
Intended to breastfeed exclusively, before birth	66.1	61.3 - 70.9	19,300	60.1	53.0 - 67.3	9,400	72.		8.100
Breastfeeding practices	00.1	02.0 ,0.9	13,300	00.1	55.5 07.5	3,400	72.	, 55.5 ,5.5	0,100
Any breastfeeding, 1 month after delivery	83.1	79.3 - 86.9	23,900	79.4	73.5 - 85.4	12,300	87.	7 83.2 - 92.2	9,700
Exclusive breastfeeding, 1 month after delivery	41.8	36.7 - 46.9	11,700	37.9	30.9 - 44.9	5,600	45.		5,000
Any breastfeeding, 3 months after delivery	63.2	57.4 - 69.0	14,000	59.4	51.2 - 67.7	7,300	68.		5,500
Exclusive breastfeeding, 3 months after delivery	25.9	20.8 - 31.0	5,700	22.2	15.5 - 29.0	2,600	28.		2,300
Exclasive breastreeding, 5 months after delivery	23.9	20.0 31.0	3,700	22.2	13.3 - 23.0	2,000	20.	U 20.5 - 37.0	2,300



MIHA Data Snapshot, Riverside County by Prenatal Health Insurance, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Riverside County				Medi-Cal		Private			
	%	95% CI	N	%	95% CI	N	%	95% CI	N	
Service Utilization and Health Insurance Coverage										
Had a usual source of pre-pregnancy care	67.4	62.7 - 72.1	19,900	55.6	48.4 - 62.9	8,800	84.0	78.5 - 89.6	9,500	
Initiated prenatal care in 1st trimester	85.7	82.2 - 89.1	24,700	81.5	76.2 - 86.7	12,700	91.3	86.7 - 95.9	10,000	
Mom or infant needed but couldn't afford care postpartum	17.5	13.6 - 21.5	5,100	23.2	17.0 - 29.3	3,700	10.5	5.6 - 15.5	1,200	
Participated in WIC during pregnancy†	58.3	54.7 - 61.9	17,000	85.1	80.9 - 89.3	13,600	20.9	14.8 - 27.0	2,300	
Received CalFresh (food stamps) during pregnancy	27.1	22.8 - 31.4	8,000	46.5	39.3 - 53.6	7,500	2.9 *	0.3 - 5.6	300	
Pre-pregnancy insurance										
Medi-Cal	24.9	20.8 - 29.1	7,400	46.1	39.0 - 53.1	7,300				
Uninsured	28.0	23.5 - 32.5	8,300	43.6	36.5 - 50.8	6,900	4.2 *	1.0 - 7.4	500	
Private	41.9	37.7 - 46.1	12,400	7.8	3.7 - 11.9	1,200	95.4 *	92.1 - 98.6	10,800	
Maternal postpartum insurance										
Medi-Cal	38.8	34.4 - 43.3	11,400	68.6	61.7 - 75.5	10,800	2.4 *	0.7 - 4.0	300	
Uninsured	16.1	12.2 - 20.0	4,700	24.7	18.3 - 31.0	3,900				
Private	40.4	36.3 - 44.5	11,900	5.4 *	1.7 - 9.0	800	95.2 *	92.2 - 98.2	10,800	
Infant health insurance										
Medi-Cal	58.9	54.8 - 63.0	16,900	93.6 *	89.9 - 97.4	14,400	16.8	10.9 - 22.7	1,900	
Uninsured	2.8	1.2 - 4.3	800				4.1 *	1.2 - 7.1	500	
Private	32.8	28.9 - 36.6	9,400				78.2	71.8 - 84.7	8,600	
Maternal Demographics										
Age										
15-19	6.7	4.2 - 9.2	2,000	10.8	6.4 - 15.2	1,700	1.1 *	0.1 - 2.0	100	
20-34	78.3	73.9 - 82.7	23,300	80.4	74.6 - 86.2	12,900	75.4	68.0 - 82.8	8,600	
35+	15.0	11.2 - 18.9	4,500	8.8	4.4 - 13.2	1,400	23.5	16.1 - 30.9	2,700	
Did not complete high school (or GED)	16.1	12.2 - 20.0	4,700	25.6	19.1 - 32.1	4,000				
Unmarried	44.1	39.4 - 48.9	13,000	63.6	56.5 - 70.7	10,100	20.1	14.0 - 26.2	2,300	
Race/Ethnicity										
White	27.5	23.2 - 31.8	8,100	17.0	11.7 - 22.4	2,700	43.4	35.5 - 51.2	4,800	
Hispanic	59.7	55.1 - 64.2	17,500	72.9	67.2 - 78.7	11,600	40.1	32.4 - 47.9	4,500	
Asian/Pacific Islander	6.5	3.6 - 9.4	1,900	2.2 *	0.2 - 4.1	300	12.3	5.7 - 18.8	1,400	
Black	5.9	5.1 - 6.7	1,700	7.4	5.9 - 8.9	1,200	3.9	2.6 - 5.1	400	
Born outside the U.S.	28.4	23.6 - 33.3	8,500	34.5	27.5 - 41.5	5,500	19.5	12.1 - 27.0	2,200	
Speaks non-English language at home	19.2	15.1 - 23.3	5,700	25.7	19.4 - 32.1	4,100	9.9	5.0 - 14.7	1,100	
Lives in a high poverty neighborhood	37.7	32.6 - 42.7	10,600	49.8	42.6 - 57.1	7,500	22.8	15.3 - 30.4	2,500	
Income as a percent of the Federal Poverty Guideline										
0-100%	44.0	39.3 - 48.7	12,200	73.5	66.9 - 80.1	11,000	6.1 *	2.2 - 10.1	700	
101-200%	25.5	20.9 - 30.1	7,100	23.7	17.4 - 30.1	3,500	25.9	18.8 - 32.9	2,800	
> 200%	30.5	26.8 - 34.2	8,400	2.8 *	0.1 - 5.4	400	68.0	60.6 - 75.4	7,300	

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Riverside County was 610. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

^{*} Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%)

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



MIHA Data Snapshot, Riverside County by Race/Ethnicity, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Riverside County		Hispanic		Black		White		Asian/Pacific Islander		
	% 95%	6 CI N	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N	
Total	100	29,700	59.7 55.1 - 64.2	17,500	5.9 5.1 - 6.7	1,700	27.5 23.2 - 31.8	8,100	6.5 3.6 - 9.4	1,900	
Health Status before Pregnancy											
In good to excellent health	92.3 89.6 -	94.9 27,300	89.6 85.5 - 93.7	15,600	91.8 87.1 - 96.6	1,600					
Chronic conditions											
Diabetes†	2.2 * 0.7 -	3.8 700	2.7 * 0.3 - 5.1	500							
Hypertension†	2.6 1.3 -	3.9 800	2.0 * 0.3 - 3.7	300	9.4 * 2.7 - 16.0	200	2.8 * 0.2 - 5.4	200			
Asthma†	8.2 5.6 -	10.8 2,400	6.0 2.7 - 9.3	1,000	12.4 5.6 - 19.2	200	12.7 7.1 - 18.4	1,000			
Nutrition and Weight											
Daily folic acid use, month before pregnancy	33.4 28.6 -	38.2 9,900	29.4 23.1 - 35.7	5,100	21.0 12.7 - 29.3	400	40.7 31.6 - 49.7	3,300	53.8 31.4 - 76.1	1,000	
Overweight before pregnancy	25.9 21.2 -	30.6 7,400	26.6 20.3 - 33.0	4,400	26.9 17.3 - 36.4	500	20.7 12.9 - 28.5	1,700	36.9 * 13.5 - 60.4	700	
Obese before pregnancy	25.4 20.9 -	29.9 7,300	28.6 22.2 - 35.1	4,700	26.2 16.8 - 35.6	400	23.3 15.4 - 31.1	1,900	10.4 * 0.2 - 20.6	200	
Inadequate weight gain during pregnancy	18.8 14.3 -	23.3 4,700	22.0 15.5 - 28.5	3,200	11.8 * 4.5 - 19.1	200	12.6 5.9 - 19.3	900	26.4 * 6.7 - 46.2	400	
Excessive weight gain during pregnancy	41.0 35.6 -	46.4 10,200	38.1 30.6 - 45.5	5,500	56.6 44.6 - 68.5	800	45.3 35.6 - 55.0	3,200	30.6 * 12.0 - 49.3	500	
Food insecurity during pregnancy	21.7 17.5 -	26.0 6,400	24.4 18.3 - 30.4	4,300	19.4 11.3 - 27.6	300	18.9 11.3 - 26.5	1,500	15.4 * 0.3 - 30.5	300	
Intimate Partner Violence (IPV) and Depressive Symptoms											
Physical or psychological IPV during pregnancy	5.9 3.7 -	8.2 1,800	5.5 2.5 - 8.4	1,000	12.1 6.0 - 18.1	200	6.3 * 1.3 - 11.3	500			
Prenatal depressive symptoms	13.1 9.8 -	16.4 3,900	12.2 7.7 - 16.6	2,100	30.6 21.4 - 39.8	500	13.4 6.7 - 20.1	1,100			
Postpartum depressive symptoms	16.0 12.3 -	19.7 4,700	17.9 12.5 - 23.3	3,100	24.2 15.1 - 33.3	400	12.5 6.9 - 18.0	1,000	9.1 * 1.0 - 17.3	200	
Hardships and Support during Pregnancy											
Experienced two or more hardships during childhood	25.6 21.2 -	30.0 7,400	25.2 19.1 - 31.3	4,300	31.2 21.9 - 40.5	500	27.1 18.7 - 35.5	2,100	16.2 * 1.0 - 31.4	300	
Homeless or did not have a regular place to sleep	2.9 1.3 -		1.5 * 0.2 - 2.8	300	9.9 * 2.9 - 16.9	200					
Moved due to problems paying rent or mortgage	7.9 5.2 -	10.6 2,300	8.2 4.4 - 12.1	1,400	16.1 7.8 - 24.4	300	6.9 * 2.2 - 11.7	600			
Woman or partner lost job	17.6 13.7 -	21.6 5,200	21.2 15.4 - 27.1	3,700	20.1 12.1 - 28.1	300	11.6 5.8 - 17.4	900			
Woman or partner had pay or hours cut back	10.5 7.6 -	13.4 3,100	11.1 7.0 - 15.2	1,900	16.5 8.7 - 24.4	300	8.8 4.3 - 13.3	700			
Became separated or divorced	9.4 6.5 -	12.2 2,800	8.9 5.2 - 12.5	1,600	18.3 10.3 - 26.3	300	10.3 * 4.0 - 16.6	800			
Had no practical or emotional support	3.5 1.7 -	5.2 1,000	2.8 * 0.7 - 4.9	500	6.5 * 0.6 - 12.3	100					
Substance Use											
Any smoking, 3 months before pregnancy	10.1 7.1 -	13.0 3,000	8.1 4.2 - 12.0	1,400	15.9 8.4 - 23.5	300	15.7 9.1 - 22.2	1,200			
Any smoking, 3rd trimester†	1.9 * 0.6 -			,			5.4 * 1.3 - 9.5	400			
Any smoking, postpartum	4.4 2.3 -	6.4 1,300	2.4 * 0.1 - 4.6	400	6.8 * 2.0 - 11.7	100	9.5 * 3.9 - 15.1	800			
Any binge drinking, 3 months before pregnancy	15.4 11.6 -	19.1 4,500	15.1 10.1 - 20.2	2,600	9.7 * 3.1 - 16.3	200	19.8 12.2 - 27.4	1,500			
Any alcohol use, 3rd trimester†	3.0 1.5 -	4.5 900	2.3 * 0.5 - 4.1	400	5.2 * 1.4 - 9.1	<100	4.9 * 1.2 - 8.6	400			
Pregnancy Intention and Family Planning											
Mistimed or unwanted pregnancy	29.2 24.6 -	33.8 8,600	31.0 24.5 - 37.4	5,400	48.8 38.4 - 59.2	800	23.0 14.9 - 31.1	1,900	20.6 * 4.3 - 36.9	400	
Unsure of pregnancy intentions	16.1 12.2 -	20.0 4,700	18.1 12.6 - 23.6	3,200	20.4 12.0 - 28.9	400	12.6 5.7 - 19.5	1,000			
Postpartum birth control use†	80.5 76.5 -		81.9 76.5 - 87.3	14,100	74.9 65.2 - 84.6	1,300	83.2 75.6 - 90.7	6,700	65.8 * 45.5 - 86.1	1,200	
Infant Sleep and Breastfeeding											
Placed infant on back to sleep	79.5 75.4 -	83.6 23,000	84.5 79.4 - 89.5	14,500	57.7 46.7 - 68.6	1,000	77.4 69.0 - 85.7	6,100	63.6 * 41.8 - 85.3	1,100	
Infant always or often shared bed	34.8 29.9 -		36.1 29.4 - 42.8	6,200	46.8 36.0 - 57.5	800	29.4 20.8 - 37.9	2,300	31.8 * 11.4 - 52.1	500	
Intended to breastfeed, before birth	94.2 92.1 -		93.9 91.0 - 96.9	16,300	91.3 * 85.9 - 96.7	1,600	94.1 * 90.1 - 98.0	7,500			
Intended to breastfeed exclusively, before birth	66.1 61.3 -		57.9 51.0 - 64.8	10,100	68.5 58.6 - 78.4	1,200	84.1 77.9 - 90.4	6,700	64.9 * 44.1 - 85.7	1,100	
Breastfeeding practices		,		•		-				•	
Any breastfeeding, 1 month after delivery	83.1 79.3 -	86.9 23,900	83.2 77.8 - 88.5	14,200	76.2 66.9 - 85.4	1,300	81.6 74.9 - 88.4	6,400			
Exclusive breastfeeding, 1 month after delivery	41.8 36.7 -	46.9 11,700	37.3 30.6 - 44.0	6,200	36.5 25.9 - 47.1	600	49.6 40.1 - 59.1	3,800	45.6 21.7 - 69.5	700	
Any breastfeeding, 3 months after delivery	63.2 57.4 -		61.9 54.2 - 69.5	8,400	50.7 38.2 - 63.1	600	64.9 53.7 - 76.1	3,700			
Exclusive breastfeeding, 3 months after delivery	25.9 20.8 -	31.0 5,700	20.6 14.4 - 26.8	2,800	20.7 11.3 - 30.1	300	38.4 27.4 - 49.4	2,200	28.7 * 2.6 - 54.7	300	



MIHA Data Snapshot, Riverside County by Race/Ethnicity, 2013-2015

Maternal and Infant Health Assessment (MIHA) Survey

	Riverside County				Hispanic			Black			White			Asian/Pacific Islander	
	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N
Service Utilization and Health Insurance Coverage															
Had a usual source of pre-pregnancy care	67.4	62.7 - 72.1	19,900	64.9	58.3 - 71.5	11,300	68.8	58.4 - 79.2	1,200	72.4	63.7 - 81.0	5,800	63.6	43.4 - 83.8	1,200
Initiated prenatal care in 1st trimester	85.7	82.2 - 89.1	24,700	86.6	82.0 - 91.2	14,700	77.4	67.7 - 87.0	1,300	86.0	79.3 - 92.7	6,800	82.1	66.2 - 98.0	1,600
Mom or infant needed but couldn't afford care postpartum	17.5	13.6 - 21.5	5,100	18.3	12.9 - 23.8	3,200	11.4	5.8 - 17.0	200	16.6	9.6 - 23.6	1,300	23.1 '	* 2.9 - 43.3	400
Participated in WIC during pregnancy†	58.3	54.7 - 61.9	17,000	71.1	65.6 - 76.6	12,300	72.7	65.8 - 79.6	1,200	36.0	27.6 - 44.4	2,900	29.8	* 9.9 - 49.8	500
Received CalFresh (food stamps) during pregnancy	27.1	22.8 - 31.4	8,000	30.4	24.0 - 36.8	5,300	51.5	41.7 - 61.4	900	19.6	11.8 - 27.4	1,600			
Pre-pregnancy insurance															
Medi-Cal	24.9	20.8 - 29.1	7,400	30.5	24.2 - 36.8	5,300	43.3	33.0 - 53.5	700	14.5	7.7 - 21.3	1,200			
Uninsured	28.0	23.5 - 32.5	8,300	33.7	27.0 - 40.3	5,800	26.0	17.0 - 34.9	400	20.9	13.2 - 28.5	1,700			
Private	41.9	37.7 - 46.1	12,400	30.0	24.2 - 35.8	5,200	23.6	15.8 - 31.5	400	61.6	52.8 - 70.3	5,000	80.0	65.7 - 94.3	1,500
Prenatal insurance															
Medi-Cal	54.1	50.3 - 58.0	16,100	66.2	60.3 - 72.1	11,600	67.8	59.7 - 76.0	1,200	33.7	25.3 - 42.1	2,700	18.0	* 2.5 - 33.6	300
Uninsured	1.7 *	0.4 - 3.0	500	2.0	0.2 - 3.9	400									
Private	38.2	34.4 - 42.0	11,300	25.5	20.3 - 30.7	4,500	24.8	17.0 - 32.6	400	60.0	51.4 - 68.7	4,800	71.6	* 53.3 - 90.0	1,400
Maternal postpartum insurance															
Medi-Cal	38.8	34.4 - 43.3	11,400	42.9	36.2 - 49.6	7,400	65.2	56.3 - 74.1	1,100	29.3	21.0 - 37.7	2,300	17.1 °	* 1.7 - 32.5	300
Uninsured	16.1	12.2 - 20.0	4,700	22.8	16.7 - 28.9	3,900	5.7	* 1.3 - 10.1	<100	7.8	2.8 - 12.7	600			
Private	40.4	36.3 - 44.5	11,900	29.7	23.9 - 35.6	5,100	23.0	15.6 - 30.4	400	59.1	50.2 - 67.9	4,700	71.6	* 53.3 - 90.0	1,400
Infant health insurance															
Medi-Cal	58.9	54.8 - 63.0	16,900	70.9	65.0 - 76.8	12,100	73.7	65.9 - 81.6	1,200	35.5	26.7 - 44.3	2,800	34.1	11.9 - 56.2	600
Uninsured	2.8	1.2 - 4.3	800	2.9	0.7 - 5.0	500									
Private	32.8	28.9 - 36.6	9,400	20.4	15.4 - 25.4	3,500	21.3	14.1 - 28.5	400	56.4	47.4 - 65.4	4,400	56.2	32.9 - 79.5	900
Maternal Demographics															
Age															
15-19	6.7	4.2 - 9.2	2,000	9.7	5.7 - 13.8	1,700	5.9	* 0.7 - 11.2	100						
20-34	78.3	73.9 - 82.7	23,300	77.0	71.0 - 83.0	13,500	77.6	68.8 - 86.4	1,400	83.1	75.2 - 90.9	6,700	64.9	43.8 - 85.9	1,200
35+	15.0	11.2 - 18.9	4,500	13.2	8.2 - 18.3	2,300	16.5	8.9 - 24.0	300	15.4	7.8 - 23.0	1,200	32.2	11.6 - 52.8	600
Did not complete high school (or GED)	16.1	12.2 - 20.0	4,700	21.4	15.5 - 27.4	3,700	12.8	* 4.7 - 20.8	200	7.6	2.5 - 12.7	600			
Unmarried	44.1	39.4 - 48.9	13,000	50.6	43.7 - 57.5	8,800	76.6	68.8 - 84.3	1,300	31.8	22.9 - 40.7	2,600			
Born outside the U.S.	28.4	23.6 - 33.3	8,500	36.6	29.8 - 43.4	6,400	5.2	* 0.2 - 10.1	<100	4.2	0.4 - 8.1	300	77.3 *	* 60.9 - 93.7	1,500
Speaks non-English language at home	19.2	15.1 - 23.3	5,700	28.0	21.8 - 34.2	4,900							39.3	17.9 - 60.8	800
Lives in a high poverty neighborhood	37.7	32.6 - 42.7	10,600	46.3	39.2 - 53.4	7,600	38.6	28.2 - 48.9	700	24.0	14.9 - 33.2	1,900			
Income as a percent of the Federal Poverty Guideline															
0-100%	44.0	39.3 - 48.7	12,200	54.9	48.0 - 61.9	8,800	57.4	47.8 - 67.1	1,000	25.3	17.0 - 33.7	2,000	18.2	* 1.7 - 34.7	300
101-200%	25.5	20.9 - 30.1	7,100	29.1	22.5 - 35.7	4,700	20.1	11.4 - 28.8	300	22.4	14.7 - 30.1	1,800	15.7	* 2.5 - 28.9	300
> 200%	30.5	26.8 - 34.2	8,400	15.9	11.5 - 20.3	2,600	22.4	15.3 - 29.5	400	52.3	43.3 - 61.3	4,100	66.1	45.5 - 86.6	1,100

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Riverside County was 610. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

^{*} Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



MIHA Data Snapshot, Riverside County by Total Live Births, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		Riverside County		First Liv	e Birth		Seco	Second Live Birth or More			
	%	95% CI	N	%	95%	CI	N	%	95% CI	N	
Total	100		29,700	35.4	30.4 -	40.3	10,500	64.6	59.7 - 69.6	19,200	
Health Status before Pregnancy											
In good to excellent health Chronic conditions	92.3	89.6 - 94.9	27,300	90.8	86.2 -	95.3	9,500	93.1	89.9 - 96.4	17,800	
Diabetes†	2.2 *	0.7 - 3.8	700	3.7 *	0.6 -	6.8	400				
Hypertension†	2.6	1.3 - 3.9	800	3.1 *	0.7 -	5.5	300	2.3 *	0.7 - 3.8	400	
Asthma†	8.2	5.6 - 10.8	2,400	9.6	5.4 -	13.9	1,000	7.4	4.1 - 10.6	1,400	
Nutrition and Weight											
Daily folic acid use, month before pregnancy	33.4	28.6 - 38.2	9,900	37.6	29.2 -	46.0	4,000	31.1	25.2 - 37.0	6,000	
Overweight before pregnancy	25.9	21.2 - 30.6	7,400	22.9	15.8 -	30.0	2,300	27.5	21.5 - 33.6	5,100	
Obese before pregnancy	25.4	20.9 - 29.9	7,300	15.4	9.4 -	21.3	1,600	30.9	24.9 - 36.9	5,700	
Inadequate weight gain during pregnancy	18.8	14.3 - 23.3	4,700	14.2	7.7 -	20.7	1,300	21.4	15.5 - 27.3	3,400	
Excessive weight gain during pregnancy	41.0	35.6 - 46.4	10,200	47.6	38.3 -	56.9	4,300	37.3	30.7 - 43.9	6,000	
Food insecurity during pregnancy	21.7	17.5 - 26.0	6,400	20.7	13.8 -	27.6	2,200	22.3	16.8 - 27.7	4,300	
Intimate Partner Violence (IPV) and Depressive Symptoms											
Physical or psychological IPV during pregnancy	5.9	3.7 - 8.2	1,800	5.9 *	2.1 -	9.7	600	6.0	3.2 - 8.8	1,100	
Prenatal depressive symptoms	13.1	9.8 - 16.4	3,900	9.6	4.5 -	14.8	1,000	15.1	10.8 - 19.3	2,900	
Postpartum depressive symptoms	16.0	12.3 - 19.7	4,700	9.9	5.5 -	14.4	1,000	19.3	14.3 - 24.4	3,700	
Hardships and Support during Pregnancy											
Experienced two or more hardships during childhood	25.6	21.2 - 30.0	7,400	19.4	13.2 -	25.5	2,000	29.0	23.1 - 34.8	5,400	
Homeless or did not have a regular place to sleep	2.9	1.3 - 4.5	900	4.7 *	0.9 -	8.5	500	1.9 *	0.7 - 3.2	400	
Moved due to problems paying rent or mortgage	7.9	5.2 - 10.6	2,300	6.8 *	2.3 -	11.4	700	8.4	5.0 - 11.8	1,600	
Woman or partner lost job	17.6	13.7 - 21.6	5,200	14.5	8.3 -	20.6	1,500	19.4	14.2 - 24.5	3,700	
Woman or partner had pay or hours cut back	10.5	7.6 - 13.4	3,100	10.3	5.7 -	14.9	1,100	10.6	7.0 - 14.3	2,000	
Became separated or divorced	9.4	6.5 - 12.2	2,800	9.8	5.4 -	14.1	1,000	9.1	5.5 - 12.8	1,800	
Had no practical or emotional support	3.5	1.7 - 5.2	1,000					4.9	2.2 - 7.6	900	
Substance Use											
Any smoking, 3 months before pregnancy	10.1	7.1 - 13.0	3,000	10.0	5.2 -	14.7	1,000	10.1	6.3 - 13.9	1,900	
Any smoking, 3rd trimester†	1.9 *	0.6 - 3.1	600					2.1 *	0.5 - 3.8	400	
Any smoking, postpartum	4.4	2.3 - 6.4	1,300	2.3 *	0.2 -	4.3	200	5.5	2.6 - 8.4	1,100	
Any binge drinking, 3 months before pregnancy	15.4	11.6 - 19.1	4,500	21.3	14.3 -	28.3	2,100	12.2	7.9 - 16.5	2,300	
Any alcohol use, 3rd trimester†	3.0	1.5 - 4.5	900	1.7 *	0.1 -	3.4	200	3.7	1.6 - 5.7	700	
Pregnancy Intention and Family Planning											
Mistimed or unwanted pregnancy	29.2	24.6 - 33.8	8,600	33.1	24.9 -	41.3	3,500	27.0	21.5 - 32.5	5,100	
Unsure of pregnancy intentions	16.1	12.2 - 20.0	4,700	14.6	8.3 -	20.8	1,500	17.0	12.0 - 21.9	3,200	
Postpartum birth control use†	80.5	76.5 - 84.6	23,500	78.6	71.5 -	85.6	8,100	81.6	76.7 - 86.5	15,400	
Infant Sleep and Breastfeeding											
Placed infant on back to sleep	79.5	75.4 - 83.6	23,000	76.7	69.3 -	84.1	7,800	81.0	76.0 - 86.0	15,200	
Infant always or often shared bed	34.8	29.9 - 39.6	10,000	36.6	28.3 -	44.9	3,700	33.8	27.7 - 39.8	6,300	
Intended to breastfeed, before birth	94.2	92.1 - 96.3	27,500	96.5 *	93.9 -	99.0	9,800	93.1	90.1 - 96.0	17,700	
Intended to breastfeed exclusively, before birth	66.1	61.3 - 70.9	19,300	71.8	63.9 -	79.6	7,300	63.1	56.9 - 69.2	12,000	
Breastfeeding practices			•				•	-		•	
Any breastfeeding, 1 month after delivery	83.1	79.3 - 86.9	23,900	84.3	78.8 -	89.8	8,600	82.4	77.4 - 87.4	15,300	
Exclusive breastfeeding, 1 month after delivery	41.8	36.7 - 46.9	11,700	40.5	31.9 -	49.2	4,000	42.5	36.2 - 48.8	7,700	
Any breastfeeding, 3 months after delivery	63.2	57.4 - 69.0	14,000	58.8	48.8 -	68.7	4,600	65.6	58.5 - 72.6	9,400	
Exclusive breastfeeding, 3 months after delivery	25.9	20.8 - 31.0	5,700	26.5	17.4 -	35.6	2,000	25.6	19.6 - 31.7	3,600	



MIHA Data Snapshot, Riverside County by Total Live Births, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		Riverside County	1		First Live Birth		Second Live Birth or More			
	%	95% CI	N	%	95% CI	N	%	95% CI	N	
Service Utilization and Health Insurance Coverage										
Had a usual source of pre-pregnancy care	67.4	62.7 - 72.1	19,900	60.0	51.6 - 68.4	6,300	71.5	65.8 - 77.2	13,700	
Initiated prenatal care in 1st trimester	85.7	82.2 - 89.1	24,700	85.9	80.1 - 91.7	8,800	85.5	81.2 - 89.9	15,900	
Mom or infant needed but couldn't afford care postpartum	17.5	13.6 - 21.5	5,100	17.0	10.6 - 23.4	1,700	17.8	12.7 - 22.9	3,400	
Participated in WIC during pregnancy†	58.3	54.7 - 61.9	17,000	54.8	46.9 - 62.8	5,700	60.2	55.1 - 65.3	11,300	
Received CalFresh (food stamps) during pregnancy	27.1	22.8 - 31.4	8,000	19.6	12.7 - 26.5	2,100	31.2	25.5 - 36.9	6,000	
Pre-pregnancy insurance										
Medi-Cal	24.9	20.8 - 29.1	7,400	12.5	6.5 - 18.5	1,300	31.8	26.3 - 37.4	6,000	
Uninsured	28.0	23.5 - 32.5	8,300	32.5	24.7 - 40.3	3,400	25.6	19.8 - 31.3	4,900	
Private	41.9	37.7 - 46.1	12,400	48.7	40.5 - 57.0	5,100	38.1	32.7 - 43.5	7,200	
Prenatal insurance										
Medi-Cal	54.1	50.3 - 58.0	16,100	47.2	39.1 - 55.3	5,000	57.9	52.7 - 63.1	11,100	
Uninsured	1.7 *	0.4 - 3.0	500				2.6 *	0.6 - 4.5	500	
Private	38.2	34.4 - 42.0	11,300	45.8	37.8 - 53.8	4,800	34.0	29.0 - 39.1	6,500	
Maternal postpartum insurance										
Medi-Cal	38.8	34.4 - 43.3	11,400	33.2	25.4 - 41.1	3,500	41.9	36.1 - 47.8	8,000	
Uninsured	16.1	12.2 - 20.0	4,700	14.4	8.2 - 20.6	1,500	17.1	11.9 - 22.2	3,200	
Private	40.4	36.3 - 44.5	11,900	45.4	37.2 - 53.6	4,700	37.6	32.2 - 43.0	7,100	
Infant health insurance										
Medi-Cal	58.9	54.8 - 63.0	16,900	52.0	43.7 - 60.2	5,300	62.7	57.3 - 68.1	11,600	
Uninsured	2.8	1.2 - 4.3	800	3.2 *	0.3 - 6.0	300	2.5 *	0.7 - 4.4	500	
Private	32.8	28.9 - 36.6	9,400	38.4	30.5 - 46.4	3,900	29.7	24.6 - 34.7	5,500	
Maternal Demographics										
Age										
15-19	6.7	4.2 - 9.2	2,000	16.4	10.0 - 22.8	1,700	1.4 *	0.1 - 2.7	300	
20-34	78.3	73.9 - 82.7	23,300	75.8	68.0 - 83.5	8,000	79.6	74.3 - 84.9	15,300	
35+	15.0	11.2 - 18.9	4,500	7.8 *	2.6 - 13.0	800	19.0	13.8 - 24.2	3,700	
Did not complete high school (or GED)	16.1	12.2 - 20.0	4,700	16.9	9.6 - 24.2	1,800	15.6	11.0 - 20.2	2,900	
Unmarried	44.1	39.4 - 48.9	13,000	56.4	48.1 - 64.6	5,900	37.4	31.4 - 43.3	7,100	
Race/Ethnicity										
White	27.5	23.2 - 31.8	8,100	30.2	22.2 - 38.2	3,200	26.0	20.9 - 31.2	4,900	
Hispanic	59.7	55.1 - 64.2	17,500	56.2	47.8 - 64.5	5,900	61.6	55.9 - 67.3	11,600	
Asian/Pacific Islander	6.5	3.6 - 9.4	1,900	6.3 *	1.5 - 11.0	700	6.6	3.0 - 10.2	1,300	
Black	5.9	5.1 - 6.7	1,700	7.1	4.9 - 9.3	700	5.3	4.2 - 6.4	1,000	
Born outside the U.S.	28.4	23.6 - 33.3	8,500	19.8	12.7 - 27.0	2,100	33.2	26.8 - 39.5	6,400	
Speaks non-English language at home	19.2	15.1 - 23.3	5,700	12.9	7.5 - 18.3	1,400	22.7	17.1 - 28.2	4,300	
Lives in a high poverty neighborhood	37.7	32.6 - 42.7	10,600	38.7	29.7 - 47.6	3,800	37.1	30.9 - 43.4	6,800	
Income as a percent of the Federal Poverty Guideline										
0-100%	44.0	39.3 - 48.7	12,200	38.8	30.2 - 47.3	3,800	46.8	40.8 - 52.9	8,400	
101-200%	25.5	20.9 - 30.1	7,100	26.3	18.6 - 34.0	2,600	25.1	19.4 - 30.8	4,500	
> 200%	30.5	26.8 - 34.2	8,400	34.9	27.0 - 42.9	3,400	28.1	23.2 - 33.0	5,000	

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Riverside County was 610. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

^{*} Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%)

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).