

MIHA Data Snapshot, San Diego County by Education, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Sai	n Diego Cou	nty		High Schoo	I	High Sch	ool Graduat	e or GED	:	Some Colleg	е	Co	llege Gradu	ate
	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N
Total	100		43,300	11.5	8.1 - 14.9	4,900	16.3	12.5 - 20.1	7,000	33.9	28.9 - 39.0	14,500	38.3	33.2 - 43.3	16,400
Health Status before Pregnancy															
In good to excellent health	94.1	91.9 - 96.2	40,700	82.7 *	70.5 - 94.9	4,100	95.5 *	92.4 - 98.7	6,700	93.3	89.7 - 96.8	13,500			
Chronic conditions															
Diabetes†	3.5 *	1.4 - 5.7	1,500												
Hypertension†	3.3	1.4 - 5.1	1,400							3.7	* 0.2 - 7.3	500			
Asthma†	9.2	5.9 - 12.5	3,900				6.8*	1.5 - 12.1	500	9.5	4.0 - 14.9	1,400	10.8 *	4.4 - 17.1	1,800
Nutrition and Weight															
Daily folic acid use, month before pregnancy	38.6	33.3 - 43.9	16,700	22.7	9.6 - 35.8	1,100	29.6	17.6 - 41.5	2,100	34.5	25.6 - 43.4	5,000	51.9	42.5 - 61.3	8,500
Overweight before pregnancy	24.0	19.3 - 28.7	9,900	22.0 *	7.0 - 37.0	900	26.2	13.9 - 38.4	1,700	26.5	18.9 - 34.1	3,800	22.1	14.1 - 30.1	3,500
Obese before pregnancy	16.7	12.7 - 20.6	6,900	23.3 *	8.5 - 38.2	900	19.1	9.9 - 28.4	1,200	22.2	14.5 - 29.8	3,200	8.1 *	3.1 - 13.2	1,300
Inadequate weight gain during pregnancy	17.6	12.9 - 22.4	6,400	26.2 *	4.5 - 47.9	700	18.3 *	6.6 - 30.0	1,000	22.1	13.2 - 30.9	2,900	12.2	6.4 - 18.1	1,800
Excessive weight gain during pregnancy	39.5	33.7 - 45.3	14,400	48.5	27.1 - 69.8	1,300	36.0	21.5 - 50.5	2,000	38.2	29.2 - 47.1	4,900	40.4	30.6 - 50.2	5,900
Food insecurity during pregnancy	14.1	10.4 - 17.9	6,100	26.2 *	10.5 - 42.0	1,300	18.9	10.2 - 27.7	1,300	20.9	12.8 - 28.9	3,000	2.6 *	0.1 - 5.2	400
Intimate Partner Violence (IPV) and Depressive Symptoms															
Physical or psychological IPV during pregnancy	7.2	4.2 - 10.1	3,100	16.0 *	4.2 - 27.8	800	6.8 *	1.2 - 12.5	500	10.6	* 4.2 - 17.1	1,500			
Prenatal depressive symptoms	13.7	9.9 - 17.6	5,900	20.7 *	5.3 - 36.0	1,000	15.7	7.1 - 24.2	1,100	17.7	10.5 - 24.9	2,600	7.2 *	2.4 - 11.9	1,200
Postpartum depressive symptoms	13.2	9.6 - 16.8	5,700	7.5 *	1.8 - 13.1	400	16.3	6.9 - 25.7	1,100	18.3	11.1 - 25.4	2,700	9.0 *	3.6 - 14.3	1,500
Hardships and Support during Pregnancy															
Experienced two or more hardships during childhood	27.5	22.7 - 32.3	11,600	42.7	25.7 - 59.6	2,000	28.1	17.4 - 38.9	1,900	40.7	31.7 - 49.7	5,800	11.0	5.0 - 17.1	1,800
Homeless or did not have a regular place to sleep	3.7	1.7 - 5.8	1,600			_,	_	0.8 - 10.3	400	_	* 0.8 - 10.2	800			_,
Moved due to problems paying rent or mortgage	5.7	3.3 - 8.0	2,400					3.5 - 18.1	800	9.9	4.2 - 15.6	1,400	1.3 *	0.1 - 2.5	200
Woman or partner lost job	12.2	8.9 - 15.4	5,200	20.1 *	6.8 - 33.5	1,000		4.0 - 17.1	700	17.9	10.9 - 24.9	2,600	_	1.9 - 8.3	800
Woman or partner had pay or hours cut back	9.3	6.5 - 12.0	4,000	_	2.0 - 23.6	600		4.7 - 19.5	800	12.0	6.8 - 17.2	1,700	_	0.9 - 7.7	700
Became separated or divorced	6.5	3.6 - 9.5	2,800	19.7 *	5.6 - 33.8	900	3.5 *	0.8 - 6.2	200	8.7	* 2.6 - 14.8	1,200			
Had no practical or emotional support	3.8	1.9 - 5.7	1,600				7.6 *	1.6 - 13.5	500	3.0	* 0.4 - 5.6	400			
Substance Use															
Any smoking, 3 months before pregnancy	12.3	8.8 - 15.8	5,200	16.1 *	2.8 - 29.4	800	17.5	9.1 - 25.9	1,200	16.8	9.7 - 23.9	2,400	4.0 *	0.9 - 7.1	600
Any smoking, 3rd trimester†	2.2	1.0 - 3.4	900						,		* 0.6 - 5.8	500			
Any smoking, postpartum	5.5	3.1 - 7.9	2,400				7.3 *	2.3 - 12.4	500	9.1	* 3.7 - 14.6	1,300			
Any binge drinking, 3 months before pregnancy		14.3 - 22.8	7,800	16.3 *	3.8 - 28.8	700	_	2.9 - 16.2	600	-	19.3 - 35.3	4,000	14.8	8.1 - 21.6	2,400
Any alcohol use, 3rd trimester†	12.3	8.3 - 16.4	5,300							13.0	6.2 - 19.8	1,900	17.0	9.5 - 24.4	2,800
Pregnancy Intention and Family Planning															
Mistimed or unwanted pregnancy	30.4	25.5 - 35.4	13,000	43 1	26.6 - 59.5	2,100	46.5	33.7 - 59.4	3,200	31.6	23.4 - 39.8	4,500	18 5	11.2 - 25.9	2,900
Unsure of pregnancy intentions		10.3 - 18.0	6,000	_	5.9 - 32.8	900		6.7 - 22.9	1,000		10.9 - 25.6	2,600		2.6 - 13.1	1,300
Postpartum birth control use†		78.3 - 86.9	35,000		68.9 - 98.4	3,800	_	83.0 - 96.2	6,100		75.7 - 89.7	11,800	_	71.0 - 86.6	,
Infant Sleep and Breastfeeding															
Placed infant on back to sleep	79 9	75.4 - 84.3	34,000	69.7	53.4 - 86.0	3,400	73.3	60.7 - 85.9	5,000	82.3	76.0 - 88.7	11,700	83.5	76.4 - 90.7	13,500
Infant always or often shared bed		25.0 - 34.6	12,700		21.3 - 53.9	1,800		20.0 - 44.7	2,200		27.1 - 43.3	5,000		14.3 - 29.5	3,500
Intended to breastfeed, before birth		90.4 - 95.9	39,600		72.8 - 97.0	4,100		84.4 - 98.5	6,300		* 90.3 - 97.6	13,400		91.2 - 99.6	15,400
Intended to breastfeed exclusively, before birth		63.4 - 73.6	29,200		44.4 - 76.5	2,900		40.5 - 66.4	3,700		61.7 - 77.8	9,900		67.6 - 84.7	12,300
Breastfeeding practices	- 0.0		-,			,	30.1		-,	-5		-,	. 0.2		.,
Any breastfeeding, 1 month after delivery	90.2	87.1 - 93.2	38,000	85.2 *	71.8 - 98.6	4,100	86.5*	78.0 - 95.1	5,800	89.2	84.6 - 93.8	12,700	94.1 *	89.9 - 98.4	15,000
Exclusive breastfeeding, 1 month after delivery		48.7 - 59.8	22,400		44.1 - 77.5	2,800		31.1 - 56.7	2,900		44.3 - 62.5	7,300	_	48.2 - 67.0	9,200
Any breastfeeding, 3 months after delivery		69.1 - 79.9	24,300		38.6 - 76.0	2,400		51.9 - 78.6	3,700		61.9 - 80.9	7,700		82.1 - 95.3	10,400
Exclusive breastfeeding, 3 months after delivery		30.6 - 42.7	11,900		10.7 - 43.2	1,100		5.0 - 25.9	900		24.0 - 43.8	3,600		40.9 - 63.4	6,200



MIHA Data Snapshot, San Diego County by Education, 2013-2015

Maternal and Infant Health Assessment (MIHA) Survey

	Sar	n Diego Cou	nty	_ -		High School Graduate or GED		e or GED			e	College Graduate		ate	
	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N
Service Utilization and Health Insurance Coverage															
Had a usual source of pre-pregnancy care	75.4	71.0 - 79.8	32,600	71.4	57.2 - 85.6	3,500	62.8	50.1 - 75.5	4,300	73.1	65.8 - 80.4	10,600	83.9	77.1 - 90.7	13,700
Initiated prenatal care in 1st trimester	89.7	86.7 - 92.7	38,200	75.0	60.4 - 89.7	3,700	82.1	73.9 - 90.3	5,400	88.5	82.9 - 94.1	12,700	98.4 *	96.9 - 99.8	16,100
Mom or infant needed but couldn't afford care postpartum	10.2	6.8 - 13.7	4,300	15.9	3.4 - 28.4	800	10.5	* 0.8 - 20.2	700	14.7	7.9 - 21.5	2,000	4.3 *	0.6 - 8.1	700
Participated in WIC during pregnancy†	44.3	40.0 - 48.5	18,600	83.5	70.5 - 96.5	4,000	71.9	59.2 - 84.6	4,800	54.4	45.9 - 62.9	7,800	10.8	6.4 - 15.2	1,700
Received CalFresh (food stamps) during pregnancy	19.3	15.5 - 23.1	8,300	45.6	29.0 - 62.1	2,200	27.0	16.4 - 37.5	1,900	25.4	17.5 - 33.2	3,700	2.3 *	0.7 - 4.0	400
Pre-pregnancy insurance															
Medi-Cal	16.4	13.0 - 19.9	7,100	30.1	16.4 - 43.8	1,500	28.8	17.7 - 39.9	2,000	20.1	13.1 - 27.1	2,900	4.2 *	0.8 - 7.5	700
Uninsured	20.4	16.2 - 24.5	8,800	47.1	30.6 - 63.7	2,300	35.2	22.6 - 47.8	2,400	18.6	11.6 - 25.6	2,700	7.1	3.3 - 11.0	1,200
Private	50.4	45.7 - 55.1	21,700	13.5	2.5 - 24.6	700	24.5	14.1 - 35.0	1,700	42.7	34.0 - 51.3	6,200	79.5	72.5 - 86.5	13,000
Prenatal insurance															
Medi-Cal	38.5	34.5 - 42.5	16,600	75.7	61.3 - 90.1	3,700	71.3	60.2 - 82.4	5,000	41.4	33.2 - 49.7	6,000	10.4	5.8 - 15.0	1,700
Uninsured	2.7 *	0.9 - 4.5	1,200												
Private	45.8	41.2 - 50.4	19,800				13.8	6.8 - 20.8	1,000	35.5	27.1 - 43.9	5,200	79.6	72.6 - 86.5	13,000
Maternal postpartum insurance															
Medi-Cal	27.7	23.7 - 31.7	11,800	66.1	50.9 - 81.3	3,200	45.7	33.4 - 58.0	3,000	31.6	23.7 - 39.5	4,600	5.5 *	2.0 - 9.0	900
Uninsured	13.5	9.9 - 17.2	5,800	19.0	6.7 - 31.3	900	22.5	11.1 - 33.8	1,500	15.0	8.2 - 21.8	2,200	6.9 *	2.4 - 11.4	1,100
Private	46.5	41.8 - 51.1	19,800				16.1	8.0 - 24.1	1,000	35.6	27.1 - 44.1	5,200	79.0	71.7 - 86.2	12,900
Infant health insurance															
Medi-Cal	41.0	36.6 - 45.3	17,400	82.6	71.0 - 94.2	3,900	72.5	61.6 - 83.3	5,000	46.4	37.9 - 54.8	6,600	10.4	5.5 - 15.2	1,700
Uninsured	4.8	2.5 - 7.1	2,000	14.1	3.0 - 25.1	700				5.2	* 1.0 - 9.4	700			
Private	41.9	37.2 - 46.5	17,800				11.9	5.6 - 18.2	800	28.5	20.4 - 36.6	4,000	78.5	71.1 - 85.8	12,700
Maternal Demographics															
Unmarried	36.3	31.5 - 41.2	15,700	81.4	71.1 - 91.8	4,000	47.8	35.1 - 60.6	3,300	41.8	33.3 - 50.4	6,000	13.2	6.6 - 19.9	2,200
Race/Ethnicity															
White	38.6	33.2 - 44.0	15,500				20.7	9.3 - 32.1	1,300	39.4	30.3 - 48.5	5,200	52.4	42.7 - 62.0	8,100
Hispanic	43.3	38.1 - 48.6	17,400	78.4	63.8 - 93.0	3,500	56.6	43.0 - 70.1	3,600	47.2	38.1 - 56.2	6,300	24.7	16.8 - 32.5	3,800
Asian/Pacific Islander	12.4	8.3 - 16.6	5,000				15.1 '	* 2.9 - 27.3	1,000	5.8	* 1.2 - 10.3	800	20.1	11.8 - 28.3	3,100
Black	5.0	4.1 - 6.0	2,000	6.0	1.3 - 10.6	300	6.9	3.5 - 10.2	400	6.6	4.3 - 8.8	900	2.8 *	1.0 - 4.5	400
Born outside the U.S.	32.8	27.6 - 37.9	14,200	40.9	25.1 - 56.6	2,000	50.3	37.6 - 63.0	3,500	22.7	14.8 - 30.6	3,300	31.6	22.8 - 40.3	5,200
Speaks non-English language at home	26.8	21.9 - 31.6	11,600	43.9	27.8 - 60.0	2,100	46.8	33.9 - 59.6	3,300	16.3	9.7 - 23.0	2,400	22.5	14.4 - 30.5	3,700
Lives in a high poverty neighborhood	28.9	24.2 - 33.7	12,100	47.4	30.6 - 64.1	2,200	42.8	30.1 - 55.5	2,900	31.4	23.1 - 39.6	4,500	15.1	8.4 - 21.9	2,400
Income as a percent of the Federal Poverty Guideline															
0-100%		28.5 - 37.6	13,300		54.7 - 90.8	3,000		47.7 - 73.7	3,700		32.3 - 50.3	5,700	5.5	2.5 - 8.5	900
101-200%	19.6	15.3 - 23.8	7,900	25.6	7.5 - 43.7	1,100	28.2	16.0 - 40.3	1,700	23.7	16.7 - 30.7	3,200	10.4	4.7 - 16.1	1,700
> 200%	47.4	42.7 - 52.1	19,100				11.1 '	* 3.0 - 19.3	700	35.0	26.5 - 43.5	4,800	84.0	77.9 - 90.2	13,400

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of San Diego County was 602. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

^{*} Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



MIHA Data Snapshot, San Diego County by Geographical Area, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

Material and mant fleaten / 155e55ment (mm/		San Diego Count	y		Url	oan			Rural/I	Frontier	
	%	95% CI	N	%	95%	CI	N	 %	95%	CI	N
Total	100		43,300	95.3 *	92.4 -	98.1	39,800	4.7 *	1.9 -	7.6	2,000
Health Status before Pregnancy											
In good to excellent health Chronic conditions	94.1	91.9 - 96.2	40,700	94.4	92.4 -	96.5	37,600				
Diabetes†	3.5 *	1.4 - 5.7	1,500	3.5 *	1.2 -	5.7	1,400				
Hypertension [†]	3.3	1.4 - 5.1	1,400	3.6	1.5 -	5.6	1,400				
Asthma†	9.2	5.9 - 12.5	3,900	9.2	5.7 -	12.6	3,600				
Nutrition and Weight											
Daily folic acid use, month before pregnancy	38.6	33.3 - 43.9	16,700	37.5	32.1 -	43.0	14,900	46.8 *	15.8 -	77.9	900
Overweight before pregnancy	24.0	19.3 - 28.7	9,900	24.1	19.3 -	29.0	9,100				
Obese before pregnancy	16.7	12.7 - 20.6	6,900	16.6	12.6 -	20.6	6,300				
Inadequate weight gain during pregnancy	17.6	12.9 - 22.4	6,400	18.0	13.1 -	22.9	6,000				
Excessive weight gain during pregnancy	39.5	33.7 - 45.3	14,400	39.5	33.5 -	45.4	13,300	48.2 *	15.6 -	80.9	900
Food insecurity during pregnancy	14.1	10.4 - 17.9	6,100	14.5	10.5 -	18.5	5,700				
Intimate Partner Violence (IPV) and Depressive Symptoms											
Physical or psychological IPV during pregnancy	7.2	4.2 - 10.1	3,100	6.8	3.8 -	9.8	2,700				
Prenatal depressive symptoms	13.7	9.9 - 17.6	5,900	14.1	10.1 -	18.1	5,600				
Postpartum depressive symptoms	13.2	9.6 - 16.8	5,700	14.0	10.2 -	17.9	5,600				
Hardships and Support during Pregnancy											
Experienced two or more hardships during childhood	27.5	22.7 - 32.3	11,600	28.3	23.3 -	33.4	11,000				
Homeless or did not have a regular place to sleep	3.7	1.7 - 5.8	1,600	3.3 *	1.3 -	5.2	1,300				
Moved due to problems paying rent or mortgage	5.7	3.3 - 8.0	2,400	5.9	3.3 -	8.4	2,300				
Woman or partner lost job	12.2	8.9 - 15.4	5,200	11.6	8.4 -	14.8	4,600				
Woman or partner had pay or hours cut back	9.3	6.5 - 12.0	4,000	8.8	6.1 -	11.5	3,500				
Became separated or divorced	6.5	3.6 - 9.5	2,800	6.2	3.2 -	9.2	2,400				
Had no practical or emotional support	3.8	1.9 - 5.7	1,600	3.7	1.8 -	5.7	1,500				
Substance Use											
Any smoking, 3 months before pregnancy	12.3	8.8 - 15.8	5,200	12.4	8.8 -	16.0	4,800				
Any smoking, 3rd trimester†	2.2	1.0 - 3.4	900	2.2	0.9 -	3.4	900				
Any smoking, postpartum	5.5	3.1 - 7.9	2,400	5.7	3.2 -	8.2	2,300				
Any binge drinking, 3 months before pregnancy	18.5	14.3 - 22.8	7,800	20.0	15.4 -	24.6	7,800				
Any alcohol use, 3rd trimester†	12.3	8.3 - 16.4	5,300	12.4	8.2 -	16.6	4,900				
Pregnancy Intention and Family Planning											
Mistimed or unwanted pregnancy	30.4	25.5 - 35.4	13,000	32.2	27.0 -	37.4	12,700				
Unsure of pregnancy intentions	14.2	10.3 - 18.0	6,000	12.7	9.1 -	16.4	5,000				
Postpartum birth control use†	82.6	78.3 - 86.9	35,000	83.1	78.7 -	87.4	32,400	60.6 *	30.7 -	90.5	1,200
Infant Sleep and Breastfeeding											
Placed infant on back to sleep	79.9	75.4 - 84.3	34,000	79.9	75.2 -	84.5	31,200				
Infant always or often shared bed	29.8	25.0 - 34.6	12,700	30.6	25.6 -	35.6	11,900				
Intended to breastfeed, before birth	93.1	90.4 - 95.9	39,600	92.7	89.7 -	95.7	36,200				
Intended to breastfeed exclusively, before birth	68.5	63.4 - 73.6	29,200	67.6	62.2 -	72.9	26,400				
Breastfeeding practices											
Any breastfeeding, 1 month after delivery	90.2	87.1 - 93.2	38,000	89.5	86.2 -	92.8	34,600				
Exclusive breastfeeding, 1 month after delivery	54.2	48.7 - 59.8	22,400	51.5	45.8 -	57.3	19,500				
Any breastfeeding, 3 months after delivery	74.5	69.1 - 79.9	24,300	73.5	67.8 -	79.3	21,700				
Exclusive breastfeeding, 3 months after delivery	36.7	30.6 - 42.7	11,900	34.3	28.2 -	40.4	10,000				



MIHA Data Snapshot, San Diego County by Geographical Area, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		San Diego Count	y			Urban			Rural/Frontier	
	%	95% CI	N		%	95% CI	N	%	95% CI	N
Service Utilization and Health Insurance Coverage										
Had a usual source of pre-pregnancy care	75.4	71.0 - 79.8	32,600		74.1	69.5 - 78.8	29,500			
Initiated prenatal care in 1st trimester	89.7	86.7 - 92.7	38,200		89.5	86.4 - 92.6	35,000			
Mom or infant needed but couldn't afford care postpartum	10.2	6.8 - 13.7	4,300		10.7	7.0 - 14.3	4,100			
Participated in WIC during pregnancy [†]	44.3	40.0 - 48.5	18,600		45.0	40.5 - 49.5	17,500	27.6 *	1.2 - 53.9	500
Received CalFresh (food stamps) during pregnancy	19.3	15.5 - 23.1	8,300		19.2	15.3 - 23.2	7,600			
Pre-pregnancy insurance										
Medi-Cal	16.4	13.0 - 19.9	7,100		17.4	13.7 - 21.1	6,900			
Uninsured	20.4	16.2 - 24.5	8,800		21.2	16.8 - 25.6	8,400			
Private	50.4	45.7 - 55.1	21,700		49.2	44.3 - 54.2	19,500	65.3 *	37.2 - 93.4	1,300
Prenatal insurance										
Medi-Cal	38.5	34.5 - 42.5	16,600		39.4	35.2 - 43.7	15,700			
Uninsured	2.7 *	0.9 - 4.5	1,200		2.7 *	0.8 - 4.6	1,100			
Private	45.8	41.2 - 50.4	19,800		44.8	40.1 - 49.6	17,800	65.3 *	37.2 - 93.4	1,300
Maternal postpartum insurance										
Medi-Cal	27.7	23.7 - 31.7	11,800		28.0	23.8 - 32.2	11,000			
Uninsured	13.5	9.9 - 17.2	5,800		14.2	10.3 - 18.1	5,600			
Private	46.5	41.8 - 51.1	19,800		45.6	40.7 - 50.4	17,900	64.7 *	36.5 - 92.8	1,300
Infant health insurance			ŕ				,			•
Medi-Cal	41.0	36.6 - 45.3	17,400		42.6	38.0 - 47.1	16,600			
Uninsured	4.8	2.5 - 7.1	2,000		4.9	2.4 - 7.3	1,900			
Private	41.9	37.2 - 46.5	17,800		40.4	35.5 - 45.2	15,700	64.7 *	36.5 - 92.8	1,300
Maternal Demographics										
Age										
15-19	4.0	2.2 - 5.8	1,700		4.3	2.3 - 6.2	1,700			
20-34	75.9	71.3 - 80.6	32,900		75.6	70.8 - 80.4	30,100	69.8 *	40.5 - 99.2	1,400
35+	20.1	15.7 - 24.5	8,700		20.1	15.6 - 24.7	8,000			
Did not complete high school (or GED)	11.5	8.1 - 14.9	4,900		11.0	7.6 - 14.4	4,300			
Unmarried	36.3	31.5 - 41.2	15,700		37.7	32.5 - 42.8	14,900			
Race/Ethnicity										
White	38.6	33.2 - 44.0	15,500		38.3	32.7 - 43.8	14,300			
Hispanic	43.3	38.1 - 48.6	17,400		43.8	38.3 - 49.2	16,300			
Asian/Pacific Islander	12.4	8.3 - 16.6	5,000		12.3	8.0 - 16.5	4,600			
Black	5.0	4.1 - 6.0	2,000		5.2	4.2 - 6.2	1,900			
Born outside the U.S.	32.8	27.6 - 37.9	14,200	:	32.6	27.3 - 37.8	12,900			
Speaks non-English language at home	26.8	21.9 - 31.6	11,600		27.8	22.8 - 32.9	11,100			
Lives in a high poverty neighborhood	28.9	24.2 - 33.7	12,100		30.0	25.1 - 35.0	12,000			
Income as a percent of the Federal Poverty Guideline							•			
0-100%	33.0	28.5 - 37.6	13,300		34.0	29.2 - 38.8	12,600			
101-200%	19.6	15.3 - 23.8	7,900		19.6	15.2 - 24.1	7,300			
> 200%	47.4	42.7 - 52.1	19,100		46.3	41.4 - 51.3		71.0 *	44.2 - 97.8	1,300

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of San Diego County was 602. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. Urban and Rural/Frontier designations are based on the population size or densities of Medical Service Study Areas. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

^{*} Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



MIHA Data Snapshot, San Diego County by Income, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	San Diego Coun	ty	0-100% Federal Poverty	Guideline	101-200% Federal Poverty	Guideline	ne > 200% Federal Poverty Guideline		
	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N	
Total	100	43,300	33.0 28.5 - 37.6	13,300	19.6 15.3 - 23.8	7,900	47.4 42.7 - 52.1	19,100	
Health Status before Pregnancy									
In good to excellent health	94.1 91.9 - 96.2	40,700	88.3 82.8 - 93.7	11,800	94.0 * 89.4 - 98.6	7,400			
Chronic conditions									
Diabetes†	3.5 * 1.4 - 5.7	1,500	8.7 * 2.7 - 14.6	1,100					
Hypertension†	3.3 1.4 - 5.1	1,400	2.6 * 0.1 - 5.1	300			2.9 * 0.3 - 5.6	600	
Asthma†	9.2 5.9 - 12.5	3,900	7.8 * 3.2 - 12.4	1,000	11.8 * 3.4 - 20.3	900	9.0 * 3.5 - 14.4	1,700	
Nutrition and Weight									
Daily folic acid use, month before pregnancy	38.6 33.3 - 43.9	16,700	30.0 20.8 - 39.1	4,000	21.9 11.4 - 32.4	1,700	52.8 44.2 - 61.4	10,100	
Overweight before pregnancy	24.0 19.3 - 28.7	9,900	28.1 19.1 - 37.1	3,400	28.2 17.6 - 38.8	2,200	19.0 12.1 - 26.0	3,500	
Obese before pregnancy	16.7 12.7 - 20.6	6,900	23.3 15.4 - 31.3	2,800	21.7 11.9 - 31.6	1,700	11.6 6.0 - 17.3	2,200	
Inadequate weight gain during pregnancy	17.6 12.9 - 22.4	6,400	21.7 11.6 - 31.8	2,300	19.6 10.5 - 28.7	1,200	14.0 7.7 - 20.3	2,400	
Excessive weight gain during pregnancy	39.5 33.7 - 45.3	14,400	36.5 26.3 - 46.6	3,900	43.1 30.0 - 56.1	2,700	40.2 31.3 - 49.1	7,000	
Food insecurity during pregnancy	14.1 10.4 - 17.9	6,100	29.5 20.3 - 38.6	3,800	22.1 11.6 - 32.6	1,700			
Intimate Partner Violence (IPV) and Depressive Symptoms									
Physical or psychological IPV during pregnancy	7.2 4.2 - 10.1	3,100	15.9 8.3 - 23.6	2,100	5.5 * 0.3 - 10.8	400			
Prenatal depressive symptoms	13.7 9.9 - 17.6	5,900	17.6 9.7 - 25.4	2,300	20.2 10.0 - 30.4	1,600	7.9 3.3 - 12.5	1,500	
Postpartum depressive symptoms	13.2 9.6 - 16.8	5,700	17.3 10.0 - 24.7	2,300	14.3 6.6 - 22.0	1,100	9.7 4.7 - 14.8	1,900	
Hardships and Support during Pregnancy									
Experienced two or more hardships during childhood	27. 5 22.7 - 32.3	11,600	37.9 28.5 - 47.3	5,000	36.2 24.1 - 48.4	2,700	17.1 10.9 - 23.4	3,200	
Homeless or did not have a regular place to sleep	3.7 1.7 - 5.8	1,600	8.2 * 2.6 - 13.9	1,100					
Moved due to problems paying rent or mortgage	5.7 3.3 - 8.0	2,400	11.4 5.1 - 17.6	1,500	7.5 * 2.3 - 12.8	600			
Woman or partner lost job	12.2 8.9 - 15.4	5,200	20.1 12.5 - 27.7	2,600	14.4 6.1 - 22.7	1,100	6.6 3.2 - 10.0	1,300	
Woman or partner had pay or hours cut back	9.3 6.5 - 12.0	4,000	13.7 8.1 - 19.2	1,800	14.6 6.1 - 23.0	1,100	5.3 * 1.8 - 8.8	1,000	
Became separated or divorced	6.5 3.6 - 9.5	2,800	15.8 7.8 - 23.9	2,100	2.1 * 0.4 - 3.7	200			
Had no practical or emotional support	3.8 1.9 - 5.7	1,600	6.7 * 1.9 - 11.6	900	5.7 * 0.6 - 10.9	400			
Substance Use									
Any smoking, 3 months before pregnancy	12.3 8.8 - 15.8	5,200	19.3 11.7 - 26.8	2,500	13.5 6.0 - 21.1	1,100	7.5 3.1 - 11.9	1,400	
Any smoking, 3rd trimester†	2.2 1.0 - 3.4	900	4.8 * 1.5 - 8.1	600					
Any smoking, postpartum	5.5 3.1 - 7.9	2,400	9.5 4.2 - 14.8	1,300	8.0 * 0.7 - 15.4	600			
Any binge drinking, 3 months before pregnancy	18.5 14.3 - 22.8	7,800	16.4 9.8 - 23.0	2,100	22.0 11.6 - 32.4	1,600	19.4 12.5 - 26.2	3,700	
Any alcohol use, 3rd trimester†	12.3 8.3 - 16.4	5,300	5.5 * 0.6 - 10.4	700	10.3 * 0.9 - 19.7	800	18.6 11.4 - 25.7	3,500	
Pregnancy Intention and Family Planning									
Mistimed or unwanted pregnancy	30.4 25.5 - 35.4	13,000	44.2 34.9 - 53.6	5,800	31.8 20.2 - 43.4	2,500	18.9 12.1 - 25.8	3,600	
Unsure of pregnancy intentions	14.2 10.3 - 18.0	6,000	20.8 13.0 - 28.6	2,700	18.7 8.9 - 28.5	1,500	7.8 * 2.8 - 12.9	1,500	
Postpartum birth control use†	82.6 78.3 - 86.9	35,000	85.4 78.2 - 92.6	11,000	76.6 65.6 - 87.6	5,900	82.0 75.3 - 88.6	15,500	
Infant Sleep and Breastfeeding									
Placed infant on back to sleep	79.9 75.4 - 84.3	34,000	79.1 71.4 - 86.8	10,200	72.7 61.3 - 84.1	5,600	83.6 77.0 - 90.2	15,900	
Infant always or often shared bed	29.8 25.0 - 34.6	12,700	35.8 26.9 - 44.8	4,600	44.4 32.0 - 56.8	3,400	19.7 13.1 - 26.4	3,700	
Intended to breastfeed, before birth	93.1 90.4 - 95.9	39,600	90.9 * 85.3 - 96.6	11,700			94.4 * 90.4 - 98.3	17,900	
Intended to breastfeed exclusively, before birth	68.5 63.4 - 73.6	29,200	60.8 51.3 - 70.2	7,800	75.3 65.4 - 85.2	5,800	73.2 65.3 - 81.0	13,900	
Breastfeeding practices									
Any breastfeeding, 1 month after delivery	90.2 87.1 - 93.2	38,000	91.3 87.2 - 95.5	11,700	81.2 70.5 - 91.9	6,200	93.5 89.7 - 97.3	17,500	
Exclusive breastfeeding, 1 month after delivery	54.2 48.7 - 59.8	22,400	49.2 39.5 - 59.0	6,100	55.5 42.9 - 68.0	4,100	58.7 50.0 - 67.3	10,900	
Any breastfeeding, 3 months after delivery	74.5 69.1 - 79.9	24,300	61.7 50.7 - 72.6	6,200	64.8 50.7 - 79.0	3,900	87.0 81.1 - 92.9	12,100	
Exclusive breastfeeding, 3 months after delivery	36.7 30.6 - 42.7	11,900	20.9 12.7 - 29.2	2,100	33.2 19.7 - 46.7	2,000	50.6 40.4 - 60.8	7,000	



MIHA Data Snapshot, San Diego County by Income, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

				0-100% Fede	ral Poverty	Guideline	101-200% Fe	ederal Poverty	y Guideline	> 200% Federal Poverty Guideline		
	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N
Service Utilization and Health Insurance Coverage												
Had a usual source of pre-pregnancy care	75.4	71.0 - 79.8	32,600	64.1 55.3	2 - 73.1	8,500	74.4 6	4.5 - 84.4	5,900	86.3	80.6 - 92.0	16,500
Initiated prenatal care in 1st trimester	89.7	86.7 - 92.7	38,200	78.7 71.4	4 - 86.1	10,300	89.5 * 8	2.1 - 96.8	6,800			
Mom or infant needed but couldn't afford care postpartum	10.2	6.8 - 13.7	4,300	20.2 11.5	5 - 28.8	2,600	11.6 *	3.7 - 19.4	900	3.2 *	0.6 - 5.9	600
Participated in WIC during pregnancy†	44.3	40.0 - 48.5	18,600	80.3 72.0	0 - 88.5	10,600	62.8 5	0.8 - 74.9	5,000	6.2	2.9 - 9.5	1,200
Received CalFresh (food stamps) during pregnancy	19.3	15.5 - 23.1	8,300	46.8 37.0	6 - 56.1	6,100	19.0	8.4 - 29.6	1,500			
Pre-pregnancy insurance												
Medi-Cal	16.4	13.0 - 19.9	7,100	42.3 33.3	2 - 51.4	5,600	13.0 *	5.2 - 20.7	1,000			
Uninsured	20.4	16.2 - 24.5	8,800	36.4 27.	1 - 45.6	4,800	30.4 1	8.6 - 42.2	2,400	3.7 *	0.9 - 6.5	700
Private	50.4	45.7 - 55.1	21,700	12.0 5.8	8 - 18.2	1,600	41.2 2	9.2 - 53.1	3,200	83.1	77.1 - 89.1	15,900
Prenatal insurance												
Medi-Cal	38.5	34.5 - 42.5	16,600	80.0 72.3	3 - 87.7	10,600	52.6 4	0.7 - 64.5	4,100	4.2 *	1.3 - 7.2	800
Uninsured	2.7	* 0.9 - 4.5	1,200	5.6 * 0.8	8 - 10.5	700						
Private	45.8	41.2 - 50.4	19,800	9.4 * 3.5	5 - 15.3	1,200	25.8 1	5.3 - 36.3	2,000	81.2	74.6 - 87.7	15,500
Maternal postpartum insurance												
Medi-Cal	27.7	23.7 - 31.7	11,800	61.1 51.8	8 - 70.3	7,800	31.2 1	9.5 - 43.0	2,500	3.2 *	0.7 - 5.7	600
Uninsured	13.5	9.9 - 17.2	5,800	24.3 15.8	8 - 32.8	3,100	21.9 1	1.1 - 32.7	1,700	2.6 *	0.2 - 5.0	500
Private	46.5	41.8 - 51.1	19,800	8.3 * 2.3	7 - 13.9	1,100	28.3 1	7.7 - 38.8	2,200	82.1	76.0 - 88.3	15,700
Infant health insurance												
Medi-Cal	41.0	36.6 - 45.3	17,400	82.5 75.2	2 - 89.8	10,500	59.1 4	7.3 - 70.8	4,600	5.7	2.4 - 9.0	1,100
Uninsured	4.8	2.5 - 7.1	2,000	8.1 * 2.9	9 - 13.3	1,000	7.7 *	0.4 - 15.0	600			
Private	41.9	37.2 - 46.5	17,800				16.1	8.1 - 24.1	1,200	80.0	73.3 - 86.6	15,200
Maternal Demographics												
Did not complete high school (or GED)	11.5	8.1 - 14.9	4,900	22.6 14.1	7 - 30.5	3,000	13.7 *	3.0 - 24.4	1,100			
Unmarried	36.3	31.5 - 41.2	15,700	62.4 53.3	3 - 71.4	8,200	46.9 3	4.5 - 59.2	3,700	12.2	6.6 - 17.8	2,300
Race/Ethnicity												
White	38.6	33.2 - 44.0	15,500	21.0 12.3	7 - 29.2	2,600	27.1 1	5.1 - 39.2	1,900	57.7	49.1 - 66.3	10,400
Hispanic	43.3	38.1 - 48.6	17,400	62.7 53.3	3 - 72.1	7,800	58.3 4	5.8 - 70.8	4,100	22.2	15.5 - 28.8	4,000
Asian/Pacific Islander	12.4	8.3 - 16.6	5,000	7.9 * 1.3	3 - 14.5	1,000	7.0 *	0.7 - 13.2	500	16.7	9.6 - 23.8	3,000
Black	5.0	4.1 - 6.0	2,000	7.9 5.:	1 - 10.7	1,000	6.8	3.8 - 9.7	500	3.0	1.5 - 4.5	500
Born outside the U.S.		27.6 - 37.9	14,200	39.7 30.		5,300		3.4 - 46.3	2,700		17.2 - 32.8	4,800
Speaks non-English language at home		21.9 - 31.6	11,600	38.0 28.3		5,000		4.3 - 34.9	1,900		10.1 - 24.0	3,300
Lives in a high poverty neighborhood		24.2 - 33.7	12,100	41.1 31.9		5,300	_	5.3 - 49.7	2,800	_	10.0 - 23.1	3,100

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of San Diego County was 602. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

^{*} Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



MIHA Data Snapshot, San Diego County by Maternal Age, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	San Diego Cour	<u> </u>			20-34 Years		35 Years or Older		
	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N	
Total	100	43,300	4.0 2.2 - 5.8	1,700	75.9 71.3 - 80.6	32,900	20.1 15.7 - 24.5	8,700	
Health Status before Pregnancy									
In good to excellent health	94.1 91.9 - 96.2	40,700			94.3 92.0 - 96.7	31,000	94.5 * 89.6 - 99.3	8,200	
Chronic conditions									
Diabetes†	3.5 * 1.4 - 5.7	1,500			4.2 * 1.4 - 7.0	1,400			
Hypertension†	3.3 1.4 - 5.1	1,400			3.3 * 1.0 - 5.5	1,100			
Asthma†	9.2 5.9 - 12.5	3,900			8.1 4.6 - 11.5	2,600	12.8 * 3.1 - 22.5	1,100	
Nutrition and Weight									
Daily folic acid use, month before pregnancy	38.6 33.3 - 43.9	16,700			36.3 30.2 - 42.4	11,900	52.7 40.3 - 65.0	4,600	
Overweight before pregnancy	24.0 19.3 - 28.7	9,900	25.9 * 5.2 - 46.7	400	24.0 18.5 - 29.4	7,600	23.7 13.0 - 34.3	1,900	
Obese before pregnancy	16.7 12.7 - 20.6	6,900			17.4 12.8 - 21.9	5,500	16.6 7.1 - 26.2	1,300	
Inadequate weight gain during pregnancy	17.6 12.9 - 22.4	6,400			19.2 13.4 - 24.9	5,400	11.7 * 4.0 - 19.4	800	
Excessive weight gain during pregnancy	39.5 33.7 - 45.3	14,400	46.0 20.4 - 71.6	600	38.1 31.6 - 44.6	10,700	44.1 29.7 - 58.4	3,000	
Food insecurity during pregnancy	14.1 10.4 - 17.9	6,100			15.4 10.7 - 20.1	5,000	9.5 * 3.3 - 15.7	800	
Intimate Partner Violence (IPV) and Depressive Symptoms									
Physical or psychological IPV during pregnancy	7.2 4.2 - 10.1	3,100	22.9 * 3.2 - 42.6	400	7.8 4.1 - 11.4	2,500			
Prenatal depressive symptoms	13.7 9.9 - 17.6	5,900			14.2 9.8 - 18.7	4,700	13.3 * 4.6 - 22.0	1,200	
Postpartum depressive symptoms	13.2 9.6 - 16.8	5,700			11.8 8.0 - 15.6	3,900	19.1 9.1 - 29.2	1,700	
Hardships and Support during Pregnancy									
Experienced two or more hardships during childhood	27.5 22.7 - 32.3	11,600	28.7 * 7.4 - 50.0	500	28.4 22.8 - 34.0	9,100	23.8 13.0 - 34.5	2,100	
Homeless or did not have a regular place to sleep	3.7 1.7 - 5.8	1,600			4.5 * 1.8 - 7.2	1,500			
Moved due to problems paying rent or mortgage	5.7 3.3 - 8.0	2,400			6.9 3.8 - 10.0	2,200	1.9 * 0.1 - 3.8	200	
Woman or partner lost job	12.2 8.9 - 15.4	5,200			12.5 8.6 - 16.4	4,100	10.1 * 3.9 - 16.2	900	
Woman or partner had pay or hours cut back	9.3 6.5 - 12.0	4,000			11.2 7.7 - 14.7	3,700	3.5 * 0.5 - 6.6	300	
Became separated or divorced	6.5 3.6 - 9.5	2,800	20.3 * 1.2 - 39.3	300	5.9 2.5 - 9.3	1,900			
Had no practical or emotional support	3.8 1.9 - 5.7	1,600			4.1 1.8 - 6.5	1,400			
Substance Use									
Any smoking, 3 months before pregnancy	12.3 8.8 - 15.8	5,200	26.2 * 4.7 - 47.8	500	13.2 9.0 - 17.3	4,300	6.0 * 1.1 - 10.9	500	
Any smoking, 3rd trimester†	2.2 1.0 - 3.4	900			2.1 * 0.7 - 3.5	700			
Any smoking, postpartum	5.5 3.1 - 7.9	2,400			6.0 3.0 - 8.9	1,900			
Any binge drinking, 3 months before pregnancy	18.5 14.3 - 22.8	7,800			18.4 13.6 - 23.2	5,900	19.1 8.9 - 29.3	1,700	
Any alcohol use, 3rd trimester†	12.3 8.3 - 16.4	5,300			13.8 8.9 - 18.6	4,500	9.5 * 1.5 - 17.6	800	
Pregnancy Intention and Family Planning									
Mistimed or unwanted pregnancy	30.4 25.5 - 35.4	13,000			31.9 26.0 - 37.7	10,400	13.8 6.4 - 21.2	1,100	
Unsure of pregnancy intentions	14.2 10.3 - 18.0	6,000			16. 5 11.8 - 21.2	5,400			
Postpartum birth control use†	82.6 78.3 - 86.9	35,000			85.7 81.1 - 90.3	27,600	69.0 57.5 - 80.5	6,000	
Infant Sleep and Breastfeeding									
Placed infant on back to sleep	79.9 75.4 - 84.3	34,000	72.9 * 53.6 - 92.2	1,200	79.3 74.1 - 84.4	25,700	83.6 * 73.2 - 94.0	7,000	
Infant always or often shared bed	29.8 25.0 - 34.6	12,700	46.0 22.8 - 69.1	800	30.7 25.0 - 36.3	9,900	23.1 12.8 - 33.3	1,900	
Intended to breastfeed, before birth	93.1 90.4 - 95.9	39,600			93.2 90.2 - 96.2	30,300	92.0 * 84.6 - 99.5	7,700	
Intended to breastfeed exclusively, before birth	68.5 63.4 - 73.6	29,200	78.4 * 60.1 - 96.7	1,300	69.8 64.1 - 75.5	22,700	61.5 48.8 - 74.3	5,200	
Breastfeeding practices									
Any breastfeeding, 1 month after delivery	90.2 87.1 - 93.2	38,000			89.4 85.7 - 93.1	28,700	92.0 * 85.9 - 98.1	7,700	
Exclusive breastfeeding, 1 month after delivery	54.2 48.7 - 59.8	22,400	74.5 * 57.5 - 91.6	1,300	55.6 49.3 - 61.8	17,600	44.5 31.2 - 57.7	3,500	
Any breastfeeding, 3 months after delivery	74.5 69.1 - 79.9	24,300	77.0 * 59.1 - 94.9	1,000	73.0 66.6 - 79.3	18,500	80.5 69.2 - 91.9	4,900	
Exclusive breastfeeding, 3 months after delivery	36.7 30.6 - 42.7	11,900	44.1 * 16.8 - 71.5	600	34.5 27.9 - 41.2	8,700	44.2 28.4 - 60.0	2,600	



MIHA Data Snapshot, San Diego County by Maternal Age, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	San Diego (County	15-19 Years		20-34 Years		35 Years or Older		
	% 95% C	N	% 95% CI	N	% 95% CI	N	% 95% CI	N	
Service Utilization and Health Insurance Coverage									
Had a usual source of pre-pregnancy care	75.4 71.0 - 7	9.8 32,600	79.9 * 66.3 - 93.5	1,400	72.6 67.3 - 78.0	23,800	84.9 76.9 - 92.9	7,400	
Initiated prenatal care in 1st trimester	89.7 86.7 - 9	2.7 38,200	61.1 39.1 - 83.1	1,100	91.4 88.3 - 94.5	29,500	88.9 * 81.3 - 96.5	7,600	
Mom or infant needed but couldn't afford care postpartum	10.2 6.8 - 1	3.7 4,300			10.3 6.1 - 14.4	3,300	10.4 * 3.7 - 17.2	900	
Participated in WIC during pregnancy†	44.3 40.0 - 4	8.5 18,600			46.0 40.8 - 51.3	14,800	28.5 18.6 - 38.4	2,400	
Received CalFresh (food stamps) during pregnancy	19.3 15.5 - 2	3.1 8,300	60.7 39.1 - 82.3	1,000	20.7 16.0 - 25.4	6,800	6.3 * 1.9 - 10.7	500	
Pre-pregnancy insurance									
Medi-Cal	16.4 13.0 - 1	9.9 7,100	64.6 * 43.7 - 85.5	1,100	16.0 12.0 - 20.1	5,200	8.3 * 2.4 - 14.1	700	
Uninsured	20.4 16.2 - 2	4.5 8,800			21.0 16.0 - 26.0	6,900	20.5 11.4 - 29.5	1,800	
Private	50.4 45.7 - 5	5.1 21,700	21.2 * 2.7 - 39.6	400	49.2 43.6 - 54.8	16,100	60.8 49.1 - 72.5	5,300	
Prenatal insurance									
Medi-Cal	38.5 34.5 - 4	2.5 16,600			40.2 35.1 - 45.2	13,200	22.8 14.0 - 31.5	2,000	
Uninsured	2.7 * 0.9 -	4.5 1,200			2.1 * 0.2 - 4.0	700			
Private	45.8 41.2 - 5	0.4 19,800			44.7 39.2 - 50.1	14,600	57.8 45.7 - 69.8	5,000	
Maternal postpartum insurance									
Medi-Cal	27.7 23.7 - 3	1.7 11,800	88.2 * 77.2 - 99.2	1,500	28.5 23.5 - 33.4	9,200	12.8 6.3 - 19.4	1,100	
Uninsured	13.5 9.9 - 1	7.2 5,800			13.2 8.9 - 17.5	4,300	16.0 7.5 - 24.5	1,400	
Private	46.5 41.8 - 5	1.1 19,800			45.0 39.5 - 50.6	14,500	61.0 49.3 - 72.6	5,300	
Infant health insurance									
Medi-Cal	41.0 36.6 - 4	5.3 17,400			42.9 37.5 - 48.3	13,900	23.4 14.2 - 32.6	2,000	
Uninsured	4.8 2.5 -	7.1 2,000			4.6 2.0 - 7.3	1,500			
Private	41.9 37.2 - 4	6.5 17,800			40.0 34.5 - 45.5	12,900	57.7 45.3 - 70.0	4,800	
Maternal Demographics									
Did not complete high school (or GED)	11.5 8.1 - 1	4.9 4,900	52.6 29.8 - 75.5	900	10.4 6.4 - 14.5	3,400	7.2 * 1.8 - 12.7	600	
Unmarried	36.3 31.5 - 4	1.2 15,700			36.2 30.4 - 41.9	11,800	24.9 14.4 - 35.3	2,200	
Race/Ethnicity									
White	38.6 33.2 - 4	4.0 15,500			38.8 32.5 - 45.0	11,800	44.4 31.4 - 57.4	3,600	
Hispanic	43.3 38.1 - 4	8.6 17,400	84.5 * 72.5 - 96.5	1,300	43.9 37.8 - 50.1	13,400	33.2 21.8 - 44.7	2,700	
Asian/Pacific Islander	12.4 8.3 - 1	6.6 5,000			11.3 6.6 - 16.0	3,500	18.6 7.9 - 29.2	1,500	
Black	5.0 4.1 -	6.0 2,000	6.6 * 0.6 - 12.6	100	5.4 4.0 - 6.7	1,600	3.4 1.4 - 5.4	300	
Born outside the U.S.	32.8 27.6 - 3	7.9 14,200			31.1 25.2 - 37.0	10,200	44.4 32.1 - 56.7	3,900	
Speaks non-English language at home	26.8 21.9 - 3	1.6 11,600	25.6 * 4.6 - 46.5	400	23.5 18.2 - 28.9	7,700	39.2 27.1 - 51.3	3,400	
Lives in a high poverty neighborhood	28.9 24.2 - 3	3.7 12,100	52.1 28.7 - 75.6	900	29.0 23.4 - 34.6	9,100	23.9 13.8 - 34.1	2,100	

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of San Diego County was 602. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

^{*} Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



MIHA Data Snapshot, San Diego County by Neighborhood Poverty, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	% 95% CI N %			.9% of Reside elow Povert		5-9.9% of Residual Below Pover		10-19.9% of Resi below Pover		≥ 20% of Residents below Poverty		
	%	95% CI	N	%	95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N
Total	100		43,300	10.0	6.6 - 13.4	4,200	22.8 17.9 - 27.6	9,500	38.3 33.0 - 43.6	16,000	28.9 24.2 - 33.7	12,100
Health Status before Pregnancy												
In good to excellent health	94.1	91.9 - 96.2	40,700						94.5 * 90.7 - 98.3	15,100	89.8 84.9 - 94.7	10,800
Chronic conditions												
Diabetes†	3.5 *	1.4 - 5.7	1,500						2.8 * 0.4 - 5.3	400	4.5 * 0.4 - 8.7	500
Hypertension†	3.3	1.4 - 5.1	1,400						4.5 * 0.7 - 8.2	700	5.0 * 0.6 - 9.4	600
Asthma†	9.2	5.9 - 12.5	3,900						12.4 6.0 - 18.8	2,000	7.8 * 2.6 - 12.9	900
Nutrition and Weight												
Daily folic acid use, month before pregnancy	38.6	33.3 - 43.9	16,700	64.5	48.0 - 80.9	2,700	41.5 29.2 - 53.8	3,900	35.7 27.1 - 44.3	5,700	29.0 19.7 - 38.3	3,500
Overweight before pregnancy	24.0	19.3 - 28.7	9,900	23.8	* 7.7 - 39.9	900	19.5 10.0 - 28.9	1,900	25.0 17.3 - 32.7	3,800	26.5 17.3 - 35.8	3,000
Obese before pregnancy	16.7	12.7 - 20.6	6,900				6.9 * 2.5 - 11.2	700	21.5 13.8 - 29.2	3,300	23.1 15.0 - 31.1	2,600
Inadequate weight gain during pregnancy	17.6	12.9 - 22.4	6,400	17.5	* 3.4 - 31.6	600	21.1 9.7 - 32.5	1,900	13.3 7.3 - 19.3	1,900	21.9 11.5 - 32.4	2,000
Excessive weight gain during pregnancy	39.5	33.7 - 45.3	14,400	36.3	17.5 - 55.1	1,200	41.6 28.7 - 54.6	3,700	40.7 31.7 - 49.8	5,700	38.3 27.4 - 49.2	3,500
Food insecurity during pregnancy	14.1	10.4 - 17.9	6,100				12.0 * 2.4 - 21.6	1,100	12.5 7.3 - 17.7	2,000	20.1 11.7 - 28.5	2,400
Intimate Partner Violence (IPV) and Depressive Symptoms												
Physical or psychological IPV during pregnancy	7.2	4.2 - 10.1	3,100				7.2 * 0.9 - 13.4	700	6.0 * 1.5 - 10.6	1,000	10.7 * 4.0 - 17.4	1,300
Prenatal depressive symptoms		9.9 - 17.6	5,900	10.1	* 0.3 - 19.9	400	15.0 6.6 - 23.5	1,400	14.5 8.4 - 20.7	2,300	13.7 5.9 - 21.6	1,700
Postpartum depressive symptoms	_	9.6 - 16.8	5,700				10.6 * 3.9 - 17.4	1,000	12.6 7.2 - 18.0	2,000	18.8 10.3 - 27.3	2,300
Hardships and Support during Pregnancy												
Experienced two or more hardships during childhood	27 5	22.7 - 32.3	11,600	9.8	* 1.1 - 18.4	400	25.9 14.8 - 37.0	2,400	27.1 19.6 - 34.6	4,200	36.5 26.5 - 46.5	4,300
Homeless or did not have a regular place to sleep	_	1.7 - 5.8	1,600		1.1 10.4	400	25.5 14.0 57.0	2,400	4.3 * 0.4 - 8.3	700	2.3 * 0.7 - 3.9	300
Moved due to problems paying rent or mortgage		3.3 - 8.0	2,400				7.2 * 1.6 - 12.9	700	3.8 * 1.1 - 6.5	600	8.2 * 2.2 - 14.1	1,000
Woman or partner lost job		8.9 - 15.4	5,200				7.5 * 2.3 - 12.8	700	14.2 8.8 - 19.5	2,200	14.0 7.0 - 21.1	1,700
Woman or partner had pay or hours cut back	9.3	6.5 - 12.0	4,000				7.9 * 2.5 - 13.2	800	10.0 5.5 - 14.6	1,600	9.2 4.0 - 14.5	1,100
Became separated or divorced		3.6 - 9.5	2,800						7.1 * 2.1 - 12.1	1,100	8.1 * 2.0 - 14.1	1,000
Had no practical or emotional support		1.9 - 5.7	1,600						3.6 * 1.0 - 6.2	600	2.9 * 0.3 - 5.4	300
Substance Use	0.0		,						5.0			
Any smoking, 3 months before pregnancy	12.3	8.8 - 15.8	5,200				11.1 * 3.2 - 19.1	1,000	12.8 7.2 - 18.5	2,000	14.0 7.5 - 20.6	1,700
Any smoking, 3rd trimester†		1.0 - 3.4	900					1,000	12.0 7.2 10.5	2,000	2.8 * 0.5 - 5.2	300
Any smoking, postpartum		3.1 - 7.9	2,400						4.8 * 1.8 - 7.8	800	4.5 * 1.1 - 7.8	500
Any binge drinking, 3 months before pregnancy		14.3 - 22.8	7,800		* 2.9 - 24.2	600	18.3 8.8 - 27.9	1,700	23.3 15.7 - 30.9	3,600	15.9 8.7 - 23.1	1,900
Any alcohol use, 3rd trimester†		8.3 - 16.4	5,300		* 1.2 - 32.4	700	12.2 * 3.4 - 21.0	,	9.8 4.2 - 15.3	1,500	15.6 7.1 - 24.2	1,900
Pregnancy Intention and Family Planning	12.5	0.5 10.4	3,300	10.0	1.2 32.4	700	12.2 5.4 21.0	1,200	J.0 4.2 15.5	1,500	13.0 7.1 24.2	1,500
Mistimed or unwanted pregnancy	20.4	25.5 - 35.4	13,000	22.0	* 7.6 - 40.1	1,000	26.8 15.6 - 38.0	2,500	31.3 23.6 - 39.0	4,900	36.4 26.7 - 46.0	4,400
Unsure of pregnancy intentions		25.5 - 35.4 10.3 - 18.0	6,000	23.9	7.6 - 40.1	1,000	26.8 15.6 - 38.0 15.1 * 6.0 - 24.3	2,500 1,400	16.3 9.5 - 23.1	2,500	13.1 7.0 - 19.3	1,600
Postpartum birth control use†		78.3 - 86.9	35,000		* 63.0 - 94.6	3,300	81.2 71.0 - 91.5	7,600	82.4 76.1 - 88.8	12,900	83.1 74.8 - 91.4	9,700
·	02.0	70.3 - 60.3	33,000	70.0	03.0 - 94.0	3,300	01.2 /1.0 - 91.3	7,000	02.4 70.1 - 88.8	12,500	03.1 74.0 - 31.4	3,700
Infant Sleep and Breastfeeding	70.0			745			72.2		00.4 == 0.00	40.000	00.0 0 -0 -	
Placed infant on back to sleep		75.4 - 84.3	34,000	_	58.4 - 90.7	3,100	72.2 60.4 - 84.0	6,900	83.4 77.2 - 89.6	13,000	82.9 75.0 - 90.7	9,700
Infant always or often shared bed Intended to breastfeed, before birth		25.0 - 34.6	12,700		11.9 - 47.3	1,200	27.1 16.3 - 37.9	2,600	33.6 25.6 - 41.5	5,200	27.9 19.5 - 36.4	3,300
•		90.4 - 95.9	39,600		* 64.4 05.2	2 200	93.7 * 87.7 - 99.7	8,900	93.4 * 89.2 - 97.6	14,600	90.0 * 83.6 - 96.4	10,600
Intended to breastfeed exclusively, before birth Breastfeeding practices		63.4 - 73.6	29,200		* 64.1 - 95.3	3,300	70.0 58.2 - 81.7	6,700	67.8 59.7 - 75.9	10,600	65.3 55.6 - 74.9	7,700
Any breastfeeding, 1 month after delivery		87.1 - 93.2	38,000				92.3 * 87.0 - 97.5	8,700	90.8 86.4 - 95.2	14,000	84.1 76.1 - 92.0	9,900
Exclusive breastfeeding, 1 month after delivery	_	48.7 - 59.8	22,400	57.7	39.4 - 75.9	2,300	57.2 44.9 - 69.5	5,400	51.8 42.9 - 60.7	8,000	50.3 39.8 - 60.7	5,600
Any breastfeeding, 3 months after delivery		69.1 - 79.9	24,300				75.7 63.6 - 87.8	•	74.3 65.7 - 82.9	8,900	68.1 57.1 - 79.1	6,200
Exclusive breastfeeding, 3 months after delivery	36.7	30.6 - 42.7	11,900	50.1	27.6 - 72.7	1,400	35.4 22.3 - 48.6	2,700	42.4 32.3 - 52.5	5,100	28.4 18.1 - 38.8	2,500



MIHA Data Snapshot, San Diego County by Neighborhood Poverty, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	San Diego Cou	inty	0-4.9% of Residents below Poverty		5-9.9% of Reside below Povert	10-19.9% of Resi below Pover		≥ 20% of Residents below Poverty		
	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N
Service Utilization and Health Insurance Coverage										
Had a usual source of pre-pregnancy care	75.4 71.0 - 79.8	32,600	84.2 * 71.1 - 97.4	3,500	70.0 58.1 - 81.8	6,700	77.6 70.9 - 84.2	12,400	71.7 63.6 - 79.8	8,700
Initiated prenatal care in 1st trimester	89.7 86.7 - 92.7	38,200					86.3 80.7 - 91.9	13,600	90.1 85.3 - 95.0	10,600
Mom or infant needed but couldn't afford care postpartun	10.2 6.8 - 13.7	4,300			11.7 * 2.5 - 20.9	1,100	10.2 5.0 - 15.4	1,600	10.0 * 3.6 - 16.3	1,200
Participated in WIC during pregnancy†	44.3 40.0 - 48.5	18,600	3.7 * 0.4 - 7.0	200	31.8 20.8 - 42.8	2,900	41.6 33.8 - 49.4	6,500	71.7 62.5 - 80.8	8,500
Received CalFresh (food stamps) during pregnancy	19.3 15.5 - 23.1	8,300			9.0 * 2.2 - 15.9	800	20.9 14.5 - 27.3	3,300	29.8 20.7 - 38.8	3,600
Pre-pregnancy insurance										
Medi-Cal	16.4 13.0 - 19.9	7,100	2.9 * 0.1 - 5.8	100	9.4 * 3.1 - 15.8	900	19.2 13.2 - 25.2	3,100	24.7 16.2 - 33.2	3,000
Uninsured	20.4 16.2 - 24.5	8,800			20.8 9.6 - 32.0	2,000	12.5 7.6 - 17.3	2,000	33.8 24.7 - 42.9	4,000
Private	50.4 45.7 - 55.1	21,700	78.4 * 65.1 - 91.6	3,300	51.0 38.9 - 63.1	4,900	55.2 47.2 - 63.3	8,800	32.2 23.3 - 41.2	3,900
Prenatal insurance										
Medi-Cal	38.5 34.5 - 42.5	16,600	4.3 * 0.8 - 7.7	200	28.6 17.3 - 40.0	2,700	36.2 28.9 - 43.6	5,800	60.9 51.7 - 70.1	7,300
Uninsured	2.7 * 0.9 - 4.5	1,200								
Private	45.8 41.2 - 50.4	19,800	77.8 * 64.5 - 91.1	3,300	48.8 36.9 - 60.7	4,600	48.6 40.3 - 56.8	7,800	28.6 19.9 - 37.3	3,400
Maternal postpartum insurance										
Medi-Cal	27.7 23.7 - 31.7	11,800	3.7 * 0.4 - 7.0	200	15.9 7.8 - 23.9	1,500	26.6 19.9 - 33.3	4,300	46.4 36.8 - 56.1	5,500
Uninsured	13.5 9.9 - 17.2	5,800	15.2 * 1.3 - 29.1	600	13.0 * 3.2 - 22.9	1,200	11.8 6.6 - 17.1	1,900	15.7 9.1 - 22.2	1,900
Private	46.5 41.8 - 51.1	19,800	72.7 56.9 - 88.4	3,000	51.6 39.6 - 63.6	4,700	50.1 41.8 - 58.4	8,000	28.4 19.7 - 37.1	3,400
Infant health insurance										
Medi-Cal	41.0 36.6 - 45.3	17,400	8.5 * 1.1 - 15.9	400	32.6 20.8 - 44.4	3,100	38.9 31.2 - 46.7	6,100	63.9 54.5 - 73.2	7,400
Uninsured	4.8 2.5 - 7.1	2,000					6.3 * 1.9 - 10.7	1,000	3.6 * 1.0 - 6.2	400
Private	41.9 37.2 - 46.5	17,800	72.8 57.1 - 88.5	3,000	47.7 35.7 - 59.7	4,500	42.2 33.7 - 50.7	6,600	24.4 15.8 - 33.0	2,800
Maternal Demographics										
Age										
15-19	4.0 2.2 - 5.8	1,700					5.0 * 1.4 - 8.5	800	7.3 * 3.0 - 11.7	900
20-34	75.9 71.3 - 80.6	32,900	67.7 50.4 - 85.0	2,800	86.9 79.4 - 94.4	8,300	70.2 61.8 - 78.6	11,200	75.6 67.4 - 83.7	9,100
35+	20.1 15.7 - 24.5	8,700	32.3 15.0 - 49.6	1,400	12.9 5.4 - 20.4	1,200	24.8 16.7 - 32.9	4,000	17.1 9.7 - 24.4	2,100
Did not complete high school (or GED)	11.5 8.1 - 14.9	4,900					11.7 6.0 - 17.3	1,900	18.1 10.2 - 26.0	2,200
Unmarried	36.3 31.5 - 41.2	15,700			30.7 18.9 - 42.5	2,900	36.6 28.4 - 44.7	5,800	51.4 41.5 - 61.2	6,200
Race/Ethnicity										
White	38.6 33.2 - 44.0	15,500	48.0 29.3 - 66.7	1,900	49.0 36.3 - 61.8	4,500	37.5 28.6 - 46.3	5,500	32.3 21.9 - 42.7	3,600
Hispanic	43.3 38.1 - 48.6	17,400	19.6 * 7.4 - 31.8	800	33.1 21.2 - 45.0	3,000	44.0 35.5 - 52.6	6,500	57.2 46.9 - 67.6	6,300
Asian/Pacific Islander	12.4 8.3 - 16.6	5,000	32.1 * 13.0 - 51.1	1,200	14.2 * 3.9 - 24.4	1,300	11.8 5.9 - 17.6	1,700		
Black	5.0 4.1 - 6.0	2,000			3.3 1.4 - 5.2	300	5.7 3.6 - 7.9	800	7.3 4.3 - 10.2	800
Born outside the U.S.	32.8 27.6 - 37.9	14,200	41.0 22.6 - 59.4	1,700	28.5 16.8 - 40.2	2,700	31.8 23.8 - 39.7	5,000	32.0 22.9 - 41.1	3,900
Speaks non-English language at home	26.8 21.9 - 31.6	11,600	33.7 16.2 - 51.2	1,400	18.9 8.0 - 29.9	1,800	23.2 16.2 - 30.2	3,700	35.4 26.1 - 44.7	4,300
Income as a percent of the Federal Poverty Guideline		•	•	•		•	-	•	•	•
0-100%	33.0 28.5 - 37.6	13,300			24.0 12.2 - 35.7	2,000	35.6 27.5 - 43.7	5,300	47.8 37.8 - 57.8	5,300
101-200%	19.6 15.3 - 23.8	7,900	15.4 * 2.0 - 28.9	600	16.4 7.4 - 25.5	1,400	17.3 11.0 - 23.6	2,600	24.9 16.0 - 33.9	2,800
> 200%	47.4 42.7 - 52.1	19,100	77.8 * 63.0 - 92.7	3,300	59.6 47.1 - 72.1	5,100	47.1 38.4 - 55.9	7,100	27.3 17.9 - 36.7	3,100

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of San Diego County was 602. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. Neighborhood poverty is the percentage of residents with incomes below poverty in a woman's census tract. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

^{*} Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



MIHA Data Snapshot, San Diego County by Prenatal Health Insurance, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		San Diego Count	у		Med	i-Cal			Private	
	%	95% CI	N	%	95%	CI	N	%	95% CI	N
Total	100		43,300	38.5	34.5 -	42.5	16,600	45.8	41.2 - 50.4	19,800
Health Status before Pregnancy										
In good to excellent health	94.1	91.9 - 96.2	40,700	88.2	83.3 -	93.2	14,700	97.9 *	96.0 - 99.7	19,400
Chronic conditions										
Diabetes†	3.5 *	1.4 - 5.7	1,500	6.7 *	1.9 -	11.5	1,100			
Hypertension†	3.3	1.4 - 5.1	1,400	3.6 *	0.2 -	7.1	600	3.5 *	0.7 - 6.3	700
Asthma†	9.2	5.9 - 12.5	3,900	8.4	3.6 -	13.2	1,400	12.4	6.5 - 18.4	2,400
Nutrition and Weight										
Daily folic acid use, month before pregnancy	38.6	33.3 - 43.9	16,700	20.4	13.1 -	27.7	3,400	56.5	48.2 - 64.8	11,200
Overweight before pregnancy	24.0	19.3 - 28.7	9,900	27.1	19.3 -	35.0	4,200	20.6	13.7 - 27.5	4,000
Obese before pregnancy	16.7	12.7 - 20.6	6,900	20.4	14.1 -	26.7	3,100	15.5	9.2 - 21.9	3,000
Inadequate weight gain during pregnancy	17.6	12.9 - 22.4	6,400	20.3	11.5 -	29.1	2,500	15.7	9.4 - 22.0	2,800
Excessive weight gain during pregnancy	39.5	33.7 - 45.3	14,400	41.8	32.1 -	51.6	5,200	39.8	31.2 - 48.5	7,000
Food insecurity during pregnancy	14.1	10.4 - 17.9	6,100	27.3	19.3 -	35.2	4,400			
Intimate Partner Violence (IPV) and Depressive Symptoms										
Physical or psychological IPV during pregnancy	7.2	4.2 - 10.1	3,100	12.3	6.1 -	18.6	2,000			
Prenatal depressive symptoms	13.7	9.9 - 17.6	5,900	16.3	9.7 -	23.0	2,700	8.3	3.7 - 12.9	1,600
Postpartum depressive symptoms	13.2	9.6 - 16.8	5,700	17.2	10.8 -	23.5	2,900	10.7	5.5 - 15.9	2,100
Hardships and Support during Pregnancy										
Experienced two or more hardships during childhood	27.5	22.7 - 32.3	11,600	38.4	30.0 -	46.8	6,300	15.1	9.6 - 20.7	2,900
Homeless or did not have a regular place to sleep	3.7	1.7 - 5.8	1,600	7.3 *	2.7 -	12.0	1,200			
Moved due to problems paying rent or mortgage	5.7	3.3 - 8.0	2,400	9.7	4.6 -	14.9	1,600	2.9 *	0.6 - 5.1	600
Woman or partner lost job	12.2	8.9 - 15.4	5,200	22.1	15.1 -	29.0	3,600	6.0	2.8 - 9.3	1,200
Woman or partner had pay or hours cut back	9.3	6.5 - 12.0	4,000	16.3	10.5 -	22.2	2,700	4.3 *	1.4 - 7.1	800
Became separated or divorced	6.5	3.6 - 9.5	2,800	11.3	5.2 -	17.3	1,800			
Had no practical or emotional support	3.8	1.9 - 5.7	1,600	5.3 *	1.9 -	8.6	900			
Substance Use										
Any smoking, 3 months before pregnancy	12.3	8.8 - 15.8	5,200	17.5	11.4 -	23.6	2,900	5.1 *	1.3 - 8.9	1,000
Any smoking, 3rd trimester†	2.2	1.0 - 3.4	900	3.9 *	1.3 -	6.5	600			,
Any smoking, postpartum	5.5	3.1 - 7.9	2,400	8.4	4.1 -	12.8	1,400			
Any binge drinking, 3 months before pregnancy	18.5	14.3 - 22.8	7,800	17.1	11.0 -	23.1	2,700	16.6	10.4 - 22.8	3,300
Any alcohol use, 3rd trimester†	12.3	8.3 - 16.4	5,300	5.8 *	1.1 -	10.5	1,000	16.9	10.1 - 23.6	3,300
Pregnancy Intention and Family Planning										
Mistimed or unwanted pregnancy	30.4	25.5 - 35.4	13,000	42.3	33.9 -	50.7	6,900	20.0	13.3 - 26.8	3,900
Unsure of pregnancy intentions	14.2	10.3 - 18.0	6,000	21.3	14.2 -	28.5	3,500	10.0	4.6 - 15.3	1,900
Postpartum birth control use†	82.6	78.3 - 86.9	35,000	84.2	77.3 -	91.1	13,600	85.2	79.4 - 91.0	16,600
Infant Sleep and Breastfeeding										
Placed infant on back to sleep	79.9	75.4 - 84.3	34,000	80.5	73.4 -	87.6	13,000	83.5	77.1 - 89.8	16,300
Infant always or often shared bed	29.8	25.0 - 34.6	12,700	38.4	30.1 -	46.7	6,200	21.9	15.1 - 28.6	4,300
Intended to breastfeed, before birth	93.1	90.4 - 95.9	39,600	93.6 *	89.6 -	97.5	15,100	96.1 *	93.4 - 98.8	18,800
Intended to breastfeed exclusively, before birth	68.5	63.4 - 73.6	29,200	64.2	56.0 -	72.5	10,300	73.0	65.4 - 80.6	14,300
Breastfeeding practices	00.5			04.2			,	75.0	, 30.0	,555
Any breastfeeding, 1 month after delivery	90.2	87.1 - 93.2	38,000	85.6	79.6 -	91.7	13,600	95.5 *	92.5 - 98.5	18,600
Exclusive breastfeeding, 1 month after delivery	54.2	48.7 - 59.8	22,400	48.3	39.5 -	57.1	7,400	60.1	51.8 - 68.4	11,700
Any breastfeeding, 3 months after delivery	74.5	69.1 - 79.9	24,300	61.9	52.2 -	71.5	7,900	87.8	81.5 - 94.0	12,700
Exclusive breastfeeding, 3 months after delivery	36.7	30.6 - 42.7	11,900	20.1	12.6 -	27.6	2,500	51.1	41.2 - 61.1	7,400
	30.7		,	20.1			_,_ 00	J1.1	01.1	.,



MIHA Data Snapshot, San Diego County by Prenatal Health Insurance, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	San Diego County				Medi-Cal		Private			
	%	95% CI	N		%	95% CI	N	%	95% CI	N
Service Utilization and Health Insurance Coverage										
Had a usual source of pre-pregnancy care	75.4	71.0 - 79.8	32,600		62.0	54.1 - 69.9	10,300	88.0	82.7 - 93.3	17,400
Initiated prenatal care in 1st trimester	89.7	86.7 - 92.7	38,200		83.6	78.0 - 89.2	13,500			
Mom or infant needed but couldn't afford care postpartum	10.2	6.8 - 13.7	4,300		15.6	8.7 - 22.4	2,500	3.6 *	0.8 - 6.5	700
Participated in WIC during pregnancy†	44.3	40.0 - 48.5	18,600		82.3	75.4 - 89.2	13,500	10.3	5.9 - 14.6	2,000
Received CalFresh (food stamps) during pregnancy	19.3	15.5 - 23.1	8,300		47.0	38.6 - 55.4	7,700			
Pre-pregnancy insurance										
Medi-Cal	16.4	13.0 - 19.9	7,100		42.0	33.8 - 50.2	6,900			
Uninsured	20.4	16.2 - 24.5	8,800		43.2	34.8 - 51.5	7,100			
Private	50.4	45.7 - 55.1	21,700		11.8	6.4 - 17.1	1,900	97.2 *	95.0 - 99.5	19,200
Maternal postpartum insurance										
Medi-Cal	27.7	23.7 - 31.7	11,800		69.3	61.7 - 76.9	11,200			
Uninsured	13.5	9.9 - 17.2	5,800		25.9	18.5 - 33.3	4,200			
Private	46.5	41.8 - 51.1	19,800		3.3 *	0.9 - 5.6	500	94.7 *	91.0 - 98.4	18,700
Infant health insurance										
Medi-Cal	41.0	36.6 - 45.3	17,400		95.3	92.5 - 98.0	15,300	6.3 *	2.6 - 10.0	1,200
Uninsured	4.8	2.5 - 7.1	2,000		3.7 *	1.2 - 6.2	600	4.0 *	0.5 - 7.4	800
Private	41.9	37.2 - 46.5	17,800					88.6	83.5 - 93.7	17,400
Maternal Demographics										
Age										
15-19	4.0	2.2 - 5.8	1,700		8.9	4.6 - 13.3	1,500			
20-34	75.9	71.3 - 80.6	32,900		79.2	73.0 - 85.3	13,200	74.1	66.7 - 81.5	14,600
35+	20.1	15.7 - 24.5	8,700		11.9	7.0 - 16.8	2,000	25.4	18.1 - 32.8	5,000
Did not complete high school (or GED)	11.5	8.1 - 14.9	4,900		22.4	15.1 - 29.7	3,700			
Unmarried	36.3	31.5 - 41.2	15,700		63.9	56.1 - 71.7	10,500	16.2	10.3 - 22.0	3,200
Race/Ethnicity										
White	38.6	33.2 - 44.0	15,500		19.2	11.8 - 26.7	2,900	53.6	45.1 - 62.1	9,900
Hispanic	43.3	38.1 - 48.6	17,400		67.0	58.6 - 75.4	10,100	25.1	18.3 - 31.9	4,600
Asian/Pacific Islander	12.4	8.3 - 16.6	5,000		6.9 *	1.3 - 12.4	1,000	18.0	10.9 - 25.1	3,300
Black	5.0	4.1 - 6.0	2,000		6.2	4.1 - 8.2	900	2.8	1.5 - 4.1	500
Born outside the U.S.	32.8	27.6 - 37.9	14,200		38.4	30.2 - 46.6	6,300	29.4	21.6 - 37.2	5,800
Speaks non-English language at home	26.8	21.9 - 31.6	11,600		37.0	28.9 - 45.2	6,200	18.2	11.4 - 24.9	3,600
Lives in a high poverty neighborhood	28.9	24.2 - 33.7	12,100		45.7	37.3 - 54.1	7,300	18.0	11.8 - 24.2	3,400
Income as a percent of the Federal Poverty Guideline										
0-100%	33.0	28.5 - 37.6	13,300		68.1	59.8 - 76.5	10,600	6.6 *	2.4 - 10.9	1,200
101-200%	19.6	15.3 - 23.8	7,900		26.7	18.7 - 34.6	4,100	10.8	5.9 - 15.8	2,000
> 200%	47.4	42.7 - 52.1	19,100		5.2 *	1.5 - 8.9	800	82.5	76.3 - 88.7	15,500

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of San Diego County was 602. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

^{*} Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%)

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



MIHA Data Snapshot, San Diego County by Race/Ethnicity, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	San Diego County			Hispanic		Black		White	Asian/Pacific Islander		
	%	95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N
Total	100		43,300	43.3 38.1 - 48.6	17,400	5.0 4.1 - 6.0	2,000	38.6 33.2 - 44.0	15,500	12.4 8.3 - 16.6	5,000
Health Status before Pregnancy											
In good to excellent health	94.1	91.9 - 96.2	40,700	91.7 87.9 - 95.5	16,000	89.6 * 82.6 - 96.5	1,800	95.6 * 92.3 - 99.0	14,900		
Chronic conditions											
Diabetes†	3.5 *	1.4 - 5.7	1,500	3.9 * 1.0 - 6.9	700						
Hypertension†	3.3	1.4 - 5.1	1,400	4.3 * 0.8 - 7.8	700						
Asthma†	9.2	5.9 - 12.5	3,900	7.7 3.3 - 12.1	1,300	7.5 * 2.2 - 12.7	100	12.2 5.4 - 18.9	1,900		
Nutrition and Weight											
Daily folic acid use, month before pregnancy	38.6	33.3 - 43.9	16,700	26.9 19.6 - 34.2	4,700	30.9 19.7 - 42.2	600	46.5 36.8 - 56.2	7,200	63.2 46.8 - 79.6	3,200
Overweight before pregnancy	24.0	19.3 - 28.7	9,900	28.7 21.0 - 36.4	4,600	26.4 14.9 - 38.0	500	22.8 14.7 - 30.9	3,500		
Obese before pregnancy	16.7	12.7 - 20.6	6,900	18.1 11.9 - 24.3	2,900	29.5 16.8 - 42.2	600	15.3 8.6 - 21.9	2,400		
Inadequate weight gain during pregnancy	17.6	12.9 - 22.4	6,400	26.1 17.3 - 34.9	3,600	11.6 * 4.0 - 19.3	200	11.9 5.6 - 18.2	1,700		
Excessive weight gain during pregnancy	39.5	33.7 - 45.3	14,400	35.8 27.0 - 44.7	4,900	50.7 36.7 - 64.7	900	46.0 36.0 - 55.9	6,500	20.7 * 4.6 - 36.9	900
Food insecurity during pregnancy	14.1	10.4 - 17.9	6,100	18.6 11.7 - 25.4	3,200	18.1 8.5 - 27.7	400	10.6 4.4 - 16.7	1,600		
Intimate Partner Violence (IPV) and Depressive Symptoms											
Physical or psychological IPV during pregnancy	7.2	4.2 - 10.1	3,100	8.1 * 3.1 - 13.1	1,400	11.5 * 3.2 - 19.8	200	5.9 * 1.4 - 10.5	900		
Prenatal depressive symptoms	13.7	9.9 - 17.6	5,900	17.9 11.5 - 24.3	3,100	13.1 * 4.0 - 22.2	300	9.4 * 3.5 - 15.2	1,500		
Postpartum depressive symptoms	13.2	9.6 - 16.8	5,700	16.5 10.4 - 22.5	2,900	17.9 * 6.3 - 29.4	400	9.8 4.2 - 15.4	1,500		
Hardships and Support during Pregnancy											
Experienced two or more hardships during childhood	27.5	22.7 - 32.3	11,600	41.5 33.3 - 49.7	7,100	27.2 16.5 - 38.0	500	19.2 11.6 - 26.8	2,900		
Homeless or did not have a regular place to sleep	3.7	1.7 - 5.8	1,600	3.1 * 0.4 - 5.7	500	8.1 * 0.2 - 16.0	200				
Moved due to problems paying rent or mortgage	5.7	3.3 - 8.0	2,400	7.0 * 2.4 - 11.5	1,200	14.0 * 4.9 - 23.0	300	5.3 * 1.6 - 8.9	800		
Woman or partner lost job	12.2	8.9 - 15.4	5,200	16.8 10.7 - 22.9	2,900	9.5 * 3.2 - 15.8	200	7.3 3.4 - 11.2	1,100	7.8 * 0.7 - 14.8	400
Woman or partner had pay or hours cut back	9.3	6.5 - 12.0	4,000	11.7 6.9 - 16.6	2,000	12.0 * 4.4 - 19.5	200	6.9 * 2.6 - 11.2	1,100		
Became separated or divorced	6.5	3.6 - 9.5	2,800	9.8 4.1 - 15.4	1,700	13.7 6.2 - 21.3	300				
Had no practical or emotional support	3.8	1.9 - 5.7	1,600	7.0 2.9 - 11.2	1,200						
Substance Use											
Any smoking, 3 months before pregnancy	12.3	8.8 - 15.8	5,200	10.7 5.9 - 15.6	1,900	14.3 * 5.4 - 23.2	300	16.0 8.9 - 23.1	2,400		
Any smoking, 3rd trimester†	2.2	1.0 - 3.4	900	1.9 * 0.2 - 3.6	300						
Any smoking, postpartum	5.5	3.1 - 7.9	2,400	2.6 * 0.7 - 4.4	400	11.7 * 3.1 - 20.3	200	8.1 * 2.8 - 13.3	1,200		
Any binge drinking, 3 months before pregnancy	18.5	14.3 - 22.8	7,800	20.6 14.1 - 27.1	3,500	12.1 * 4.5 - 19.7	200	19.4 11.5 - 27.3	3,000	8.5 * 0.7 - 16.2	400
Any alcohol use, 3rd trimester†	12.3	8.3 - 16.4	5,300	6.8 * 2.3 - 11.3	1,200			21.8 13.2 - 30.4	3,400		
Pregnancy Intention and Family Planning											
Mistimed or unwanted pregnancy	30.4	25.5 - 35.4	13,000	37.8 30.0 - 45.7	6,500	40.8 27.9 - 53.7	800	25.2 16.6 - 33.8	3,800	26.3 * 10.0 - 42.5	1,300
Unsure of pregnancy intentions	14.2	10.3 - 18.0	6,000	16.5 10.4 - 22.6	2,800	18.6 9.4 - 27.8	400	7.5 * 2.6 - 12.4	1,100		
Postpartum birth control use†	82.6	78.3 - 86.9	35,000	86.8 80.9 - 92.8	14,800	69.7 56.6 - 82.8	1,300	79.8 71.8 - 87.8	12,200	83.0 * 69.8 - 96.2	4,200
Infant Sleep and Breastfeeding											
Placed infant on back to sleep	79.9	75.4 - 84.3	34,000	86.4 81.3 - 91.5	14,900	74.5 64.2 - 84.9	1,400	79.7 71.9 - 87.6	12,100	69.1 * 50.8 - 87.3	3,500
Infant always or often shared bed	29.8	25.0 - 34.6	12,700	36.5 28.9 - 44.2	6,300	52.5 39.3 - 65.7	1,000	13.5 7.2 - 19.8	2,100	41.7 23.5 - 59.9	2,100
Intended to breastfeed, before birth	93.1	90.4 - 95.9	39,600	90.4 85.1 - 95.8	15,600			94.9 * 91.1 - 98.6	14,400		
Intended to breastfeed exclusively, before birth	68.5	63.4 - 73.6	29,200	60.7 52.8 - 68.6	10,500	72.5 61.1 - 84.0	1,400	79.0 70.9 - 87.1	12,000	54.9 36.4 - 73.4	2,700
Breastfeeding practices											
Any breastfeeding, 1 month after delivery	90.2	87.1 - 93.2	38,000	88.1 83.5 - 92.7	15,000	87.8 81.4 - 94.1	1,700	91.1 * 85.1 - 97.2	13,700	90.8 * 82.0 - 99.6	4,500
Exclusive breastfeeding, 1 month after delivery	54.2	48.7 - 59.8	22,400	56.0 48.1 - 64.0	9,400	42.1 28.5 - 55.7	800	59.4 49.7 - 69.2	9,000	39.8 21.5 - 58.1	1,900
Any breastfeeding, 3 months after delivery	74.5	69.1 - 79.9	24,300	64.2 55.2 - 73.2	8,500	76.9 66.6 - 87.2	1,100	82.5 73.4 - 91.6	9,400	80.2 * 65.4 - 95.0	3,100
Exclusive breastfeeding, 3 months after delivery	36.7	30.6 - 42.7	11,900	28.1 20.1 - 36.2	3,800	22.1 10.3 - 34.0	300	49.2 37.7 - 60.6	5,600	27.1 * 8.9 - 45.2	1,000



MIHA Data Snapshot, San Diego County by Race/Ethnicity, 2013-2015

Maternal and Infant Health Assessment (MIHA) Survey

	San Diego County		Hispanic		Black			White			Asian/Pacific Islander				
•	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N
Service Utilization and Health Insurance Coverage															
Had a usual source of pre-pregnancy care	75.4	71.0 - 79.8	32,600	71.2 6	1.4 - 78.0	12,400	69.3	58.0 - 80.6	1,400	82.2	74.9 - 89.4	12,700	75.3 *	58.0 - 92.5	3,800
Initiated prenatal care in 1st trimester	89.7	86.7 - 92.7	38,200	82.9 7	5.8 - 88.9	14,200	88.8	81.2 - 96.4	1,800	96.3 *	93.3 - 99.3	14,900			
Mom or infant needed but couldn't afford care postpartum	10.2	6.8 - 13.7	4,300	16.4	9.7 - 23.1	2,800	12.0	4.1 - 19.9	200	2.1 *	0.2 - 3.9	300			
Participated in WIC during pregnancy†	44.3	40.0 - 48.5	18,600	67.3 5	9.9 - 74.6	11,300	68.4	59.0 - 77.7	1,400	20.4	13.4 - 27.3	3,100	24.6 *	9.4 - 39.8	1,200
Received CalFresh (food stamps) during pregnancy	19.3	15.5 - 23.1	8,300	25.5 1	3.6 - 32.4	4,400	35.7	23.3 - 48.2	700	11.8	5.5 - 18.0	1,800			
Pre-pregnancy insurance															
Medi-Cal	16.4	13.0 - 19.9	7,100	26.0 19	9.1 - 32.9	4,500	37.0	23.8 - 50.1	700	6.5 *	2.2 - 10.7	1,000			
Uninsured	20.4	16.2 - 24.5	8,800	30.9 2	3.5 - 38.3	5,400	10.7	5.3 - 16.1	200	11.8	5.3 - 18.2	1,800			
Private	50.4	45.7 - 55.1	21,700	32.9 2	5.8 - 40.0	5,700	28.5	17.0 - 39.9	600	66.2	57.5 - 74.9	10,300	69.8	52.7 - 86.8	3,500
Prenatal insurance															
Medi-Cal	38.5	34.5 - 42.5	16,600	57.7 5	0.4 - 64.9	10,100	45.8	33.7 - 57.8	900	18.6	11.6 - 25.6	2,900	20.6 *	5.4 - 35.7	1,000
Uninsured	2.7 *	0.9 - 4.5	1,200	4.8 *	0.8 - 8.7	800									
Private	45.8	41.2 - 50.4	19,800	26.6 20	0.1 - 33.2	4,600	25.7	14.8 - 36.7	500	63.9	54.9 - 72.8	9,900	66.6	49.5 - 83.8	3,300
Maternal postpartum insurance															
Medi-Cal	27.7	23.7 - 31.7	11,800	42.1 3	1.5 - 49.6	7,300	45.5	32.8 - 58.1	900	12.2	6.3 - 18.1	1,900	10.6 *	0.7 - 20.4	500
Uninsured	13.5	9.9 - 17.2	5,800	21.9 1	5.0 - 28.8	3,800	7.1	1.4 - 12.9	100	5.3 *	0.9 - 9.6	800			
Private	46.5	41.8 - 51.1	19,800	26.0 19	9.5 - 32.5	4,500	23.2	12.6 - 33.8	500	67.2	58.7 - 75.7	10,400	71.3	55.4 - 87.1	3,300
Infant health insurance															
Medi-Cal	41.0	36.6 - 45.3	17,400	61.9 5	1.6 - 69.2	10,700	51.9	39.6 - 64.1	1,000	19.9	12.5 - 27.2	3,000	23.5 *	7.9 - 39.1	1,200
Uninsured	4.8	2.5 - 7.1	2,000	7.9	3.4 - 12.3	1,400									
Private	41.9	37.2 - 46.5	17,800	21.4 1	5.6 - 27.3	3,700	18.1	8.1 - 28.0	300	62.8	53.7 - 71.9	9,500	63.2	45.8 - 80.6	3,200
Maternal Demographics															
Age															
15-19	4.0	2.2 - 5.8	1,700	7.6	3.4 - 11.8	1,300	5.1	* 0.9 - 9.4	100						
20-34	75.9	71.3 - 80.6	32,900	76.8 7	0.1 - 83.6	13,400	81.1	72.3 - 89.9	1,600	76.1	67.7 - 84.6	11,800	69.0	52.4 - 85.6	3,500
35+	20.1	15.7 - 24.5	8,700	15.6	9.8 - 21.3	2,700	13.8	6.0 - 21.6	300	23.4	15.0 - 31.8	3,600	30.3	13.7 - 46.9	1,500
Did not complete high school (or GED)	11.5	8.1 - 14.9	4,900	20.4 1	3.6 - 27.2	3,500	13.3	* 3.9 - 22.8	300						
Unmarried	36.3	31.5 - 41.2	15,700	50.9 4	3.0 - 58.8	8,800	54.1	40.9 - 67.3	1,100	21.7	13.9 - 29.6	3,300	20.1 *	5.7 - 34.5	1,000
Born outside the U.S.	32.8	27.6 - 37.9	14,200	42.1 3	1.2 - 50.1	7,400	21.7	10.6 - 32.7	400	11.6	6.0 - 17.2	1,800	68.2	52.9 - 83.5	3,400
Speaks non-English language at home	26.8	21.9 - 31.6	11,600	37.1 2	9.4 - 44.8	6,500	11.4	3.8 - 19.0	200	9.4 *	3.8 - 15.0	1,500	56.9	39.2 - 74.6	2,800
Lives in a high poverty neighborhood	28.9	24.2 - 33.7	12,100	38.0 3	0.1 - 45.9	6,300	40.8	27.6 - 53.9	800	23.0	14.6 - 31.4	3,600			
Income as a percent of the Federal Poverty Guideline															
0-100%	33.0	28.5 - 37.6	13,300	49.0 4	0.9 - 57.1	7,800	49.0	36.1 - 61.9	1,000	17.4	10.4 - 24.4	2,600	21.8 *	5.0 - 38.5	1,000
101-200%	19.6	15.3 - 23.8	7,900	25.8 1	3.4 - 33.1	4,100	23.9	14.1 - 33.7	500	12.8	6.3 - 19.2	1,900	10.9 *	1.0 - 20.8	500
> 200%	47. <u>4</u>	42.7 - 52.1	19,100	25.2 1	3.5 - 32.0	4,000	27.1	15.4 - 38.8	500	69.8	61.3 - 78.4	10,400	67.3	49.5 - 85.1	3,000

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of San Diego County was 602. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

^{*} Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



MIHA Data Snapshot, San Diego County by Total Live Births, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		San Diego Count	у		First Liv	e Birth		Second Live Birth or More			
	%	95% CI	N	%	95%	CI	N	%	95% CI	N	
Total	100		43,300	43.1	37.7 -	48.6	18,700	56.9	51.4 - 62.3	24,600	
Health Status before Pregnancy											
In good to excellent health	94.1	91.9 - 96.2	40,700	93.8	90.4 -	97.1	17,500	94.3	91.5 - 97.1	23,200	
Chronic conditions											
Diabetes†	3.5 *	1.4 - 5.7	1,500					3.2 *	1.2 - 5.1	800	
Hypertension [†]	3.3	1.4 - 5.1	1,400					4.4 *	1.7 - 7.1	1,100	
Asthma†	9.2	5.9 - 12.5	3,900	14.5	8.0 -	21.0	2,700	5.2	2.4 - 8.0	1,200	
Nutrition and Weight											
Daily folic acid use, month before pregnancy	38.6	33.3 - 43.9	16,700	42.0	33.4 -	50.6	7,800	35.9	29.1 - 42.7	8,800	
Overweight before pregnancy	24.0	19.3 - 28.7	9,900	21.4	14.4 -	28.5	3,900	26.0	19.7 - 32.4	6,000	
Obese before pregnancy	16.7	12.7 - 20.6	6,900	13.0	7.2 -	18.7	2,400	19.6	14.1 - 25.0	4,500	
Inadequate weight gain during pregnancy	17.6	12.9 - 22.4	6,400	9.6	4.5 -	14.8	1,600	24.1	16.9 - 31.3	4,800	
Excessive weight gain during pregnancy	39.5	33.7 - 45.3	14,400	47.2	38.2 -	56.2	7,700	33.3	26.0 - 40.5	6,700	
Food insecurity during pregnancy	14.1	10.4 - 17.9	6,100	12.8	6.9 -	18.7	2,300	15.1	10.1 - 20.2	3,700	
Intimate Partner Violence (IPV) and Depressive Symptoms											
Physical or psychological IPV during pregnancy	7.2	4.2 - 10.1	3,100	8.1 *	3.3 -	12.9	1,500	6.4	2.8 - 10.1	1,600	
Prenatal depressive symptoms	13.7	9.9 - 17.6	5,900	12.1	6.3 -	17.9	2,300	15.0	9.9 - 20.1	3,700	
Postpartum depressive symptoms	13.2	9.6 - 16.8	5,700	11.7	6.5 -	16.9	2,200	14.3	9.4 - 19.2	3,500	
Hardships and Support during Pregnancy											
Experienced two or more hardships during childhood	27.5	22.7 - 32.3	11,600	27.1	19.4 -	34.9	4,900	27.7	21.5 - 34.0	6,700	
Homeless or did not have a regular place to sleep	3.7	1.7 - 5.8	1,600	2.9 *	0.1 -	5.7	500	4.4 *	1.4 - 7.3	1,100	
Moved due to problems paying rent or mortgage	5.7	3.3 - 8.0	2,400	4.7 *	1.6 -	7.8	900	6.4	3.0 - 9.8	1,600	
Woman or partner lost job	12.2	8.9 - 15.4	5,200	12.4	7.7 -	17.2	2,300	12.0	7.5 - 16.5	2,900	
Woman or partner had pay or hours cut back	9.3	6.5 - 12.0	4,000	10.6	6.0 -	15.3	2,000	8.2	4.8 - 11.6	2,000	
Became separated or divorced	6.5	3.6 - 9.5	2,800	7.8 *	2.8 -	12.8	1,400	5.6 *	2.0 - 9.2	1,400	
Had no practical or emotional support	3.8	1.9 - 5.7	1,600	3.2 *	0.1 -	6.4	600	4.2	1.9 - 6.5	1,000	
Substance Use											
Any smoking, 3 months before pregnancy	12.3	8.8 - 15.8	5,200	15.9	9.8 -	22.0	2,900	9.7	5.7 - 13.6	2,400	
Any smoking, 3rd trimester†	2.2	1.0 - 3.4	900	1.9 *	0.4 -	3.3	300	2.4 *	0.6 - 4.2	600	
Any smoking, postpartum	5.5	3.1 - 7.9	2,400	6.4 *	2.4 -	10.4	1,200	4.8 *	1.9 - 7.7	1,200	
Any binge drinking, 3 months before pregnancy	18.5	14.3 - 22.8	7,800	26.4	18.8 -	34.1	4,800	12.5	8.1 - 17.0	3,000	
Any alcohol use, 3rd trimester†	12.3	8.3 - 16.4	5,300	15.2	8.0 -	22.5	2,800	10.1	5.8 - 14.5	2,400	
Pregnancy Intention and Family Planning											
Mistimed or unwanted pregnancy	30.4	25.5 - 35.4	13,000	32.9	24.9 -	41.0	6,200	28.5	22.2 - 34.7	6,800	
Unsure of pregnancy intentions	14.2	10.3 - 18.0	6,000	13.8	8.2 -	19.5	2,600	14.4	9.1 - 19.6	3,400	
Postpartum birth control use†	82.6	78.3 - 86.9	35,000	82.9	76.1 -	89.8	15,200	82.4	76.9 - 87.9	19,800	
Infant Sleep and Breastfeeding											
Placed infant on back to sleep	79.9	75.4 - 84.3	34,000	80.3	73.3 -	87.4	14,900	79.5	73.7 - 85.3	19,100	
Infant always or often shared bed	29.8	25.0 - 34.6	12,700	29.5	22.0 -	37.0	5,500	30.0	23.6 - 36.3	7,200	
Intended to breastfeed, before birth	93.1	90.4 - 95.9	39,600	93.7 *	89.2 -	98.1	17,300	92.7	89.3 - 96.2	22,300	
Intended to breastfeed exclusively, before birth	68.5	63.4 - 73.6	29,200	78.4	70.9 -	85.8	14,500	60.9	54.0 - 67.8	14,600	
Breastfeeding practices	00.3		,_00	, 01		0	,_ 50	00.5	3 373	_ ,,000	
Any breastfeeding, 1 month after delivery	90.2	87.1 - 93.2	38,000	89.8	84.7 -	94.9	16,500	90.4	86.7 - 94.1	21,500	
Exclusive breastfeeding, 1 month after delivery	54.2	48.7 - 59.8	22,400	48.7	40.2 -	57.3	9,000	58.7	51.6 - 65.7	13,400	
Any breastfeeding, 3 months after delivery	74.5	69.1 - 79.9	24,300	76.2	68.1 -	84.3	10,800	73.3	65.9 - 80.6	13,600	
Exclusive breastfeeding, 3 months after delivery	36.7	30.6 - 42.7	11,900	30.7	21.6 -	39.8	4,300	41.2	33.0 - 49.4	7,500	
	30.7	23.0 .2.7	11,500	50.7		33.0	.,500	71.2	22.0 .3.4	.,550	



MIHA Data Snapshot, San Diego County by Total Live Births, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		San Diego Count		First Live Birth		Second Live Birth or More			
	%	95% CI	N	%	95% CI	N	%	95% CI	N
Service Utilization and Health Insurance Coverage									
Had a usual source of pre-pregnancy care	75.4	71.0 - 79.8	32,600	69.1	61.4 - 76.7	12,800	80.2	75.0 - 85.4	19,700
Initiated prenatal care in 1st trimester	89.7	86.7 - 92.7	38,200	93.4	89.6 - 97.1	17,400	86.8	82.4 - 91.2	20,800
Mom or infant needed but couldn't afford care postpartum	10.2	6.8 - 13.7	4,300	7.9 *	3.0 - 12.8	1,500	12.0	7.2 - 16.8	2,800
Participated in WIC during pregnancy†	44.3	40.0 - 48.5	18,600	35.4	28.3 - 42.5	6,400	51.0	44.7 - 57.3	12,200
Received CalFresh (food stamps) during pregnancy	19.3	15.5 - 23.1	8,300	15.7	9.7 - 21.6	2,900	22.0	16.7 - 27.3	5,400
Pre-pregnancy insurance									
Medi-Cal	16.4	13.0 - 19.9	7,100	10.1	5.9 - 14.2	1,900	21.2	15.9 - 26.6	5,200
Uninsured	20.4	16.2 - 24.5	8,800	21.1	14.3 - 27.8	3,900	19.9	14.4 - 25.3	4,900
Private	50.4	45.7 - 55.1	21,700	56.7	48.7 - 64.6	10,600	45.7	39.1 - 52.2	11,200
Prenatal insurance									
Medi-Cal	38.5	34.5 - 42.5	16,600	34.8	27.8 - 41.8	6,500	41.3	35.1 - 47.5	10,100
Uninsured	2.7 *	0.9 - 4.5	1,200				3.0 *	0.6 - 5.5	700
Private	45.8	41.2 - 50.4	19,800	48.8	40.7 - 56.8	9,100	43.5	37.1 - 49.9	10,700
Maternal postpartum insurance									
Medi-Cal	27.7	23.7 - 31.7	11,800	26.8	20.2 - 33.4	4,900	28.4	22.5 - 34.3	6,900
Uninsured	13.5	9.9 - 17.2	5,800	10.4	5.3 - 15.4	1,900	15.9	10.6 - 21.1	3,900
Private	46.5	41.8 - 51.1	19,800	49.6	41.6 - 57.7	9,100	44.1	37.5 - 50.6	10,700
Infant health insurance									
Medi-Cal	41.0	36.6 - 45.3	17,400	36.3	29.0 - 43.7	6,700	44.6	38.1 - 51.1	10,700
Uninsured	4.8	2.5 - 7.1	2,000	3.4 *	0.7 - 6.0	600	5.9 *	2.4 - 9.5	1,400
Private	41.9	37.2 - 46.5	17,800	46.3	38.1 - 54.4	8,600	38.5	32.0 - 44.9	9,200
Maternal Demographics									
Age									
15-19	4.0	2.2 - 5.8	1,700	7.3	3.7 - 11.0	1,400			
20-34	75.9	71.3 - 80.6	32,900	79.8	72.7 - 86.9	14,900	72.9	66.8 - 79.1	18,000
35+	20.1	15.7 - 24.5	8,700	12.9	6.5 - 19.3	2,400	25.6	19.5 - 31.6	6,300
Did not complete high school (or GED)	11.5	8.1 - 14.9	4,900	11.1	5.7 - 16.6	2,000	11.8	7.3 - 16.3	2,900
Unmarried	36.3	31.5 - 41.2	15,700	43.1	34.9 - 51.2	8,000	31.2	25.0 - 37.5	7,700
Race/Ethnicity									
White	38.6	33.2 - 44.0	15,500	46.1	37.2 - 55.0	8,000	32.8	26.1 - 39.6	7,500
Hispanic	43.3	38.1 - 48.6	17,400	34.8	27.0 - 42.7	6,100	49.8	42.9 - 56.8	11,400
Asian/Pacific Islander	12.4	8.3 - 16.6	5,000	13.7	6.9 - 20.4	2,400	11.5	6.3 - 16.6	2,600
Black	5.0	4.1 - 6.0	2,000	4.5	2.9 - 6.2	800	5.4	3.7 - 7.0	1,200
Born outside the U.S.	32.8	27.6 - 37.9	14,200	23.8	16.3 - 31.4	4,400	39.6	32.7 - 46.5	9,700
Speaks non-English language at home	26.8	21.9 - 31.6	11,600	20.4	13.3 - 27.5	3,800	31.6	25.1 - 38.2	7,800
Lives in a high poverty neighborhood	28.9	24.2 - 33.7	12,100	30.3	22.4 - 38.2	5,400	27.9	21.8 - 34.0	6,700
Income as a percent of the Federal Poverty Guideline									
0-100%	33.0	28.5 - 37.6	13,300	25.8	18.6 - 33.1	4,400	38.4	31.9 - 44.8	8,900
101-200%	19.6	15.3 - 23.8	7,900	16.1	9.9 - 22.3	2,800	22.1	16.2 - 28.0	5,100
> 200%	47.4	42.7 - 52.1	19,100	58.1	49.8 - 66.3	9,900	39.5	33.0 - 46.0	9,200

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of San Diego County was 602. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

^{*} Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%)

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).