

MIHA Data Snapshot, San Francisco County by Education, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		San Francisco Co	ounty	<high school<="" th=""><th>High School Graduate</th><th>or GED</th><th>Some College</th><th>College Gradua</th><th>ate</th></high>	High School Graduate	or GED	Some College	College Gradua	ate		
Health Status before Pregnancy 1,000 1,0		% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N
In good to excellent health	tal	100	8,800	8.9 6.7 - 11.0	800	8.4 6.1 - 10.7	700	16.5 12.9 - 20.1	1,400	66.2 62.3 - 70.1	5,800
Chiomic conditions	alth Status before Pregnancy										
Diabetes	n good to excellent health	95.3 93.6 - 97.0	8,400	87.5 * 78.7 - 96.3	700	80.8 69.8 - 91.8	600	93.9 * 89.7 - 98.0	1,400		
Mysterians	Chronic conditions										
Matthmar Meight Daily folic acid use, month before pregnancy 42,6 86,6 86,8 3,00 22,6 116,937,20 200 29,9 11,1 4,15 20,0 25,6 162,94,9 40, 52,9 46,3 9,4 10,0 0 20 20 20 20 20 20 20 20 20 20 20 20	Diabetes†	2.4 1.2 - 3.6	200	5.6 * 0.2 - 11.1	<100	6.9 * 0.9 - 12.9	<100			1.7 * 0.3 - 3.1	<100
Nutrition and Weight Daily folio acid use, month before pregnancy 13,6 386 - 486 3000 22,6 115 387 200 29,9 17.4 625 200 25,6 182 349 400 52,9 46.5 98.7 Overweight before pregnancy 15,4 117 - 191 1,7 100 28,2 138 44,6 200 17.1 6.8 72 4 100 17.0 81 25.8 200 11.3 71 36.5 16.2 17.8 10.1 17.1 18.1 18.1 18.1 18.1 18.1 18.1 18	Hypertension†	2.0 0.9 - 3.1	200					4.3 * 0.3 - 8.4	<100		
Daily folic acid use, month before pregnancy 43,6 846 840 380 22,6 116 327 200 29,9 17.4 42,5 200 25,6 16.2 34.9 400 52,9 46.3 93.0 00 00 00 00 00 00 00	Asthma†	6.9 4.7 - 9.2	600	6.3 * 0.7 - 12.0	<100	14.9 * 4.2 - 25.6	100	4.3 * 1.6 - 7.1	<100	6.4 3.5 - 9.4	400
Desero before pregnancy	trition and Weight										
Does before pregnancy 9.1 6.5 1.7 8.0 2.82 13.4 46 200 17.1 6.8 2.74 2.00 1.70 8.1 2.58 2.00 4.4 2.1 6.8 1.6 2.1 2.2 2.00 3.7 2.5 3.8 3.0 3.	Daily folic acid use, month before pregnancy	43.6 38.6 - 48.6	3,800	22.6 11.6 - 33.7	200	29.9 17.4 - 42.5	200	25.6 16.2 - 34.9	400	52.9 46.3 - 59.4	3,000
Indequate weight gain during pregnancy 16.2 12.2 0.20 32.8 16.3 4.93 2.00 22.2 7.5 3.68 1.00 17.9 7.8 2.81 2.00 38.6 31.2 3.25 5.00 3.00 15.1 4.4 4.55 3.00 37.7 37.7 4.98 3.00 38.6 31.2 5.3 5.00 37.6 37.7 37.7 4.98 3.00 38.6 31.2 3.3 3.0 3.5 3.4 4.4 4.5 3.0 37.6 37.7 37.7 4.98 3.00 38.6 31.2 3.3 3.0 3.5 3.4 3.5 3.0 3.0 3.5 3.0 3.0 3.5 3.0	Overweight before pregnancy	15.4 11.7 - 19.1	1,300	29.0 15.1 - 42.9	200	21.8 10.1 - 33.4	100	24.2 12.7 - 35.7	300	11.3 7.1 - 15.6	600
Excessive weight gain during pregnancy 37,5 33 - 427 2,900 30,0 15.1 - 449 100 30,0 14.4 - 455 200 37,7 15.7 - 488 500 38,6 31.9 - 45.2	Obese before pregnancy	9.1 6.5 - 11.7	800	28.2 13.8 - 42.6	200	17.1 * 6.8 - 27.4	100	17.0 8.1 - 25.8	200	4.4 2.1 - 6.8	300
Food insecurity during pregnancy 9,2 6,9 1,15 800 36,6 23,8 4,93 300 18,8 8,5 - 29,1 100 13,0 6,0 - 19,9 200 3,3 * 1,2 - 5,3 * 1.5 * 1	nadequate weight gain during pregnancy	16.2 12.2 - 20.2	1,200	32.8 16.3 - 49.3	200	22.2 * 7.5 - 36.8	100	17.9 7.8 - 28.1	200	13.8 9.2 - 18.5	700
Priyaled are psychological IPV during pregnancy 54, 34 - 74 50, 376, 247 - 504 300 17.2 * 60 - 283 100 21.2 * 10.5 * 31.9 300 9,0 5.0 * 13.0 7.2 * 60 - 283 100 21.2 * 10.5 * 31.9 300 9,0 5.0 * 13.0 7.2 * 60 - 283 100 21.2 * 10.5 * 31.9 300 9,0 5.0 * 13.0 7.2 * 60 - 283 100 21.2 * 10.5 * 31.9 300 9,0 5.0 * 13.0 7.2 * 60 - 283 300 21.2 * 10.5 * 31.9 300 9,0 5.0 * 13.0 7.2 * 60 - 283 300 21.5 * 10.5 * 22.0 30.5 * 12.5 * 10.0 * 21.2 * 10.5 * 31.9 300 9,0 5.0 * 13.0 7.2 * 60 - 283 7.2 * 60	Excessive weight gain during pregnancy	37.5 32.3 - 42.7	2,900	30.0 15.1 - 44.9	100	30.0 14.4 - 45.5	200	37.7 25.7 - 49.8	500	38.6 31.9 - 45.2	2,000
Physical or psychological IPV during pregnancy	Food insecurity during pregnancy	9.2 6.9 - 11.5	800	36.6 23.8 - 49.3	300	18.8 8.5 - 29.1	100	13.0 6.0 - 19.9	200	3.3 * 1.2 - 5.3	200
Physical or psychological IPV during pregnancy	imate Partner Violence (IPV) and Depressive Symptoms										
Pershal degressive symptoms 12.0 86 153 100 27.0 17.0 17.0 17.0 17.0 17.0 17.0 17.0 1		5.4 3.4 - 7.4	500	17.4 * 6.7 - 28.1	100	8.5 * 0.2 - 16.8	<100	12.0 * 4.5 - 19.5	200	1.8 * 0.4 - 3.3	100
Postpartum depressive symptoms 12.0 8.6 1.53 1.00 20.9 10.2 1.15 200 17.6 7.4 27.7 100 19.1 8.2 30. 300 8.1 4.4 - 1.18 Hardships and Support during Pregnancy Experienced two or more hardships during childhood Horneless or did not have a regular place to sleep 2.8 16 4.0 200 9.5 2.0 17.0 10 10. 13.1 2.0 20 30.5 19.2 41.8 400 9.6 5.8 13.4 40.0 200 9.5 2.0 17.0 10. 10. 13.1 2.0 15.3 14. 92 41.0 1.2 0.2 20 40.0 20 4.2 20 40.0 1.2 0.2 20 40.0 1		-						-	300		500
Experienced two or more hardships during Pregnancy Experienced two or more hardships during childhood 1.1 1.5 2.0 2.0 41.3 28.1 54.5 300 21.5 11.0 32.0 20. 30.5 19.2 41.8 400 9.5 5.8 13.4 Moved due to problems paying rent or mortgage 3.1 1.7 4.5 300 10.5 30.1 9.7 3.2 30.0 16.4 6.4 2.64 100 15.8 6.1 25.4 300 8.7 4.5 30.0 20.5 30.1 9.7 3.2 30.0 16.4 6.4 2.64 100 15.8 6.1 25.4 30.0 8.7 4.5 10.0 20.0 23.4 3.0 30.5 19.2 41.8 40.0 20.0 8.7 4.5 10.0 20.0 23.4 4.5 2.5 2.5 11.1 20.0 23.5 2.5	• • •				200		100		300	8.1 4.4 - 11.8	500
Experienced two or more hardships during childhood Homeless or did not have a regular place to sleep 2.8 1.6 4.0 20 9.5 2.0 17.0 4.00 Homeless or did not have a regular place to sleep 3.1 1.7 4.5 300 10.5 3.0 17.9 4.00 Word our or partner lost job Word and or partner lost job Word and or partner lost job Word and or partner had pay or hours cut back 8.2 5.2 11.1 700 8.6 3.0 18.6 4.6 22.7 100 16.4 6.4 26.4 100 15.8 6.1 1.5 1.2 4.00 0.8 2.1 1.1 1.6 4.00 Word and or partner had pay or hours cut back 8.2 5.2 11.1 700 8.6 3.0 18.6 4.6 2.7 100 5.5 12 9.7 4.00 15.8 6.1 1.5 8.3 2.46 200 18.8 2.1 1.5 1.6 1.0 1.0 1.2 1.1 1.2 1.2 1.0 1.0 1.2 1.2 1.2 1.0 1.0 1.2 1.2 1.2 1.0 1.0 1.2 1.2 1.2 1.2 1.0 1.0 1.2 1.2 1.2 1.2 1.2 1.2 1.2 1.2 1.2 1.2											
Homeless or did not have a regular place to sleep 2.8 1.6 - 4.0 200 9.5 * 2.0 - 17.0		17 1 13 5 - 20 6	1 500	41 3 28 1 - 54 5	300	21 5 11 0 - 32 0	200	30 5 19 2 - 41 8	400	96 58 - 134	500
Moved due to problems paying rent or mortgage Moved no prattner lost job 11,7 8,2 - 15,2 1,000 21,0 00 21,0			,	-							500
Woman or partner lost job	9 , ,	_				-	-100				<100
Moman or partner had pay or hours cut back 8.2 5.2 1.11 700 8.6 ° 0.3 1.68 <100 11.2 ° 1.7 2.07 <100 16.5 ° 8.3 2.46 2.00 5.8 ° 2.1 9.5	,	-					100				500
Became separated or divorced Had no practical or emotional support Became separated or divorced Had no practical or emotional support Bustance Use Any smoking, 3 months before pregnancy Any smoking, 3 dr trimester† 1.9° 0.7° 3.1° 200 4.5° 0.3° 8.6° 1.00 3.8° 0.6° 6.9° 1.00				-						-	300
Had no practical or emotional support 2.3 1.0 3.6 200	• • • •	-	300						<100		<100
Substance Use Any smoking, 3 months before pregnancy Any smoking, postpartum 3.6 2.0 5.1 300 9.5 2.3 16.6 400 8.1 12.1 15.0 400 5.1 1.3 8.9 4100 1.8 0.1 3.5 Any binge drinking, 3 months before pregnancy Any smoking, 3 months before pregnancy Any smoking, 3 months before pregnancy Any binge drinking, 3 months before pregnancy Any binge drinking, 3 months before pregnancy Any alcohol use, 3rd trimester† 28.1 23.4 32.9 2,400		_						-		0.8 * 0.1 - 1.6	<100
Any smoking, 3 months before pregnancy Any smoking, 3 months before pregnancy Any smoking, 3rd trimester† 1,9 ° 07 ° 3.1 200 4.5 ° 0.3 ° 8.6 < 100 3.8 ° 0.6 ° 6.9 < 100											
Any smoking, 3rd trimester† 1.9 ° 0.7 ° 3.1 ° 200		8.9 6.3 - 11.5	800	10.1 * 2.9 - 17.4	<100	17.0 * 6.1 - 27.9	100	8.8 3.7 - 13.8	100	7.8 4.5 - 11.2	400
Any smoking, postpartum 3.6 2.0 - 5.1 300 9.5 * 2.3 - 16.6 <100 8.1 * 1.2 - 15.0 <100 5.1 * 1.3 - 8.9 <100 1.8 * 0.1 - 3.5 Any binge drinking, 3 months before pregnancy Any alcohol use, 3rd trimester† 28.1 23.4 - 32.9 2,400 7.5 * 0.2 - 14.9 <100 1.1.1 * 3.6 - 18.7 200 21.5 16.0 - 27.0 Any alcohol use, 3rd trimester† 8.8 1 14.5 - 21.7 1,600 27.3 15.6 - 39.1 200 26.7 14.1 - 39.3 200 30.7 19.9 - 41.4 400 12.9 8.7 - 17.1 Unsure of pregnancy intentions 11.2 8.1 - 14.3 1,000 21.0 10.5 - 31.6 200 24.3 11.8 - 36.8 200 19.5 10.2 - 28.8 300 5.6 * 2.2 - 8.9 Postpartum birth control use† Placed infant on back to sleep 8.9 5 8.6 6 - 92.4 7,800 82.7 73.0 - 92.4 600 85.7 * 75.3 - 96.0 600 75.9 66.1 85.8 1,100 71.4 65.3 - 77.5 Infant always or often shared bed 1.6 1 64.6 - 73.7 6,000 43.0 29.7 - 56.3 300 37.4 22.7 - 52.1 300 46.2 34.1 - 58.4 600 29.8 23.7 - 36.0 Intended to breastfeed exclusively, before birth 8.7 5 73.2 98.0 89.0 79.3 68.3 - 90.3 600 84.7 76.4 93.0 600 87.0 * 78.6 - 95.3 1,200 96.8 * 94.8 - 98.9 Exclusive breastfeeding, 1 month after delivery 92.8 90.5 - 95.0 8,000 79.3 68.3 - 90.3 600 84.7 76.4 - 93.0 600 87.0 * 78.6 - 95.3 1,200 96.8 * 94.8 - 98.9 Exclusive breastfeeding, 1 month after delivery 45.1 40.0 - 50.3 3,800 25.2 13.8 - 36.7 200 45.6 30.8 - 60.4 300 31.7 21.3 - 42.0 400 51.0 44.3 - 57.7				-		-			100		.00
Any binge drinking, 3 months before pregnancy Any alcohol use, 3rd trimester† 28.1 23.4 - 32.9 2,400	, ,,							5 1 * 1.3 - 8.9	<100	18* 0.1 - 3.5	100
Any alcohol use, 3rd trimester† 28.1 23.4 - 32.9 2,400 7.5 * 0.2 - 14.9 <100 39.3 32.7 - 45.9 Pregnancy Intention and Family Planning Mistimed or unwanted pregnancy 18.1 14.5 - 21.7 1,600 27.3 15.6 - 39.1 200 26.7 14.1 - 39.3 200 30.7 19.9 - 41.4 400 12.9 8.7 - 17.1 Unsure of pregnancy intentions 11.2 8.1 - 14.3 1,000 21.0 10.5 - 31.6 200 24.3 11.8 - 36.8 200 19.5 10.2 - 28.8 300 5.6 * 2.2 - 8.9 Postpartum birth control use† 74.2 69.6 - 78.7 6,400 79.8 68.5 - 91.1 600 85.7 *75.3 - 96.0 600 75.9 66.1 - 85.8 1,100 71.4 65.3 - 77.5 Infant Sleep and Breastfeeding Placed infant on back to sleep 89.5 86.6 - 92.4 7,800 82.7 73.0 - 92.4 600 86.9 *79.3 - 94.6 600 85.6 *82.2 - 93.0 1,200 91.6 87.8 - 95.3 Infant always or often shared bed 34.5 29.7 - 39.3 3,000 43.0 29.7 - 56.3 300 37.4 22.7 - 52.1 300 46.2 34.1 - 58.4 600 29.8 29.8 29.5 - 95.0 Intended to breastfeed, before birth 93.7 91.4 - 96.0 8,200 90.0 *82.1 - 97.9 700 78.4 65.8 - 91.0 600 95.5 *92.1 - 98.9 1,300 95.6 *92.9 - 98.3 Intended to breastfeed exclusively, before birth 93.7 6.40 90.0 *49.0 35.6 - 62.3 400 38.4 24.7 - 52.2 300 60.7 *48.5 72.9 800 78.5 73.2 - 83.8 Breastfeeding, 1 month after delivery 92.8 90.5 - 95.0 8,000 79.3 68.3 - 90.3 600 84.7 76.4 - 93.0 600 87.0 *78.6 - 95.3 1,200 96.8 *94.8 - 98.9 Exclusive breastfeeding, 1 month after delivery 45.1 40.0 - 50.3 3,800 25.2 13.8 - 36.7 200 45.6 30.8 - 60.4 300 31.7 21.3 - 42.0 400 51.0 44.3 - 57.7	,					-		-		_	1,200
Pregnancy Intention and Family Planning Mistimed or unwanted pregnancy 18.1 14.5 - 21.7 1,600 27.3 15.6 - 39.1 200 26.7 14.1 - 39.3 200 30.7 19.9 - 41.4 400 12.9 8.7 - 17.1 Unsure of pregnancy intentions 11.2 8.1 - 14.3 1,000 21.0 10.5 - 31.6 200 24.3 11.8 - 36.8 200 19.5 10.2 - 28.8 300 5.6 * 2.2 - 8.9 Postpartum birth control use† 74.2 69.6 - 78.7 6,400 79.8 68.5 - 91.1 600 85.7 *75.3 - 96.0 600 75.9 66.1 85.8 1,100 71.4 65.3 - 77.5 Infant Sleep and Breastfeeding Placed infant on back to sleep 89.5 86.6 - 92.4 7,800 82.7 73.0 - 92.4 600 86.9 *79.3 - 94.6 600 85.6 78.2 - 93.0 1,200 91.6 87.8 - 95.3 Infant always or often shared bed 19.3 19.4 - 96.0 8,200 90.0 * 82.1 - 97.9 700 78.4 65.8 - 91.0 600 95.5 *92.1 - 98.9 1,300 95.6 *92.9 - 98.3 Intended to breastfeed, before birth 93.7 91.4 - 96.0 8,200 90.0 *82.1 - 97.9 700 78.4 65.8 - 91.0 600 95.5 *92.1 - 98.9 1,300 95.6 *92.9 - 98.3 Intended to breastfeed exclusively, before birth 80.1 64.6 - 73.7 6,000 49.0 35.6 62.3 400 38.4 24.7 - 52.2 300 60.7 48.5 - 72.9 800 78.5 73.2 - 83.8 Breastfeeding, 1 month after delivery 92.8 90.5 - 95.0 8,000 79.3 68.3 - 90.3 600 84.7 76.4 93.0 600 87.0 *78.6 - 95.3 1,200 96.8 *94.8 - 98.9 Exclusive breastfeeding, 1 month after delivery 45.1 40.0 - 50.3 3,800 25.2 13.8 - 36.7 200 45.6 30.8 - 60.4 300 31.7 21.3 - 42.0 400 51.0 44.3 - 57.7			,								2,200
Mistimed or unwanted pregnancy 18.1 14.5 - 21.7 1,600 27.3 15.6 - 39.1 200 26.7 14.1 - 39.3 200 30.7 19.9 - 41.4 400 12.9 8.7 - 17.1 Unsure of pregnancy intentions 11.2 8.1 - 14.3 1,000 21.0 10.5 - 31.6 200 24.3 11.8 - 36.8 200 19.5 10.2 - 28.8 300 5.6 * 2.2 - 8.9 Postpartum birth control use† 74.2 69.6 - 78.7 6,400 79.8 68.5 - 91.1 600 85.7 * 75.3 - 96.0 600 75.9 66.1 - 85.8 1,100 71.4 65.3 - 77.5 Infant Sleep and Breastfeeding Placed infant on back to sleep 89.5 86.6 - 92.4 7,800 82.7 73.0 - 92.4 600 86.9 * 79.3 - 94.6 600 85.6 78.2 - 93.0 1,200 91.6 87.8 - 95.3 Infant always or often shared bed 34.5 29.7 - 39.3 3,000 43.0 29.7 - 56.3 300 37.4 22.7 - 52.1 300 46.2 34.1 - 58.4 600 29.8 23.7 - 36.0 Intended to breastfeed, before birth 93.7 91.4 - 96.0 8,200 90.0 * 82.1 - 97.9 700 78.4 65.8 - 91.0 600 95.5 * 92.1 - 98.9 1,300 95.6 * 92.9 - 98.3 Intended to breastfeed exclusively, before birth 69.1 64.6 - 73.7 6,000 49.0 35.6 - 62.3 400 38.4 24.7 - 52.2 300 60.7 48.5 - 72.9 800 78.5 73.2 - 83.8 Breastfeeding practices Any breastfeeding, 1 month after delivery 92.8 90.5 - 95.0 8,000 79.3 68.3 - 90.3 600 84.7 76.4 - 93.0 600 87.0 * 78.6 - 95.3 1,200 96.8 * 94.8 - 98.9 Exclusive breastfeeding, 1 month after delivery 45.1 40.0 - 50.3 3,800 25.2 13.8 - 36.7 200 45.6 30.8 - 60.4 300 31.7 21.3 - 42.0 400 51.0 44.3 - 57.7			,			7.0				00.0	
Unsure of pregnancy intentions 11.2 8.1 - 14.3 1,000 21.0 10.5 - 31.6 200 24.3 11.8 - 36.8 200 19.5 10.2 - 28.8 300 5.6 * 2.2 - 8.9 Postpartum birth control use† 74.2 69.6 - 78.7 6,400 79.8 68.5 - 91.1 600 85.7 * 75.3 - 96.0 600 75.9 66.1 - 85.8 1,100 71.4 65.3 - 77.5 Infant Sleep and Breastfeeding Placed infant on back to sleep 89.5 86.6 - 92.4 7,800 82.7 73.0 - 92.4 600 86.9 * 79.3 - 94.6 600 85.6 78.2 - 93.0 1,200 91.6 87.8 - 95.3 Infant always or often shared bed 34.5 29.7 - 39.3 3,000 43.0 29.7 - 56.3 300 37.4 22.7 - 52.1 300 46.2 34.1 - 58.4 600 29.8 23.7 - 36.0 Intended to breastfeed, before birth 93.7 91.4 - 96.0 8,200 90.0 * 82.1 - 97.9 700 78.4 65.8 - 91.0 600 95.5 * 92.1 - 98.9 1,300 95.6 * 92.9 - 98.3 Intended to breastfeed exclusively, before birth 69.1 64.6 - 73.7 6,000 49.0 35.6 - 62.3 400 38.4 24.7 - 52.2 300 60.7 48.5 - 72.9 800 78.5 73.2 - 83.8 Breastfeeding practices Any breastfeeding, 1 month after delivery 92.8 90.5 - 95.0 8,000 79.3 68.3 - 90.3 600 84.7 76.4 - 93.0 600 87.0 * 78.6 - 95.3 1,200 96.8 * 94.8 - 98.9 Exclusive breastfeeding, 1 month after delivery 45.1 40.0 - 50.3 3,800 25.2 13.8 - 36.7 200 45.6 30.8 - 60.4 300 31.7 21.3 - 42.0 400 51.0 44.3 - 57.7	• •	18 1 14 5 - 21 7	1 600	27 3 15 6 - 39 1	200	26 7 14 1 - 39 3	200	30 7 199 - 414	400	12 9 87 - 171	700
Postpartum birth control use† 74.2 69.6 - 78.7 6,400 79.8 68.5 - 91.1 600 85.7 * 75.3 - 96.0 600 75.9 66.1 - 85.8 1,100 71.4 65.3 - 77.5 1	, ,			_		-				-	300
Placed infant on back to sleep 89.5 86.6 - 92.4 7,800 82.7 73.0 - 92.4 600 86.9 * 79.3 - 94.6 600 85.6 78.2 - 93.0 1,200 91.6 87.8 - 95.3 Infant always or often shared bed 34.5 29.7 - 39.3 3,000 43.0 29.7 - 56.3 300 37.4 22.7 - 52.1 300 46.2 34.1 - 58.4 600 29.8 23.7 - 36.0 Intended to breastfeed, before birth 93.7 91.4 - 96.0 8,200 90.0 * 82.1 - 97.9 700 78.4 65.8 - 91.0 600 95.5 * 92.1 - 98.9 1,300 95.6 * 92.9 - 98.3 Intended to breastfeed exclusively, before birth 69.1 64.6 - 73.7 6,000 49.0 35.6 - 62.3 400 38.4 24.7 - 52.2 300 60.7 48.5 - 72.9 800 78.5 73.2 - 83.8 Breastfeeding practices Any breastfeeding, 1 month after delivery 92.8 90.5 - 95.0 8,000 79.3 68.3 - 90.3 600 84.7 76.4 - 93.0 600 87.0 * 78.6 - 95.3 1,200 96.8 * 94.8 - 98.9 Exclusive breastfeeding, 1 month after delivery 45.1 40.0 - 50.3 3,800 25.2 13.8 - 36.7 200 45.6 30.8 - 60.4 300 31.7 21.3 - 42.0 400 51.0 44.3 - 57.7			•	-							4,100
Placed infant on back to sleep 89.5 86.6 - 92.4 7,800 82.7 73.0 - 92.4 600 86.9 *79.3 - 94.6 600 85.6 78.2 - 93.0 1,200 91.6 87.8 - 95.3 1nfant always or often shared bed 34.5 29.7 - 39.3 3,000 43.0 29.7 - 56.3 300 37.4 22.7 - 52.1 300 46.2 34.1 - 58.4 600 29.8 23.7 - 36.0 1ntended to breastfeed, before birth 93.7 91.4 - 96.0 8,200 90.0 *82.1 - 97.9 700 78.4 65.8 - 91.0 600 95.5 *92.1 - 98.9 1,300 95.6 *92.9 - 98.3 1ntended to breastfeed exclusively, before birth 69.1 64.6 - 73.7 6,000 49.0 35.6 - 62.3 400 38.4 24.7 - 52.2 300 60.7 48.5 - 72.9 800 78.5 73.2 - 83.8 Breastfeeding practices Any breastfeeding, 1 month after delivery 92.8 90.5 - 95.0 8,000 79.3 68.3 - 90.3 600 84.7 76.4 - 93.0 600 87.0 *78.6 - 95.3 1,200 96.8 *94.8 - 98.9 Exclusive breastfeeding, 1 month after delivery 45.1 40.0 - 50.3 3,800 25.2 13.8 - 36.7 200 45.6 30.8 - 60.4 300 31.7 21.3 - 42.0 400 51.0 44.3 - 57.7		, II.2 00:0	-,	75.0		03.7		75.5	_,	72.1	.,
Infant always or often shared bed 34.5 29.7 - 39.3 3,000 43.0 29.7 - 56.3 300 37.4 22.7 - 52.1 300 46.2 34.1 - 58.4 600 29.8 23.7 - 36.0 lntended to breastfeed, before birth 93.7 91.4 - 96.0 8,200 90.0 * 82.1 - 97.9 700 78.4 65.8 - 91.0 600 95.5 * 92.1 - 98.9 1,300 95.6 * 92.9 - 98.3 lntended to breastfeed exclusively, before birth 69.1 64.6 - 73.7 6,000 49.0 35.6 - 62.3 400 38.4 24.7 - 52.2 300 60.7 48.5 - 72.9 800 78.5 73.2 - 83.8 Breastfeeding practices Any breastfeeding, 1 month after delivery 92.8 90.5 - 95.0 8,000 79.3 68.3 - 90.3 600 84.7 76.4 - 93.0 600 87.0 * 78.6 - 95.3 1,200 96.8 * 94.8 - 98.9 Exclusive breastfeeding, 1 month after delivery 45.1 40.0 - 50.3 3,800 25.2 13.8 - 36.7 200 45.6 30.8 - 60.4 300 31.7 21.3 - 42.0 400 51.0 44.3 - 57.7		90 E 866 - 924	7 900	92 7 72 0 - 92 4	600	96 0 * 70 2 - 04 6	600	QE 6 78 2 - 02 0	1 200	016 979 052	5,300
Intended to breastfeed, before birth 93.7 91.4 - 96.0 8,200 90.0 * 82.1 - 97.9 700 78.4 65.8 - 91.0 600 95.5 * 92.1 - 98.9 1,300 95.6 * 92.9 - 98.3 Intended to breastfeed exclusively, before birth 69.1 64.6 - 73.7 6,000 49.0 35.6 - 62.3 400 38.4 24.7 - 52.2 300 60.7 48.5 - 72.9 800 78.5 73.2 - 83.8 Breastfeeding practices Any breastfeeding, 1 month after delivery 92.8 90.5 - 95.0 8,000 79.3 68.3 - 90.3 600 84.7 76.4 - 93.0 600 87.0 * 78.6 - 95.3 1,200 96.8 * 94.8 - 98.9 Exclusive breastfeeding, 1 month after delivery 45.1 40.0 - 50.3 3,800 25.2 13.8 - 36.7 200 45.6 30.8 - 60.4 300 31.7 21.3 - 42.0 400 51.0 44.3 - 57.7	•		,						,		1,700
Intended to breastfeed exclusively, before birth 69.1 64.6 - 73.7 6,000 49.0 35.6 - 62.3 400 38.4 24.7 - 52.2 300 60.7 48.5 - 72.9 800 78.5 73.2 - 83.8 Breastfeeding practices Any breastfeeding, 1 month after delivery 92.8 90.5 - 95.0 8,000 79.3 68.3 - 90.3 600 84.7 76.4 - 93.0 600 87.0 78.6 - 95.3 1,200 96.8 94.8 - 98.9 Exclusive breastfeeding, 1 month after delivery 45.1 40.0 - 50.3 3,800 25.2 13.8 - 36.7 200 45.6 30.8 - 60.4 300 31.7 21.3 - 42.0 400 51.0 44.3 - 57.7	•		,					-			5,500
Breastfeeding practices Any breastfeeding, 1 month after delivery 92.8 90.5 - 95.0 8,000 79.3 68.3 - 90.3 600 84.7 76.4 - 93.0 600 87.0 78.6 - 95.3 1,200 96.8 94.8 - 98.9 Exclusive breastfeeding, 1 month after delivery 45.1 40.0 - 50.3 3,800 25.2 13.8 - 36.7 200 45.6 30.8 - 60.4 300 31.7 21.3 - 42.0 400 51.0 44.3 - 57.7	•		,						,		4,500
Any breastfeeding, 1 month after delivery 92.8 90.5 - 95.0 8,000 79.3 68.3 - 90.3 600 84.7 76.4 - 93.0 600 87.0 * 78.6 - 95.3 1,200 96.8 * 94.8 - 98.9 Exclusive breastfeeding, 1 month after delivery 45.1 40.0 - 50.3 3,800 25.2 13.8 - 36.7 200 45.6 30.8 - 60.4 300 31.7 21.3 - 42.0 400 51.0 44.3 - 57.7		05.1 04.0 75.7	0,000	-J.U JJ.0 02.3	400	30.7 27.7 32.2	300	00.7 40.3 72.9	000	, 0.5 / 5.2 65.6	4,500
Exclusive breastfeeding, 1 month after delivery 45.1 40.0 - 50.3 3,800 25.2 13.8 - 36.7 200 45.6 30.8 - 60.4 300 31.7 21.3 - 42.0 400 51.0 44.3 - 57.7		928 905 - 950	8,000	79 3 683 - 903	600	847 764 - 930	600	87 0 * 78 6 - 95 3	1.200	968*948-989	5,500
			,			-					2,800
THE DISCUSSION OF THE PROPERTY AND THE PROPERTY OF THE PROPERT	Any breastfeeding, 3 months after delivery	82.4 78.2 - 86.5	5,500	57.0 42.3 - 71.7	400	58.9 43.0 - 74.8	400	73.3 61.0 - 85.6	800	92.0 87.9 - 96.1	3,900
		-								44.2 36.5 - 51.9	1,900



MIHA Data Snapshot, San Francisco County by Education, 2013-2015

Maternal and Infant Health Assessment (MIHA) Survey

	San	rancisco Co	unty	y <high school<="" th=""><th colspan="3"></th><th colspan="3">Some College</th><th>Co</th><th>llege Gradua</th><th>ate</th></high>						Some College			Co	llege Gradua	ate
	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N
Service Utilization and Health Insurance Coverage															
Had a usual source of pre-pregnancy care	75.4	70.9 - 79.8	6,600	65.7	53.2 - 78.3	500	69.9	56.0 - 83.8	500	69.4	58.5 - 80.2	1,000	78.7	73.1 - 84.3	4,500
Initiated prenatal care in 1st trimester	94.5	92.6 - 96.4	8,200	80.5	68.8 - 92.3	600				87.6	81.1 - 94.1	1,300	97.8 *	* 96.2 - 99.4	5,600
Mom or infant needed but couldn't afford care postpartur	n 7.0	4.8 - 9.1	600	8.2	2.4 - 14.1	<100	14.3	* 4.0 - 24.5	100	12.1	5.5 - 18.7	200	4.4	2.0 - 6.8	200
Participated in WIC during pregnancy†	31.3	27.6 - 35.1	2,700	87.4	78.8 - 95.9	700	68.3	54.2 - 82.5	500	63.1	51.2 - 75.0	900	10.4	6.2 - 14.5	600
Received CalFresh (food stamps) during pregnancy	10.3	8.3 - 12.2	900	35.0	22.6 - 47.4	300	27.1	16.1 - 38.2	200	21.0	13.7 - 28.3	300	2.1	0.9 - 3.2	100
Pre-pregnancy insurance															
Medi-Cal	14.8	12.2 - 17.5	1,300	44.1	31.1 - 57.2	300	42.7	28.7 - 56.7	300	25.9	16.5 - 35.2	400	3.9	2.0 - 5.9	200
Uninsured	12.6	9.4 - 15.9	1,100	32.5	19.8 - 45.2	300	28.2	14.6 - 41.8	200	21.3	12.1 - 30.5	300	6.0 *	* 2.4 - 9.7	300
Private	69.8	66.2 - 73.4	6,100	13.5	4.2 - 22.8	100	26.8	12.6 - 41.1	200	51.0	39.3 - 62.8	700	88.1	83.8 - 92.4	5,100
Prenatal insurance															
Medi-Cal	31.2	27.7 - 34.8	2,700	84.4	74.5 - 94.4	600	78.8	* 66.2 - 91.4	600	59.0	47.1 - 70.9	800	10.1	6.3 - 13.9	600
Uninsured	1.7 *	0.3 - 3.0	100												
Private	65.0	61.4 - 68.6	5,700				18.2	* 6.0 - 30.5	100	35.8	24.1 - 47.4	500	87.3	83.1 - 91.5	5,000
Maternal postpartum insurance															
Medi-Cal	24.9	21.3 - 28.5	2,200	64.1	51.5 - 76.8	500	57.1	42.3 - 72.0	400	48.8	36.9 - 60.7	700	8.5	4.8 - 12.2	500
Uninsured	5.7	3.5 - 8.0	500	15.1	5.7 - 24.5	100	10.5	* 3.1 - 17.8	<100	10.7	* 3.0 - 18.5	200	2.7 *	* 0.3 - 5.2	200
Private	65.9	62.1 - 69.6	5,800				26.8	11.8 - 41.8	200	38.3	26.7 - 50.0	500	86.6	82.1 - 91.2	5,000
Infant health insurance															
Medi-Cal	33.2	29.6 - 36.9	2,900				84.1	* 72.6 - 95.7	600	57.3	45.1 - 69.4	800	12.3	8.1 - 16.6	700
Uninsured															
Private	64.4	60.7 - 68.0	5,600				12.0	* 0.8 - 23.3	<100	36.4	24.4 - 48.3	500	86.1	81.8 - 90.5	5,000
Maternal Demographics															
Unmarried	25.2	21.2 - 29.1	2,200	59.0	45.8 - 72.2	500	52.6	37.8 - 67.3	400	44.2	32.4 - 56.1	600	12.3	8.0 - 16.7	700
Race/Ethnicity															
White	41.7	37.0 - 46.5	3,600							21.3	11.1 - 31.6	300	56.7	50.1 - 63.2	3,200
Hispanic	18.8	15.2 - 22.3	1,600	67.0	54.9 - 79.1	500	44.5	30.2 - 58.8	300	26.9	15.7 - 38.0	400	7.2	3.6 - 10.7	400
Asian/Pacific Islander	33.8	28.9 - 38.7	3,000	19.7	8.3 - 31.2	200	27.2	12.7 - 41.8	200	37.2	25.3 - 49.1	500	34.9	28.6 - 41.2	2,000
Black	5.4	4.7 - 6.2	500	12.0	5.8 - 18.2	<100	13.8	7.6 - 20.1	100	14.2	9.4 - 19.1	200	1.2	0.6 - 1.8	<100
Born outside the U.S.	45.5	40.7 - 50.4	4,000	78.5	68.9 - 88.2	600	67.6	54.9 - 80.3	500	47.7	35.6 - 59.8	700	37.3	30.9 - 43.7	2,100
Speaks non-English language at home	31.0	26.7 - 35.3	2,700	76.7	66.7 - 86.7	600	54.8	40.6 - 69.1	400	43.3	31.0 - 55.7	600	18.0	13.0 - 23.0	1,000
Lives in a high poverty neighborhood	12.0	9.0 - 15.0	1,000	24.8	13.4 - 36.2	200	29.8	17.2 - 42.5	200	24.8	13.9 - 35.6	300	5.0	2.2 - 7.8	300
Income as a percent of the Federal Poverty Guideline															
0-100%	20.1	16.9 - 23.3	1,700	71.5	57.8 - 85.2	500	59.9	43.9 - 75.9	400	31.7	22.1 - 41.4	400	6.5	2.9 - 10.1	400
101-200%	12.1	9.1 - 15.1	1,000	25.6	12.4 - 38.8	200	18.4	* 7.5 - 29.2	100	31.6	20.0 - 43.2	400	5.0	2.3 - 7.7	300
> 200%	67.8	64.1 - 71.5	5,600				21.7	* 5.7 - 37.8	100	36.7	23.9 - 49.5	500	88.5	84.3 - 92.8	5,000

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of San Francisco County was 594. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

^{*} Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).

Geographical Area Snapshot is not provided due to suppression criteria.

This page is intentionally left blank



MIHA Data Snapshot, San Francisco County by Income, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	San Francisco Cou	0-100% Federal Poverty	Guideline	101-200% Federal Poverty	Guideline	> 200% Federal Poverty	Guideline	
	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N
Total	100	8,800	20.1 16.9 - 23.3	1,700	12.1 9.1 - 15.1	1,000	67.8 64.1 - 71.5	5,600
Health Status before Pregnancy								
In good to excellent health	95.3 93.6 - 97.0	8,400	85.6 79.4 - 91.7	1,400	93.1 * 86.7 - 99.5	900		
Chronic conditions								
Diabetes†	2.4 1.2 - 3.6	200	4.0 * 1.2 - 6.9	<100	6.9 * 1.1 - 12.7	<100		
Hypertension†	2.0 0.9 - 3.1	200	4.5 * 0.9 - 8.1	<100				
Asthma†	6.9 4.7 - 9.2	600	10.5 5.3 - 15.7	200	4.4 * 0.6 - 8.2	<100	6.5 3.5 - 9.4	400
Nutrition and Weight								
Daily folic acid use, month before pregnancy	43.6 38.6 - 48.6	3,800	31.5 21.8 - 41.3	500	33.8 21.1 - 46.5	300	50.0 43.3 - 56.7	2,800
Overweight before pregnancy	15.4 11.7 - 19.1	1,300	26.3 17.6 - 34.9	400	25.1 11.9 - 38.3	200	10.7 6.2 - 15.2	600
Obese before pregnancy	9.1 6.5 - 11.7	800	15.2 8.9 - 21.4	200	17.1 * 6.9 - 27.4	200	6.0 3.0 - 8.9	300
Inadequate weight gain during pregnancy	16.2 12.2 - 20.2	1,200	25.1 15.0 - 35.3	300	14.4 * 4.7 - 24.1	100	14.2 9.3 - 19.1	700
Excessive weight gain during pregnancy	37. 5 32.3 - 42.7	2,900	35.7 25.6 - 45.8	500	44.1 29.5 - 58.7	400	38.1 31.3 - 44.8	2,000
Food insecurity during pregnancy	9.2 6.9 - 11.5	800	29.5 21.3 - 37.6	500	25.2 12.9 - 37.4	300		
Intimate Partner Violence (IPV) and Depressive Symptoms								
Physical or psychological IPV during pregnancy	5.4 3.4 - 7.4	500	14.4 7.4 - 21.4	200	19.1 8.4 - 29.9	200		
Prenatal depressive symptoms	14.4 10.8 - 17.9	1,300	23.4 15.9 - 31.0	400	30.4 17.3 - 43.5	300	8.4 4.2 - 12.6	500
Postpartum depressive symptoms	12.0 8.6 - 15.3	1,000	17.3 11.0 - 23.6	300	21. 5 9.9 - 33.0	200	7.8 3.7 - 11.8	400
Hardships and Support during Pregnancy								
Experienced two or more hardships during childhood	17.1 13.5 - 20.6	1,500	30.9 22.9 - 39.0	500	18.5 8.7 - 28.3	200	12.9 8.3 - 17.6	700
Homeless or did not have a regular place to sleep	2.8 1.6 - 4.0	200	10.0 5.0 - 14.9	200				
Moved due to problems paying rent or mortgage	3.1 1.7 - 4.5	300	8.2 4.2 - 12.1	100				
Woman or partner lost job	11.7 8.2 - 15.2	1,000	22.5 12.7 - 32.3	400	13.1 * 4.5 - 21.6	100	7.4 3.5 - 11.2	400
Woman or partner had pay or hours cut back	8.2 5.2 - 11.1	700	18.1 8.3 - 27.9	300	16.7 * 6.1 - 27.2	200	4.0 * 1.3 - 6.7	200
Became separated or divorced	3.1 1.9 - 4.3	300	9.4 4.9 - 14.0	200				
Had no practical or emotional support	2.3 1.0 - 3.6	200	5.9 * 1.9 - 10.0	<100				
Substance Use								
Any smoking, 3 months before pregnancy	8.9 6.3 - 11.5	800	16.3 9.1 - 23.5	300			7.7 4.4 - 10.9	400
Any smoking, 3rd trimester†	1.9 * 0.7 - 3.1	200	4.6 1.9 - 7.3	<100				
Any smoking, postpartum	3.6 2.0 - 5.1	300	9.8 5.2 - 14.5	200			1.9 * 0.1 - 3.7	100
Any binge drinking, 3 months before pregnancy	18.1 14.1 - 22.0	1,600	7.6 3.9 - 11.2	100	10.7 * 1.6 - 19.9	100	23.3 17.6 - 29.1	1,300
Any alcohol use, 3rd trimester†	28.1 23.4 - 32.9	2,400					38.1 31.5 - 44.8	2,100
Pregnancy Intention and Family Planning								
Mistimed or unwanted pregnancy	18.1 14.5 - 21.7	1,600	36.7 27.5 - 45.9	600	24.4 12.4 - 36.4	200	11.6 7.5 - 15.7	700
Unsure of pregnancy intentions	11.2 8.1 - 14.3	1,000	20.1 12.7 - 27.5	300	16.4 6.8 - 26.0	200	6.9 3.1 - 10.6	400
Postpartum birth control use†	74.2 69.6 - 78.7	6,400	78.3 68.6 - 88.0	1,300	78.8 67.2 - 90.4	800	71.9 65.9 - 77.9	4,000
Infant Sleep and Breastfeeding								
Placed infant on back to sleep	89.5 86.6 - 92.4	7,800	84.8 78.8 - 90.8	1,400	85.0 76.4 - 93.5	800	92.4 88.6 - 96.1	5,200
Infant always or often shared bed	34.5 29.7 - 39.3	3,000	44.7 35.0 - 54.4	700	45.5 31.5 - 59.5	400	29.8 23.5 - 36.1	1,700
Intended to breastfeed, before birth	93.7 91.4 - 96.0	8,200	88.0 81.5 - 94.6	1,400	95.9 * 92.2 - 99.5	900	96.3 94.2 - 98.4	5,400
Intended to breastfeed exclusively, before birth	69.1 64.6 - 73.7	6,000	56.2 46.6 - 65.8	900	50.3 36.3 - 64.3	500	78.4 72.7 - 84.0	4,400
Breastfeeding practices								
Any breastfeeding, 1 month after delivery	92.8 90.5 - 95.0	8,000	84.7 78.5 - 91.0	1,400	85.8 * 77.4 - 94.2	800	96.9 * 94.6 - 99.2	5,400
Exclusive breastfeeding, 1 month after delivery	45.1 40.0 - 50.3	3,800	39.4 28.8 - 50.0	600	30.4 18.0 - 42.8	300	49.9 43.2 - 56.7	2,700
Any breastfeeding, 3 months after delivery	82.4 78.2 - 86.5	5,500	67.3 57.6 - 77.1	900	56.7 40.9 - 72.4	400	93.2 89.3 - 97.2	3,900
Exclusive breastfeeding, 3 months after delivery	35.6 30.0 - 41.1	2,300	16.6 8.7 - 24.5	200	16.3 6.9 - 25.8	100	45.7 38.0 - 53.4	1,900



MIHA Data Snapshot, San Francisco County by Income, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	San Francisco County			0-100% Fede	ral Poverty	Guideline	101-200% F	ederal Povert	y Guideline	> 200% Fe	deral Poverty	Guideline
	%	95% CI	N	% 9	95% CI	N	%	95% CI	N	%	95% CI	N
Service Utilization and Health Insurance Coverage												
Had a usual source of pre-pregnancy care	75.4	70.9 - 79.8	6,600	73.7 65.8	- 81.6	1,200	64.8	51.5 - 78.0	700	78.9	73.1 - 84.7	4,400
Initiated prenatal care in 1st trimester	94.5	92.6 - 96.4	8,200	89.0 83.6	- 94.5	1,400	89.8 *	81.6 - 97.9	900	97.8 *	96.1 - 99.5	5,500
Mom or infant needed but couldn't afford care postpartum	7.0	4.8 - 9.1	600	12.5 7.7	- 17.3	200	19.1	8.1 - 30.1	200	3.0 *	0.8 - 5.2	200
Participated in WIC during pregnancy†	31.3	27.6 - 35.1	2,700	84.8 77.6	- 92.0	1,400	78.6	67.5 - 89.6	800	4.9 *	1.6 - 8.3	300
Received CalFresh (food stamps) during pregnancy	10.3	8.3 - 12.2	900	38.3 29.6	- 47.1	600	16.9	8.7 - 25.1	200			
Pre-pregnancy insurance												
Medi-Cal	14.8	12.2 - 17.5	1,300	47.9 38.2	57.7	800	28.8	16.4 - 41.2	300			
Uninsured	12.6	9.4 - 15.9	1,100	31.8 21.6	- 42.0	500	33.0	19.7 - 46.4	300	2.0 *	0.3 - 3.7	100
Private	69.8	66.2 - 73.4	6,100	12.6 5.9	- 19.2	200	32.4	19.5 - 45.2	300	96.8*	94.4 - 99.2	5,400
Prenatal insurance												
Medi-Cal	31.2	27.7 - 34.8	2,700	88.2 81.4	- 95.0	1,500	74.0	61.6 - 86.3	700	4.9 *	1.8 - 8.0	300
Uninsured	1.7	* 0.3 - 3.0	100									
Private	65.0	61.4 - 68.6	5,700	4.9 * 1.4	- 8.5	<100	15.8 *	5.9 - 25.6	200	94.1	90.6 - 97.5	5,300
Maternal postpartum insurance												
Medi-Cal	24.9	21.3 - 28.5	2,200	77.3 70.0	- 84.5	1,300	46.8	32.8 - 60.7	500	3.9 *	1.0 - 6.7	200
Uninsured	5.7	3.5 - 8.0	500	9.4 4.8	- 14.0	200	27.5	13.7 - 41.3	300			
Private	65.9	62.1 - 69.6	5,800	5.6 * 1.9	- 9.2	<100	22.0	10.9 - 33.1	200	94.4 *	90.9 - 97.9	5,300
Infant health insurance												
Medi-Cal	33.2	29.6 - 36.9	2,900	94.2 * 90.8	- 97.7	1,500	79.0	66.8 - 91.2	800	5.8	2.5 - 9.2	300
Uninsured												
Private	64.4	60.7 - 68.0	5,600	3.4 * 0.5	- 6.2	<100	13.2 *	3.5 - 22.8	100	92.6	88.9 - 96.4	5,200
Maternal Demographics												
Did not complete high school (or GED)	8.9	6.7 - 11.0	800	27.6 19.3	- 36.0	500	16.4	7.2 - 25.7	200			
Unmarried	25.2	21.2 - 29.1	2,200	56.2 46.1	- 66.3	900	32.2	20.1 - 44.3	300	13.5	8.5 - 18.4	800
Race/Ethnicity												
White	41.7	37.0 - 46.5	3,600	12.5 * 2.1	- 22.9	200	17.7	7.4 - 28.0	200	56.6	49.9 - 63.3	3,200
Hispanic	18.8	15.2 - 22.3	1,600	42.5 33.3	- 51.7	700	21.9	11.2 - 32.7	200	8.7	4.4 - 13.1	500
Asian/Pacific Islander	33.8	28.9 - 38.7	3,000	26.6 18.3	- 35.0	400	52.2	38.5 - 65.8	500	33.2	26.8 - 39.6	1,900
Black	5.4	4.7 - 6.2	500	17.8 13.3	- 22.3	300	8.2	3.9 - 12.5	<100	1.2	0.6 - 1.9	<100
Born outside the U.S.	45.5	40.7 - 50.4	4,000	61.9 52.5	- 71.3	1,000	71.2	59.6 - 82.7	700	33.5	27.0 - 40.0	1,900
Speaks non-English language at home	31.0	26.7 - 35.3	2,700	56.1 46.5	- 65.8	900	66.8	54.6 - 79.1	700	15.0	9.9 - 20.1	800
Lives in a high poverty neighborhood	12.0	9.0 - 15.0	1,000	29.5 21.0	- 38.0	500	21.5	10.1 - 32.9	200	4.9 *	1.7 - 8.1	300

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of San Francisco County was 594. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

^{*} Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



MIHA Data Snapshot, San Francisco County by Maternal Age, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	San Francisco Cou	ınty	15-19 Years		20-34 Years		35 Years or Olde	er
	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N
Total	100	8,800	1.1 * 0.5 - 1.8	100	60.6 55.7 - 65.6	5,300	38.2 33.3 - 43.2	3,400
Health Status before Pregnancy								
In good to excellent health	95.3 93.6 - 97.0	8,400			93.0 90.3 - 95.8	4,900		
Chronic conditions								
Diabetes†	2.4 1.2 - 3.6	200			2.5 1.1 - 4.0	100	2.3 * 0.2 - 4.5	<100
Hypertension†	2.0 0.9 - 3.1	200			1.8 * 0.5 - 3.1	<100	2.4 * 0.3 - 4.5	<100
Asthma†	6.9 4.7 - 9.2	600			6.7 4.3 - 9.2	400	6.3 * 2.1 - 10.4	200
Nutrition and Weight								
Daily folic acid use, month before pregnancy	43.6 38.6 - 48.6	3,800			44.0 37.8 - 50.2	2,300	44.0 35.3 - 52.7	1,500
Overweight before pregnancy	15.4 11.7 - 19.1	1,300	35.9 * 3.1 - 68.6	<100	15.1 10.4 - 19.7	800	15.2 8.8 - 21.7	500
Obese before pregnancy	9.1 6.5 - 11.7	800			10.1 6.8 - 13.4	500	7.8 3.5 - 12.2	300
Inadequate weight gain during pregnancy	16.2 12.2 - 20.2	1,200			17.6 12.5 - 22.8	800	14.3 7.9 - 20.6	400
Excessive weight gain during pregnancy	37.5 32.3 - 42.7	2,900	50.8 * 20.2 - 81.5	<100	36.8 30.4 - 43.1	1,700	38.3 29.2 - 47.4	1,200
Food insecurity during pregnancy	9.2 6.9 - 11.5	800			12.4 8.9 - 16.0	700	3.6 * 1.3 - 5.9	100
Intimate Partner Violence (IPV) and Depressive Symptoms								
Physical or psychological IPV during pregnancy	5.4 3.4 - 7.4	500			7.2 4.2 - 10.2	400	2.0 * 0.2 - 3.8	<100
Prenatal depressive symptoms	14.4 10.8 - 17.9	1,300			14.9 10.8 - 18.9	800	13.6 6.8 - 20.4	500
Postpartum depressive symptoms	12.0 8.6 - 15.3	1,000			13.6 9.5 - 17.7	700	9.5 * 3.6 - 15.4	300
Hardships and Support during Pregnancy								
Experienced two or more hardships during childhood	17.1 13.5 - 20.6	1,500	55.3 * 24.8 - 85.9	<100	17.9 13.6 - 22.2	900	14.6 8.1 - 21.1	500
Homeless or did not have a regular place to sleep	2.8 1.6 - 4.0	200			3.4 1.7 - 5.1	200		
Moved due to problems paying rent or mortgage	3.1 1.7 - 4.5	300			3.8 1.9 - 5.7	200		
Woman or partner lost job	11.7 8.2 - 15.2	1,000			10.4 6.5 - 14.4	500	13.3 6.8 - 19.9	400
Woman or partner had pay or hours cut back	8.2 5.2 - 11.1	700			8.7 4.7 - 12.6	500	7.4 * 2.8 - 11.9	200
Became separated or divorced	3.1 1.9 - 4.3	300			4.0 2.2 - 5.8	200		
Had no practical or emotional support	2.3 1.0 - 3.6	200			3.4 * 1.4 - 5.4	200		
Substance Use	0.0				40.4		70	
Any smoking, 3 months before pregnancy	8.9 6.3 - 11.5	800			10.1 6.6 - 13.6	500	7.0 3.1 - 10.9	200
Any smoking, 3rd trimester†	1.9 * 0.7 - 3.1	200			1.8 0.8 - 2.8	<100		
Any smoking, postpartum	3.6 2.0 - 5.1	300			3.9 2.1 - 5.7	200	17.2 10.6 24.0	600
Any binge drinking, 3 months before pregnancy Any alcohol use, 3rd trimester†	18.1 14.1 - 22.0 28.1 23.4 - 32.9	1,600 2,400			18.6 13.6 - 23.5 23.7 17.8 - 29.6	1,000 1,300	17.3 10.6 - 24.0 36.1 27.6 - 44.6	600 1,200
·	28.1 23.4 - 32.9	2,400			23.7 17.8 - 29.0	1,300	30.1 27.0 - 44.0	1,200
Pregnancy Intention and Family Planning	40.4.4.5. 24.7	4.500	24.0 * 4.0	400	24.0 40.7 20.4	4 200	74 22 444	200
Mistimed or unwanted pregnancy Unsure of pregnancy intentions	18.1 14.5 - 21.7	1,600	24.8 * 4.8 - 44.7	<100	24.9 19.7 - 30.1	1,300	7.1 3.2 - 11.1	200
Postpartum birth control use†	11.2 8.1 - 14.3 74.2 69.6 - 78.7	1,000 6,400	57.3 * 30.3 - 84.4 	<100	11.8 7.9 - 15.7 80.2 75.1 - 85.3	600 4,200	8.8 3.7 - 13.9 64.4 56.0 - 72.8	300 2,100
·	74.2 09.0 - 76.7	0,400			00.2 /5.1 - 65.5	4,200	04.4 30.0 - 72.8	2,100
Infant Sleep and Breastfeeding	00 5 000		72.7****	400	00.4 05.4 00.7	. =	00.4 0.4	2 222
Placed infant on back to sleep Infant always or often shared bed	89.5 86.6 - 92.4	7,800	73.7 * 49.9 - 97.4	<100	89.4 86.1 - 92.7	4,700	90.1 84.6 - 95.6	3,000
Intended to breastfeed, before birth	34.5 29.7 - 39.3 93.7 91.4 - 96.0	3,000	23.1 * 1.9 - 44.2	<100	36.7 30.6 - 42.9 94.3 91.7 - 96.8	1,900	31.4 23.5 - 39.3 92.9 * 88.6 - 97.2	1,100
Intended to breastfeed, before birth		8,200	 30.0 * 6.1 - 54.0	~100		5,000		3,100
Breastfeeding practices	69.1 64.6 - 73.7	6,000	SU.U 0.1 - 54.U	<100	68.0 62.4 - 73.6	3,600	72.1 64.1 - 80.1	2,400
Any breastfeeding, 1 month after delivery	92.8 90.5 - 95.0	8,000	68.4 * 40.8 - 96.1	<100	92.4 89.8 - 95.0	4,800	94.0 * 89.9 - 98.2	3,100
Exclusive breastfeeding, 1 month after delivery	45.1 40.0 - 50.3	3,800	46.6 * 15.7 - 77.4	<100	44.5 38.1 - 50.9	2,300	46.0 37.3 - 54.8	1,500
Any breastfeeding, 3 months after delivery	82.4 78.2 - 86.5	5,500	40.0 15.7 - 77.4	/100	78.4 72.7 - 84.1	3,100	89.9 84.1 - 95.7	2,400
Exclusive breastfeeding, 3 months after delivery	35.6 30.0 - 41.1	2,300			30.3 23.7 - 36.8	1,200	44.1 34.4 - 53.9	1,200
Exclasive breastreeding, 5 months after delivery	JJ.U JU.U - 41.1	2,300	==		JU.J 23.7 - 30.0	1,200	44.1 J4.4 - J3.3	1,200



MIHA Data Snapshot, San Francisco County by Maternal Age, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	San Francisco Co	unty	15-19 Years		20-34 Years		35 Years or Olde	er
	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N
Service Utilization and Health Insurance Coverage								
Had a usual source of pre-pregnancy care	75.4 70.9 - 79.8	6,600	57.4 * 26.3 - 88.4	<100	72. 9 67.4 - 78.4	3,900	79.9 72.5 - 87.2	2,700
Initiated prenatal care in 1st trimester	94.5 92.6 - 96.4	8,200			93.1 90.4 - 95.7	4,900	96.8 * 94.2 - 99.4	3,200
Mom or infant needed but couldn't afford care postpartum	7.0 4.8 - 9.1	600			9.2 6.1 - 12.2	500	3.4 * 0.5 - 6.3	100
Participated in WIC during pregnancy†	31.3 27.6 - 35.1	2,700	74.5 * 50.1 - 98.8	<100	41.2 35.8 - 46.6	2,100	14.2 8.1 - 20.3	500
Received CalFresh (food stamps) during pregnancy	10.3 8.3 - 12.2	900	44.0 * 15.0 - 73.0	<100	12.9 10.1 - 15.7	700	5.0 2.3 - 7.8	200
Pre-pregnancy insurance								
Medi-Cal	14.8 12.2 - 17.5	1,300	73.0 * 46.9 - 99.1	<100	18.0 14.2 - 21.8	1,000	8.1 4.3 - 11.8	300
Uninsured	12.6 9.4 - 15.9	1,100			18.5 13.6 - 23.5	1,000	3.3 * 0.5 - 6.2	100
Private	69.8 66.2 - 73.4	6,100			59.8 54.4 - 65.2	3,200	87.1 82.3 - 91.9	2,900
Prenatal insurance								
Medi-Cal	31.2 27.7 - 34.8	2,700			39.6 34.3 - 45.0	2,100	16.1 10.3 - 21.8	500
Uninsured	1.7 * 0.3 - 3.0	100			2.8 * 0.6 - 5.0	100		
Private	65.0 61.4 - 68.6	5,700			54.2 48.8 - 59.6	2,900	83.9 78.2 - 89.7	2,800
Maternal postpartum insurance								
Medi-Cal	24.9 21.3 - 28.5	2,200			31.8 26.6 - 37.1	1,700	11.9 6.6 - 17.2	400
Uninsured	5.7 3.5 - 8.0	500			6.7 3.8 - 9.7	400	4.2 * 0.5 - 8.0	100
Private	65.9 62.1 - 69.6	5,800			57.6 52.3 - 63.0	3,100	80.9 74.3 - 87.4	2,700
Infant health insurance								
Medi-Cal	33.2 29.6 - 36.9	2,900			42.5 37.2 - 47.8	2,200	17.2 11.1 - 23.3	600
Uninsured								
Private	64.4 60.7 - 68.0	5,600			54.9 49.5 - 60.2	2,900	81.1 74.8 - 87.3	2,700
Maternal Demographics								
Did not complete high school (or GED)	8.9 6.7 - 11.0	800	29.5 * 2.3 - 56.7	<100	10.8 7.8 - 13.8	600	5.2 * 2.0 - 8.5	200
Unmarried	25.2 21.2 - 29.1	2,200			27.1 22.3 - 31.9	1,400	20.2 13.0 - 27.4	700
Race/Ethnicity								
White	41.7 37.0 - 46.5	3,600			36.5 30.6 - 42.4	1,900	51.5 42.7 - 60.2	1,700
Hispanic	18.8 15.2 - 22.3	1,600	37.4 * 6.4 - 68.4	<100	21.9 17.2 - 26.6	1,200	13.1 7.2 - 19.1	400
Asian/Pacific Islander	33.8 28.9 - 38.7	3,000			34.9 28.8 - 41.1	1,900	32.8 24.5 - 41.1	1,100
Black	5.4 4.7 - 6.2	500	53.1 * 22.9 - 83.3	<100	6.3 5.0 - 7.6	300	2.6 1.3 - 3.9	<100
Born outside the U.S.	45.5 40.7 - 50.4	4,000	15.9 * 2.8 - 29.0	<100	47.4 41.4 - 53.4	2,500	43.5 34.6 - 52.3	1,500
Speaks non-English language at home	31.0 26.7 - 35.3	2,700	14.1 * 1.9 - 26.4	<100	37.2 31.6 - 42.9	1,900	21.4 13.9 - 29.0	700
Lives in a high poverty neighborhood	12.0 9.0 - 15.0	1,000	54.2 * 23.8 - 84.6	<100	13.3 9.5 - 17.1	700	8.8 * 3.6 - 14.0	300

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of San Francisco County was 594. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

^{*} Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



MIHA Data Snapshot, San Francisco County by Neighborhood Poverty, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	San Francisco County 95% CI N				of Reside w Poverty			9% of Reside elow Povert		10-19.9% of Resi below Pover		≥ 20% of Reside below Povert	
	% 959	%CI N	%	9	5% CI	N	%	95% CI	N	% 95% CI	N	% 95% CI	N
Total	100	8,	800 8	3.8 6.	0 - 11.6	800	40.6	35.6 - 45.6	3,500	38.6 33.7 - 43.5	3,300	12.0 9.0 - 15.0	1,000
Health Status before Pregnancy													
In good to excellent health Chronic conditions	95.3 93.6	97.0 8,	400				97.3*	94.8 - 99.9	3,400	95.0 92.2 - 97.7	3,200	87.8 * 80.5 - 95.1	900
Diabetes†	2.4 1.2	3.6	200				2.4 *	0.4 - 4.4	<100	2.3 * 0.5 - 4.0	<100		
Hypertension†										2.3 * 0.3 - 4.3	<100		
Asthma†							6.2	2.6 - 9.8	200	7.6 4.0 - 11.3	200	9.1 * 1.9 - 16.4	<100
Nutrition and Weight													
Daily folic acid use, month before pregnancy	43.6 38.6	48.6 3,	800 39	9.0 23.	3 - 54.7	300	49.7	41.4 - 58.1	1,700	45.7 37.8 - 53.7	1,500	24.7 13.4 - 36.0	300
Overweight before pregnancy	15.4 11.7	19.1 1,	300 14	1.5 * 4.	0 - 24.9	100	10.7	4.9 - 16.4	400	17.9 11.9 - 23.9	600	22.4 * 8.5 - 36.3	200
Obese before pregnancy	9.1 6.5						8.9	4.2 - 13.6	300	9.5 5.4 - 13.6	300	15.3 7.4 - 23.1	100
Inadequate weight gain during pregnancy	16.2 12.2		200					11.0 - 24.5	500	14.6 8.9 - 20.3	400	19.1 * 6.9 - 31.3	200
Excessive weight gain during pregnancy	37.5 32.3	42.7 2,	900 34	1.5 18.	3 - 50.6	200	33.9	25.5 - 42.2	1,000	41.9 33.6 - 50.3	1,200	36.6 21.9 - 51.3	300
Food insecurity during pregnancy	9.2 6.9		_					3.2 - 9.2	200	9.2 5.1 - 13.2	300	24.3 13.5 - 35.2	200
Intimate Partner Violence (IPV) and Depressive Symptoms													
Physical or psychological IPV during pregnancy	5.4 3.4	7.4	500				2.3 *	0.4 - 4.3	<100	5.0 * 1.7 - 8.3	200	17.4 * 6.9 - 27.8	200
Prenatal depressive symptoms	14.4 10.8	17.9 1,	300 17	7.3 * 5.	3 - 29.3	100	10.0	5.0 - 15.0	300	17.4 11.3 - 23.6	600	17.6 * 5.3 - 29.9	200
Postpartum depressive symptoms	12.0 8.6	15.3 1,		-	7 - 31.5	100	8.4	3.6 - 13.2	300	12.3 7.5 - 17.1	400	19.1 * 6.0 - 32.3	200
Hardships and Support during Pregnancy													
Experienced two or more hardships during childhood	17.1 13.5	20.6 1	500 8	3 .2 * 0.	2 - 16.1	<100	12.9	7.3 - 18.4	400	16.6 11.5 - 21.6	500	35.8 22.1 - 49.4	400
Homeless or did not have a regular place to sleep					_ 10.1	1200	2.7 *		<100	2.7 * 0.7 - 4.7	<100	4.2 * 0.6 - 7.8	<100
Moved due to problems paying rent or mortgage	_							1.0 - 6.6	100	2.8 * 1.0 - 4.7	<100		
Woman or partner lost job	11.7 8.2							5.7 - 16.0	400	11.7 5.8 - 17.6	400	20.7 * 7.5 - 33.9	200
Woman or partner had pay or hours cut back	8.2 5.2							2.0 - 10.4	200	9.3 * 3.7 - 14.9	300	16.6 * 6.8 - 26.3	200
Became separated or divorced	3.1 1.9						2.3 *		<100	2.8 * 0.8 - 4.9	<100	6.4 * 1.9 - 10.9	<100
Had no practical or emotional support	_									4.0 * 1.1 - 7.0	100		
Substance Use													
Any smoking, 3 months before pregnancy	8.9 6.3	11.5	800				5.9	2.6 - 9.2	200	11.1 6.2 - 15.9	400	11.7 * 3.4 - 19.9	100
Any smoking, 3rd trimester†	1.9 * 0.7							2.0 3.2	200	2.1 * 0.3 - 3.9	<100		100
Any smoking, postpartum	_									5.0 2.1 - 7.8	200	9.2 * 1.4 - 16.9	<100
Any binge drinking, 3 months before pregnancy	18.1 14.1) 9 * 7.	8 - 34.1	200	19 5	12.5 - 26.5	700	19.3 12.9 - 25.6	600	9.1 * 3.0 - 15.1	<100
Any alcohol use, 3rd trimester†	28.1 23.4			-	8 - 63.5	400		20.1 - 36.1	1,000	30.1 22.2 - 38.0	1,000	7.3 * 0.8 - 13.9	<100
Pregnancy Intention and Family Planning		,							,	00.2	,	,,,	
Mistimed or unwanted pregnancy	18.1 14.5	217 1	600 13	37* 2	8 - 24.6	100	11.0	6.3 - 15.7	400	24.1 17.4 - 30.8	800	22.3 12.7 - 32.0	200
Unsure of pregnancy intentions	11.2 8.1	,			0 24.0	100	9.7	4.6 - 14.7	300	9.1 4.3 - 13.8	300	24.1 13.4 - 34.9	200
Postpartum birth control use†	74.2 69.6			5 58.	1 - 86.8	500		66.4 - 80.9	2,500	70.6 62.7 - 78.5	2,300	85.6 * 76.7 - 94.5	900
Infant Sleep and Breastfeeding	, 1.2	, , ,	,,,				73.0		_,	70.0	_,	03.0	
Placed infant on back to sleep	89.5 86.6	02/1 7	800				99.6	83.6 - 93.6	3,100	87.9 82.5 - 93.2	2,800	89.8 84.0 - 95.5	900
Infant always or often shared bed	34.5 29.7				6 - 48.2	200		28.1 - 44.6	1,300	32.1 24.8 - 39.3	1,000	29.7 18.0 - 41.3	300
Intended to breastfeed, before birth	93.7 91.4	,		<u>/.9</u> 1/. 	0 +0.2	200		89.2 - 97.3	3,300	95.4 * 92.4 - 98.3	3,100	88.8 * 81.3 - 96.4	900
Intended to breastfeed exclusively, before birth	69.1 64.6	,			8 - 90.2	600		61.0 - 76.2	2,400	75.6 69.0 - 82.3	2,500	47.6 34.2 - 61.0	500
Breastfeeding practices	05.1 04.0	. 5. ,	,,	,, J 00.	50.2	000	00.0	02.0 70.2	2,400	, 5.0 05.0 02.5	2,500	77.0 54.2 01.0	300
Any breastfeeding, 1 month after delivery	92.8 90.5	95.0 8,	000				95.2*	91.4 - 99.1	3,300	92.2 88.6 - 95.8	3,000	80.9 72.3 - 89.5	800
Exclusive breastfeeding, 1 month after delivery	45.1 40.0	50.3 3,	800 43	3.5 27.	0 - 60.0	300	43.5	34.9 - 52.0	1,500	48.2 39.9 - 56.4	1,500	41.0 27.6 - 54.5	400
Any breastfeeding, 3 months after delivery	82.4 78.2						87.4	81.0 - 93.8	2,300	82.8 76.2 - 89.3	2,000	63.9 50.6 - 77.3	600
Exclusive breastfeeding, 3 months after delivery	35.6 30.0	41.1 2,	300 31	L. 3 13.	8 - 48.7	200	33.0	24.0 - 42.0	900	39.9 30.9 - 48.9	1,000	31.9 17.5 - 46.3	300



MIHA Data Snapshot, San Francisco County by Neighborhood Poverty, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	San Francisco Co	unty	0-4.9% of Reside below Poverty		5-9.9% of Reside		10-19.9% of Resid		≥ 20% of Reside below Poverty	
	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N
Service Utilization and Health Insurance Coverage										
Had a usual source of pre-pregnancy care	75.4 70.9 - 79.8	6,600	79.5 * 66.1 - 92.9	600	75.2 68.0 - 82.4	2,600	75.7 69.0 - 82.5	2,500	79.6 68.5 - 90.6	800
Initiated prenatal care in 1st trimester	94.5 92.6 - 96.4	8,200			95.4 * 92.3 - 98.5	3,200	92.2 88.5 - 95.8	3,000	94.4 * 90.2 - 98.6	1,000
Mom or infant needed but couldn't afford care postpartun	7.0 4.8 - 9.1	600			2.4 * 0.2 - 4.6	<100	10.7 6.5 - 14.9	300	13.8 * 4.4 - 23.3	100
Participated in WIC during pregnancy†	31.3 27.6 - 35.1	2,700	19.7 * 6.8 - 32.6	100	17.8 12.2 - 23.4	600	37.4 30.3 - 44.4	1,200	65.8 52.9 - 78.7	700
Received CalFresh (food stamps) during pregnancy	10.3 8.3 - 12.2	900			4.8 2.3 - 7.3	200	11.3 8.0 - 14.6	400	29.4 18.8 - 39.9	300
Pre-pregnancy insurance										
Medi-Cal	14.8 12.2 - 17.5	1,300	9.0 * 2.1 - 15.8	<100	6.4 3.5 - 9.4	200	20.1 14.5 - 25.7	700	30.9 20.2 - 41.7	300
Uninsured	12.6 9.4 - 15.9	1,100			9.0 4.9 - 13.2	300	15.4 9.0 - 21.8	500	22.6 11.5 - 33.7	200
Private	69.8 66.2 - 73.4	6,100	87.8 * 80.1 - 95.6	700	82.3 76.9 - 87.6	2,900	60.8 53.6 - 68.1	2,000	43.0 28.9 - 57.1	400
Prenatal insurance										
Medi-Cal	31.2 27.7 - 34.8	2,700	16.9 * 6.0 - 27.7	100	17.5 12.4 - 22.6	600	37.0 29.9 - 44.1	1,200	69.5 56.9 - 82.0	700
Uninsured	1.7 * 0.3 - 3.0	100								
Private	65.0 61.4 - 68.6	5,700	83.1 * 72.3 - 94.0	600	79.4 73.8 - 84.9	2,700	57.2 49.8 - 64.6	1,900	28.0 15.6 - 40.4	300
Maternal postpartum insurance										
Medi-Cal	24.9 21.3 - 28.5	2,200	15.7 * 5.0 - 26.4	100	12.8 8.5 - 17.1	400	31.6 24.4 - 38.8	1,000	52.4 38.8 - 66.1	500
Uninsured	5.7 3.5 - 8.0	500			3.7 * 0.7 - 6.6	100	7.1 * 2.9 - 11.4	200	11.4 * 2.3 - 20.5	100
Private	65.9 62.1 - 69.6	5,800	84.3 * 73.6 - 95.0	600	79.9 74.1 - 85.7	2,800	57.0 49.6 - 64.4	1,900	32.5 19.8 - 45.3	300
Infant health insurance										
Medi-Cal	33.2 29.6 - 36.9	2,900	15.7 * 5.0 - 26.4	100	18.9 13.6 - 24.2	700	40.3 33.0 - 47.7	1,300	71.2 58.8 - 83.6	700
Uninsured										
Private	64.4 60.7 - 68.0	5,600	84.3 * 73.6 - 95.0	600	78.4 72.7 - 84.1	2,700	57.1 49.6 - 64.5	1,800	25.8 13.5 - 38.1	300
Maternal Demographics										
Age										
15-19	1.1 * 0.5 - 1.8	100							5.3 * 1.4 - 9.2	<100
20-34	60.6 55.7 - 65.6	5,300	70.5 56.4 - 84.7	500	50.5 42.1 - 58.9	1,800	65.1 57.4 - 72.7	2,200	66.2 52.3 - 80.0	700
35+	38.2 33.3 - 43.2	3,400	29.5 15.3 - 43.6	200	49.4 41.0 - 57.8	1,700	33.6 26.0 - 41.3	1,100	28.6 14.5 - 42.6	300
Did not complete high school (or GED)	8.9 6.7 - 11.0	800			5.6 2.5 - 8.6	200	10.8 6.8 - 14.8	400	18.1 9.1 - 27.2	200
Unmarried	25.2 21.2 - 29.1	2,200	12.2 * 1.8 - 22.7	<100	20.7 14.1 - 27.3	700	23.6 17.8 - 29.4	800	50.6 37.1 - 64.1	500
Race/Ethnicity										
White	41.7 37.0 - 46.5	3,600	54.6 38.3 - 71.0	400	46.3 37.9 - 54.6	1,600	41.8 34.1 - 49.6	1,400	21.5 * 8.6 - 34.4	200
Hispanic	18.8 15.2 - 22.3	1,600			14.3 9.5 - 19.1	500	19.3 13.7 - 24.9	600	36.4 23.1 - 49.6	400
Asian/Pacific Islander	33.8 28.9 - 38.7	3,000	31.2 16.1 - 46.3	200	38.5 30.1 - 47.0	1,300	32.4 24.8 - 40.0	1,100	22.7 11.2 - 34.1	200
Black	5.4 4.7 - 6.2	500	4.3 * 0.2 - 8.4	<100	0.6 * 0.1 - 1.1	<100	6.2 4.2 - 8.2	200	19.5 12.8 - 26.1	200
Born outside the U.S.	45.5 40.7 - 50.4	4,000	41.7 25.7 - 57.7	300	44.5 36.1 - 52.9	1,600	42.9 35.3 - 50.5	1,400	64.4 53.0 - 75.7	700
Speaks non-English language at home	31.0 26.7 - 35.3	2,700	13.1 * 1.4 - 24.7	<100	27.3 20.2 - 34.3	900	32.5 25.2 - 39.7	1,000	52.3 39.2 - 65.4	500
Income as a percent of the Federal Poverty Guideline										
0-100%	20.1 16.9 - 23.3	1,700	10.5 * 0.5 - 20.5	<100	8.6 5.4 - 11.8	300	25.1 18.3 - 31.9	800	49.9 35.8 - 63.9	500
101-200%	12.1 9.1 - 15.1	1,000			7.1 3.3 - 10.9	200	15.3 9.5 - 21.1	500	22.0 10.5 - 33.4	200
> 200%	67.8 64.1 - 71.5	5,600	81.4 * 68.6 - 94.3	600	84.3 79.5 - 89.1	2,800	59.6 52.0 - 67.2	1,800	28.2 13.4 - 43.0	300

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of San Francisco County was 594. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. Neighborhood poverty is the percentage of residents with incomes below poverty in a woman's census tract. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

^{*} Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



MIHA Data Snapshot, San Francisco County by Prenatal Health Insurance, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	S	an Francisco Cour	nty		Medi-Cal		Private		
	%	95% CI	N	%	95% CI	N	%	95% CI	N
Total	100		8,800	31.2	27.7 - 34.8	2,700	65.0	61.4 - 68.6	5,700
Health Status before Pregnancy									
In good to excellent health	95.3	93.6 - 97.0	8,400	89.8	85.6 - 94.0	2,500	97.9 *	96.2 - 99.6	5,500
Chronic conditions									
Diabetes†	2.4	1.2 - 3.6	200	4.5	2.0 - 6.9	100	1.6 *	0.2 - 3.1	<100
Hypertension†	2.0	0.9 - 3.1	200	4.3 *	1.5 - 7.1	100			
Asthma†	6.9	4.7 - 9.2	600	8.4	4.7 - 12.2	200	6.7	3.7 - 9.6	400
Nutrition and Weight									
Daily folic acid use, month before pregnancy	43.6	38.6 - 48.6	3,800	28.7	21.1 - 36.3	800	51.7	45.1 - 58.2	2,900
Overweight before pregnancy	15.4	11.7 - 19.1	1,300	28.1	19.8 - 36.4	700	9.4	5.4 - 13.4	500
Obese before pregnancy	9.1	6.5 - 11.7	800	15.7	10.2 - 21.2	400	6.8	3.8 - 9.9	400
Inadequate weight gain during pregnancy	16.2	12.2 - 20.2	1,200	19.4	12.4 - 26.3	400	14.0	9.2 - 18.8	700
Excessive weight gain during pregnancy	37.5	32.3 - 42.7	2,900	38.3	29.7 - 46.9	800	38.7	32.0 - 45.4	2,000
Food insecurity during pregnancy	9.2	6.9 - 11.5	800	23.3	17.2 - 29.4	600	1.4 *	0.2 - 2.6	<100
Intimate Partner Violence (IPV) and Depressive Symptoms									
Physical or psychological IPV during pregnancy	5.4	3.4 - 7.4	500	13.9	8.5 - 19.2	400			
Prenatal depressive symptoms	14.4	10.8 - 17.9	1,300	24.1	17.0 - 31.1	700	8.9	5.0 - 12.9	500
Postpartum depressive symptoms	12.0	8.6 - 15.3	1,000	17.4	11.1 - 23.6	500	8.4	4.5 - 12.4	500
Hardships and Support during Pregnancy									
Experienced two or more hardships during childhood	17.1	13.5 - 20.6	1,500	30.6	23.2 - 38.0	800	10.9	6.8 - 15.0	600
Homeless or did not have a regular place to sleep	2.8	1.6 - 4.0	200	8.0	4.5 - 11.6	200			
Moved due to problems paying rent or mortgage	3.1	1.7 - 4.5	300	7.0	4.0 - 10.0	200			
Woman or partner lost job	11.7	8.2 - 15.2	1,000	23.7	15.8 - 31.7	600	6.3	3.0 - 9.7	400
Woman or partner had pay or hours cut back	8.2	5.2 - 11.1	700	16.6	9.5 - 23.7	500	4.2 *	1.5 - 6.8	200
Became separated or divorced	3.1	1.9 - 4.3	300	7.4	4.2 - 10.6	200			
Had no practical or emotional support	2.3	1.0 - 3.6	200	4.4 *	1.6 - 7.2	100			
Substance Use									
Any smoking, 3 months before pregnancy	8.9	6.3 - 11.5	800	11.5	7.4 - 15.6	300	7.3	4.1 - 10.5	400
Any smoking, 3rd trimester†	1.9 *	0.7 - 3.1	200	3.3	1.4 - 5.2	<100			
Any smoking, postpartum	3.6	2.0 - 5.1	300	7.5	4.1 - 10.9	200			
Any binge drinking, 3 months before pregnancy	18.1	14.1 - 22.0	1,600	9.8	5.4 - 14.2	300	22.6	17.0 - 28.2	1,300
Any alcohol use, 3rd trimester†	28.1	23.4 - 32.9	2,400	11.3 *	4.4 - 18.2	300	37.6	31.0 - 44.1	2,100
Pregnancy Intention and Family Planning									
Mistimed or unwanted pregnancy	18.1	14.5 - 21.7	1,600	30.5	23.6 - 37.5	800	12.0	7.8 - 16.2	700
Unsure of pregnancy intentions	11.2	8.1 - 14.3	1,000	19.7	13.7 - 25.6	500	7.2	3.5 - 10.9	400
Postpartum birth control use†	74.2	69.6 - 78.7	6,400	81.0	73.9 - 88.1	2,200	70.7	64.7 - 76.8	4,000
Infant Sleep and Breastfeeding									
Placed infant on back to sleep	89.5	86.6 - 92.4	7,800	85.3	80.6 - 90.1	2,200	92.4	88.7 - 96.1	5,300
Infant always or often shared bed	34.5	29.7 - 39.3	3,000	45.0	37.0 - 53.0	1,200	28.3	22.1 - 34.5	1,600
Intended to breastfeed, before birth	93.7	91.4 - 96.0	8,200	89.9	85.4 - 94.3	2,400	95.1	92.3 - 97.9	5,400
Intended to breastfeed exclusively, before birth	69.1	64.6 - 73.7	6,000	51.1	42.9 - 59.3	1,400	77.6	72.2 - 83.1	4,400
Breastfeeding practices									
Any breastfeeding, 1 month after delivery	92.8	90.5 - 95.0	8,000	86.1	81.3 - 90.9	2,300	96.2 *	93.7 - 98.8	5,400
Exclusive breastfeeding, 1 month after delivery	45.1	40.0 - 50.3	3,800	38.9	30.7 - 47.1	1,000	48.9	42.3 - 55.6	2,700
Any breastfeeding, 3 months after delivery	82.4	78.2 - 86.5	5,500	69.1	61.1 - 77.2	1,400	90.2	85.6 - 94.7	3,900
Exclusive breastfeeding, 3 months after delivery	35.6	30.0 - 41.1	2,300	18.7	12.3 - 25.2	400	44.9	37.3 - 52.5	1,900



MIHA Data Snapshot, San Francisco County by Prenatal Health Insurance, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	San Francisco County				Medi-Cal		Private			
	%	95% CI	N	%	95% CI	N	%	95% CI	N	
Service Utilization and Health Insurance Coverage										
Had a usual source of pre-pregnancy care	75.4	70.9 - 79.8	6,600	66.4	59.1 - 73.7	1,800	81.1	75.5 - 86.7	4,600	
Initiated prenatal care in 1st trimester	94.5	92.6 - 96.4	8,200	85.5	80.3 - 90.7	2,300				
Mom or infant needed but couldn't afford care postpartum	7.0	4.8 - 9.1	600	14.3	9.2 - 19.3	400	3.6 *	1.3 - 5.8	200	
Participated in WIC during pregnancy†	31.3	27.6 - 35.1	2,700	83.6	78.4 - 88.9	2,300	5.4	2.7 - 8.2	300	
Received CalFresh (food stamps) during pregnancy	10.3	8.3 - 12.2	900	30.6	24.2 - 37.0	800				
Pre-pregnancy insurance										
Medi-Cal	14.8	12.2 - 17.5	1,300	47.0	39.0 - 54.9	1,300				
Uninsured	12.6	9.4 - 15.9	1,100	30.8	23.0 - 38.7	800				
Private	69.8	66.2 - 73.4	6,100	16.9	9.9 - 23.9	500				
Maternal postpartum insurance										
Medi-Cal	24.9	21.3 - 28.5	2,200	74.9	67.9 - 81.9	2,000				
Uninsured	5.7	3.5 - 8.0	500	13.5	7.9 - 19.1	400				
Private	65.9	62.1 - 69.6	5,800	5.4 *	1.5 - 9.3	100				
Infant health insurance										
Medi-Cal	33.2	29.6 - 36.9	2,900	96.9 *	95.0 - 98.8	2,600	2.4 *	0.7 - 4.2	100	
Uninsured										
Private	64.4	60.7 - 68.0	5,600				96.8 *	94.8 - 98.8	5,500	
Maternal Demographics										
Age										
15-19	1.1 *	0.5 - 1.8	100	3.6 *	1.4 - 5.8	<100				
20-34	60.6	55.7 - 65.6	5,300	76.6	69.4 - 83.8	2,100	50.3	43.8 - 56.9	2,900	
35+	38.2	33.3 - 43.2	3,400	19.8	12.7 - 26.8	500	49.7	43.1 - 56.2	2,800	
Did not complete high school (or GED)	8.9	6.7 - 11.0	800	24.5	18.1 - 30.9	600				
Unmarried	25.2	21.2 - 29.1	2,200	53.3	45.0 - 61.5	1,400	12.5	8.0 - 17.1	700	
Race/Ethnicity										
White	41.7	37.0 - 46.5	3,600	13.7	6.7 - 20.7	400	55.7	49.1 - 62.3	3,100	
Hispanic	18.8	15.2 - 22.3	1,600	37.2	29.7 - 44.7	1,000	9.3	5.4 - 13.3	500	
Asian/Pacific Islander	33.8	28.9 - 38.7	3,000	34.6	26.8 - 42.5	900	33.3	26.9 - 39.7	1,900	
Black	5.4	4.7 - 6.2	500	13.9	11.0 - 16.8	400	1.6	1.0 - 2.2	<100	
Born outside the U.S.	45.5	40.7 - 50.4	4,000	64.5	57.2 - 71.9	1,800	34.0	27.6 - 40.4	1,900	
Speaks non-English language at home	31.0	26.7 - 35.3	2,700	60.7	53.1 - 68.3	1,600	14.1	9.5 - 18.7	800	
Lives in a high poverty neighborhood	12.0	9.0 - 15.0	1,000	26.9	19.5 - 34.3	700	5.2	2.5 - 7.9	300	
Income as a percent of the Federal Poverty Guideline										
0-100%	20.1	16.9 - 23.3	1,700	59.0	50.5 - 67.4	1,500	1.5 *	0.4 - 2.6	<100	
101-200%	12.1	9.1 - 15.1	1,000	30.0	22.2 - 37.8	700	2.9 *	1.0 - 4.8	200	
> 200%	67.8	64.1 - 71.5	5,600	11.0 *	4.4 - 17.7	300	95.6	93.4 - 97.8	5,300	

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of San Francisco County was 594. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

^{*} Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%)

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



MIHA Data Snapshot, San Francisco County by Race/Ethnicity, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	San F	rancisco Co	unty		Hispanic			Black			White		Asia	n/Pacific Isla	nder
	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N
Total	100		8,800	18.8	15.2 - 22.3	1,600	5.4	4.7 - 6.2	500	41.7	37.0 - 46.5	3,600	33.8	28.9 - 38.7	3,000
Health Status before Pregnancy															
In good to excellent health	95.3	93.6 - 97.0	8,400	88.0	81.8 - 94.1	1,400	89.4 *	82.2 - 96.5	400				95.3	* 91.8 - 98.7	2,800
Chronic conditions															
Diabetes†	2.4	1.2 - 3.6	200	5.4	* 1.6 - 9.1	<100	9.2 *	* 1.4 - 16.9	<100						
Hypertension†	2.0	0.9 - 3.1	200					3.1 - 21.0	<100						
Asthma†	6.9	4.7 - 9.2	600	7.2	* 2.0 - 12.4	100	20.3	11.6 - 29.0	<100	8.6	4.2 - 12.9	300	2.7	* 0.4 - 4.9	<100
Nutrition and Weight															
Daily folic acid use, month before pregnancy	43.6	38.6 - 48.6	3,800	25.2	16.7 - 33.7	400	31.4	21.8 - 41.0	100	53.4	45.2 - 61.6	1,900	42.9	33.9 - 51.8	1,300
Overweight before pregnancy	15.4	11.7 - 19.1	1,300	30.3	18.8 - 41.7	400	26.0	15.4 - 36.7	100	9.3	5.0 - 13.7	300	14.1	6.8 - 21.5	400
Obese before pregnancy	9.1	6.5 - 11.7	800	22.2	12.6 - 31.8	300	24.4	15.9 - 32.8	100	5.9	* 2.1 - 9.8	200	4.5	* 1.4 - 7.5	100
Inadequate weight gain during pregnancy	16.2	12.2 - 20.2	1,200	21.7	10.9 - 32.4	300	13.1	5.8 - 20.5	<100	16.1	9.7 - 22.4	500	14.9	8.5 - 21.3	400
Excessive weight gain during pregnancy	37.5	32.3 - 42.7	2,900	29.5	18.1 - 40.9	400	52.6	39.9 - 65.3	200	42.3	33.8 - 50.7	1,400	32.6	23.7 - 41.5	900
Food insecurity during pregnancy	9.2	6.9 - 11.5	800	26.5	18.0 - 34.9	400	19.5	11.3 - 27.7	<100				6.6	* 2.4 - 10.9	200
Intimate Partner Violence (IPV) and Depressive Symptoms															
Physical or psychological IPV during pregnancy	5.4	3.4 - 7.4	500	9.1	4.0 - 14.1	100	11.1 *	* 3.8 - 18.5	<100	2.7	* 0.3 - 5.0	<100	5.7	* 1.6 - 9.9	200
Prenatal depressive symptoms	14.4	10.8 - 17.9	1,300	26.7	16.6 - 36.7	400	21.4	14.0 - 28.9	<100	9.5	4.5 - 14.5	300	11.8	5.7 - 17.9	300
Postpartum depressive symptoms	12.0	8.6 - 15.3	1,000	18.7	8.7 - 28.7	300	16.0	9.3 - 22.8	<100	6.8	* 2.6 - 11.0	200	14.4	8.2 - 20.6	400
Hardships and Support during Pregnancy															
Experienced two or more hardships during childhood	17.1	13.5 - 20.6	1,500	35.5	25.0 - 46.0	600	39.8	28.5 - 51.2	200	11.7	6.5 - 17.0	400	9.8	4.4 - 15.1	300
Homeless or did not have a regular place to sleep	2.8	1.6 - 4.0	200				23.0	12.3 - 33.6	100				1.4	* 0.2 - 2.6	<100
Moved due to problems paying rent or mortgage	3.1	1.7 - 4.5	300	3.9	* 0.6 - 7.2	<100	13.5	5.8 - 21.2	<100				3.5	* 0.7 - 6.2	100
Woman or partner lost job	11.7	8.2 - 15.2	1,000	20.4	11.1 - 29.7	300	18.1	9.8 - 26.3	<100	11.0	4.8 - 17.2	400	6.8	* 2.6 - 11.1	200
Woman or partner had pay or hours cut back	8.2	5.2 - 11.1	700	12.2	5.5 - 18.9	200	15.0	7.0 - 23.1	<100	7.4	* 1.9 - 12.9	300	5.6	* 1.6 - 9.7	200
Became separated or divorced	3.1	1.9 - 4.3	300	8.8	3.7 - 13.9	100	18.0	9.5 - 26.5	<100						
Had no practical or emotional support	2.3	1.0 - 3.6	200	5.2	1.1 - 9.3	<100	4.4 *	1.3 - 7.6	<100						
Substance Use															
Any smoking, 3 months before pregnancy	8.9	6.3 - 11.5	800	6.8	* 2.0 - 11.5	100	27.3	17.0 - 37.6	100	8.8	4.1 - 13.6	300	7.3	3.3 - 11.3	200
Any smoking, 3rd trimester†	1.9 *	0.7 - 3.1	200				13.3	6.1 - 20.5	<100						
Any smoking, postpartum	3.6	2.0 - 5.1	300				25.5	15.3 - 35.8	100				2.9	* 0.7 - 5.1	<100
Any binge drinking, 3 months before pregnancy	_	14.1 - 22.0	1,600	11.6		200	16.0	7.6 - 24.3	<100		18.4 - 32.2	900	13.9		400
Any alcohol use, 3rd trimester†	28.1	23.4 - 32.9	2,400	16.7	* 6.5 - 27.0	300	8.1 *	1.6 - 14.6	<100	47.2	39.0 - 55.4	1,700	14.7	7.2 - 22.2	400
Pregnancy Intention and Family Planning															
Mistimed or unwanted pregnancy	18.1	14.5 - 21.7	1,600	27.6	18.9 - 36.4	500	24.2	16.1 - 32.3	100	12.4	7.4 - 17.4	500	19.3	12.0 - 26.6	600
Unsure of pregnancy intentions	11.2	8.1 - 14.3	1,000	10.5	4.8 - 16.1	200	33.1	21.6 - 44.7	200	10.1	4.5 - 15.7	400	9.8	4.8 - 14.7	300
Postpartum birth control use†	74.2	69.6 - 78.7	6,400	86.5	78.6 - 94.4	1,400	75.6	66.5 - 84.6	300	75.3	68.0 - 82.6	2,700	65.9	57.3 - 74.5	1,900
Infant Sleep and Breastfeeding															
Placed infant on back to sleep	89.5	86.6 - 92.4	7,800	83.2	75.4 - 91.0	1,300	82.3	74.5 - 90.1	400	95.0	* 91.4 - 98.5	3,400	87.2	81.2 - 93.1	2,500
Infant always or often shared bed	34.5	29.7 - 39.3	3,000	30.1	20.3 - 40.0	500	52.4	41.3 - 63.4	200	23.3	16.5 - 30.0	800	49.0	39.8 - 58.2	1,400
Intended to breastfeed, before birth	93.7	91.4 - 96.0	8,200	92.9	* 88.2 - 97.6	1,500	84.4	76.5 - 92.3	400	96.4	* 93.2 - 99.7	3,500		* 89.3 - 97.2	2,700
Intended to breastfeed exclusively, before birth	69.1	64.6 - 73.7	6,000	61.2	50.4 - 71.9	1,000	57.0	45.5 - 68.5	300	85.9	80.2 - 91.7	3,100	55.1	46.1 - 64.1	1,600
Breastfeeding practices															
Any breastfeeding, 1 month after delivery		90.5 - 95.0	8,000		83.4 - 95.2	1,400		60.2 - 81.5	300					89.8 - 97.2	2,700
Exclusive breastfeeding, 1 month after delivery	_	40.0 - 50.3	3,800		23.7 - 44.3	500		34.3 - 55.7	200		53.0 - 69.2	2,200		22.4 - 39.6	900
Any breastfeeding, 3 months after delivery	_	78.2 - 86.5	5,500		66.4 - 85.2	1,100		37.5 - 62.9	200		* 86.7 - 97.0	2,500		71.7 - 88.8	1,600
Exclusive breastfeeding, 3 months after delivery	35.6	30.0 - 41.1	2,300	21.8	11.6 - 31.9	300	32.3	21.4 - 43.1	100	49.6	40.2 - 58.9	1,400	25.9	15.9 - 35.9	500



MIHA Data Snapshot, San Francisco County by Race/Ethnicity, 2013-2015

Maternal and Infant Health Assessment (MIHA) Survey

	San F	rancisco Co	unty	<u> </u>			Black			White		Asia	n/Pacific Isla	nder	
	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N
Service Utilization and Health Insurance Coverage															
Had a usual source of pre-pregnancy care	75.4	70.9 - 79.8	6,600	72.8	63.3 - 82.2	1,200	81.4	73.1 - 89.7	400	77.3	70.2 - 84.4	2,800	74.0	66.1 - 81.9	2,200
Initiated prenatal care in 1st trimester	94.5	92.6 - 96.4	8,200	90.5	85.0 - 96.1	1,400	84.6	* 75.2 - 94.0	400				93.8	* 90.1 - 97.6	2,800
Mom or infant needed but couldn't afford care postpartum	7.0	4.8 - 9.1	600	8.1 '	* 3.3 - 12.8	100	13.1	6.4 - 19.8	<100	4.3 *	0.9 - 7.6	200	9.0	4.8 - 13.3	300
Participated in WIC during pregnancy†	31.3	27.6 - 35.1	2,700	68.0	57.3 - 78.7	1,100	70.6	60.9 - 80.2	300	7.0 *	1.9 - 12.1	200	33.9	25.9 - 41.9	1,000
Received CalFresh (food stamps) during pregnancy	10.3	8.3 - 12.2	900	23.5	15.5 - 31.4	400	60.9	50.1 - 71.6	300	2.7 *	0.7 - 4.8	<100	3.6	1.5 - 5.8	100
Pre-pregnancy insurance															
Medi-Cal	14.8	12.2 - 17.5	1,300	31.2	22.2 - 40.2	500	48.7	38.2 - 59.3	200	2.2 *	0.2 - 4.2	<100	15.8	10.0 - 21.6	500
Uninsured	12.6	9.4 - 15.9	1,100	23.7	15.7 - 31.6	400	16.2	7.1 - 25.4	<100	8.0 *	2.5 - 13.5	300	10.7	5.5 - 15.8	300
Private	69.8	66.2 - 73.4	6,100	38.8	27.4 - 50.2	600	31.2	21.5 - 41.0	100	88.0	82.0 - 94.0	3,200	71.6	64.3 - 79.0	2,100
Prenatal insurance															
Medi-Cal	31.2	27.7 - 34.8	2,700	62.1	51.3 - 73.0	1,000	79.9	72.6 - 87.2	400	10.3	4.8 - 15.7	400	32.4	24.8 - 40.0	900
Uninsured	1.7 *	0.3 - 3.0	100												
Private	65.0	61.4 - 68.6	5,700	32.2	21.3 - 43.0	500	18.5	11.6 - 25.5	<100	86.2	80.1 - 92.3	3,100	64.3	56.4 - 72.1	1,900
Maternal postpartum insurance															
Medi-Cal	24.9	21.3 - 28.5	2,200	47.6	36.8 - 58.5	800	78.7	71.1 - 86.4	400	9.1 *	3.6 - 14.7	300	23.2	16.5 - 29.8	700
Uninsured	5.7	3.5 - 8.0	500	10.0	4.9 - 15.1	200							8.8	* 3.3 - 14.2	300
Private	65.9	62.1 - 69.6	5,800	31.1	20.3 - 41.9	500	19.2	12.0 - 26.4	<100	86.1	79.7 - 92.4	3,100	67.4	59.6 - 75.2	2,000
Infant health insurance															
Medi-Cal	33.2	29.6 - 36.9	2,900	70.0	59.3 - 80.8	1,100	81.6	74.4 - 88.9	400	11.5	5.7 - 17.4	400	32.7	25.0 - 40.4	900
Uninsured															
Private	64.4	60.7 - 68.0	5,600	27.7	16.9 - 38.4	500	15.0	8.2 - 21.8	<100	86.3	80.2 - 92.4	3,100	64.7	56.7 - 72.6	1,900
Maternal Demographics															
Age															
15-19	1.1 *	0.5 - 1.8	100				11.3	* 3.8 - 18.8	<100						
20-34	60.6	55.7 - 65.6	5,300	71.3	60.6 - 82.0	1,200	70.6	60.3 - 80.9	300	53.4	45.2 - 61.7	1,900	63.1	54.2 - 71.9	1,900
35+	38.2	33.3 - 43.2	3,400	26.4	15.7 - 37.0	400	18.1	9.4 - 26.9	<100	46.6	38.3 - 54.8	1,700	36.6	27.8 - 45.4	1,100
Did not complete high school (or GED)	8.9	6.7 - 11.0	800	31.7	22.4 - 41.0	500	19.9	10.4 - 29.4	<100				5.3	* 1.9 - 8.8	200
Unmarried	25.2	21.2 - 29.1	2,200	53.6	42.7 - 64.5	900	77.1	68.4 - 85.8	400	14.4	8.3 - 20.5	500	14.6	8.6 - 20.5	400
Born outside the U.S.	45.5	40.7 - 50.4	4,000	63.6	52.5 - 74.7	1,000	11.5	* 4.1 - 18.8	<100	23.0	15.5 - 30.5	800	70.3	61.9 - 78.8	2,100
Speaks non-English language at home	31.0	26.7 - 35.3	2,700	56.7	45.8 - 67.6	900				10.2	4.9 - 15.4	400	49.1	40.0 - 58.3	1,400
Lives in a high poverty neighborhood	12.0	9.0 - 15.0	1,000	23.6	14.0 - 33.3	400	43.7	32.5 - 54.8	200	6.1 *	1.9 - 10.3	200	8.1	3.6 - 12.6	200
Income as a percent of the Federal Poverty Guideline															
0-100%	20.1	16.9 - 23.3	1,700	50.0	38.2 - 61.7	700	66.4	56.3 - 76.5	300	5.9 *	0.6 - 11.1	200	15.7	10.4 - 21.0	400
101-200%	12.1	9.1 - 15.1	1,000	15.5	7.7 - 23.3	200	18.3	9.6 - 27.1	<100	5.0 *	* 1.9 - 8.0	200	18.5	11.5 - 25.4	500
> 200%	67.8	64.1 - 71.5	5,600	34.6	21.7 - 47.4	500	15.3	7.2 - 23.4	<100	89.2	83.3 - 95.0	3,200	65.8	58.0 - 73.7	1,900

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of San Francisco County was 594. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

^{*} Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



MIHA Data Snapshot, San Francisco County by Total Live Births, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	S	San Francisco County			First Live Birth				Second Live Birth or More		
	%	95% CI	N	%	95% (CI	N	%	95% CI	N	
Total	100		8,800	55.7	50.8 -	60.7	4,900	44.3	39.3 - 49.2	3,900	
Health Status before Pregnancy											
In good to excellent health Chronic conditions	95.3	93.6 - 97.0	8,400	96.5 *	94.3 -	98.6	4,700	93.8	90.9 - 96.6	3,700	
Diabetes†	2.4	1.2 - 3.6	200	1.5 *	0.3 -	2.7	<100	3.6 *	1.4 - 5.8	100	
Hypertension [†]	2.0	0.9 - 3.1	200	1.9 *	0.5 -	3.4	<100	2.1 *	0.5 - 3.7	<100	
Asthma†	6.9	4.7 - 9.2	600	6.6	3.8 -	9.3	300	7.4	3.7 - 11.1	300	
Nutrition and Weight											
Daily folic acid use, month before pregnancy	43.6	38.6 - 48.6	3,800	50.6	43.8 -	57.5	2,500	34.9	27.9 - 41.9	1,400	
Overweight before pregnancy	15.4	11.7 - 19.1	1,300	11.2	6.8 -	15.6	500	20.9	14.5 - 27.3	800	
Obese before pregnancy	9.1	6.5 - 11.7	800	6.5	3.4 -	9.6	300	12.6	8.1 - 17.0	500	
Inadequate weight gain during pregnancy	16.2	12.2 - 20.2	1,200	16.4	11.0 -	21.9	700	15.9	10.3 - 21.6	500	
Excessive weight gain during pregnancy	37.5	32.3 - 42.7	2,900	38.5	31.5 -	45.5	1,700	36.2	28.3 - 44.1	1,200	
Food insecurity during pregnancy	9.2	6.9 - 11.5	800	7.1	4.1 -	10.0	300	11.9	8.0 - 15.8	500	
Intimate Partner Violence (IPV) and Depressive Symptoms											
Physical or psychological IPV during pregnancy	5.4	3.4 - 7.4	500	5.3	2.6 -	8.0	300	5.5	2.5 - 8.5	200	
Prenatal depressive symptoms	14.4	10.8 - 17.9	1,300	12.0	7.6 -	16.4	600	17.3	11.6 - 23.1	700	
Postpartum depressive symptoms	12.0	8.6 - 15.3	1,000	11.9	7.5 -	16.3	600	12.1	7.0 - 17.3	500	
Hardships and Support during Pregnancy											
Experienced two or more hardships during childhood	17.1	13.5 - 20.6	1,500	15.5	10.8 -	20.2	800	19.1	13.5 - 24.6	700	
Homeless or did not have a regular place to sleep	2.8	1.6 - 4.0	200	3.1	1.4 -	4.8	200	2.4 *	0.8 - 4.1	<100	
Moved due to problems paying rent or mortgage	3.1	1.7 - 4.5	300	3.6 *	1.5 -	5.7	200	2.5 *	0.9 - 4.1	<100	
Woman or partner lost job	11.7	8.2 - 15.2	1,000	12.8	7.7 -	17.9	600	10.4	5.8 - 14.9	400	
Woman or partner had pay or hours cut back	8.2	5.2 - 11.1	700	8.9	4.3 -	13.4	400	7.3	3.8 - 10.8	300	
Became separated or divorced	3.1	1.9 - 4.3	300	2.0	0.9 -	3.1	<100	4.4	2.1 - 6.8	200	
Had no practical or emotional support	2.3	1.0 - 3.6	200	2.8 *	0.7 -	4.9	100	1.6 *	0.4 - 2.9	<100	
Substance Use											
Any smoking, 3 months before pregnancy	8.9	6.3 - 11.5	800	10.2	6.3 -	14.1	500	7.3	4.0 - 10.5	300	
Any smoking, 3rd trimester†	1.9 *	0.7 - 3.1	200	1.3 *	0.1 -	2.5	<100	2.7 *	0.4 - 4.9	100	
Any smoking, postpartum	3.6	2.0 - 5.1	300	2.8 *	1.0 -	4.7	100	4.5	1.9 - 7.1	200	
Any binge drinking, 3 months before pregnancy	18.1	14.1 - 22.0	1,600	24.8	18.7 -	30.9	1,200	9.6	5.5 - 13.8	400	
Any alcohol use, 3rd trimester†	28.1	23.4 - 32.9	2,400	30.1	23.4 -	36.7	1,500	25.7	18.8 - 32.6	1,000	
Pregnancy Intention and Family Planning											
Mistimed or unwanted pregnancy	18.1	14.5 - 21.7	1,600	18.8	13.8 -	23.9	900	17.2	12.2 - 22.2	700	
Unsure of pregnancy intentions	11.2	8.1 - 14.3	1,000	10.4	6.2 -	14.6	500	12.2	7.5 - 16.8	500	
Postpartum birth control use†	74.2	69.6 - 78.7	6,400	70.4	63.9 -	76.8	3,400	78.9	72.7 - 85.1	3,000	
Infant Sleep and Breastfeeding											
Placed infant on back to sleep	89.5	86.6 - 92.4	7,800	91.8	88.3 -	95.3	4,400	86.6	81.7 - 91.5	3,300	
Infant always or often shared bed	34.5	29.7 - 39.3	3,000	34.5	28.0 -	41.1	1,700	34.5	27.3 - 41.6	1,300	
Intended to breastfeed, before birth	93.7	91.4 - 96.0	8,200	95.0	92.1 -	97.9	4,600	92.1	88.5 - 95.7	3,600	
Intended to breastfeed exclusively, before birth	69.1	64.6 - 73.7	6,000	75.2	69.3 -	81.0	3,700	61.6	54.4 - 68.7	2,400	
Breastfeeding practices											
Any breastfeeding, 1 month after delivery	92.8	90.5 - 95.0	8,000	94.8	92.0 -	97.5	4,500	90.3	86.5 - 94.0	3,400	
Exclusive breastfeeding, 1 month after delivery	45.1	40.0 - 50.3	3,800	42.8	35.8 -	49.7	2,000	48.1	40.5 - 55.7	1,800	
Any breastfeeding, 3 months after delivery	82.4	78.2 - 86.5	5,500	86.6	81.5 -	91.6	3,200	77.2	70.3 - 84.0	2,300	
Exclusive breastfeeding, 3 months after delivery	35.6	30.0 - 41.1	2,300	36.1	28.6 -	43.6	1,300	34.9	26.6 - 43.2	1,000	



MIHA Data Snapshot, San Francisco County by Total Live Births, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	S	San Francisco County			First Live Birth		Second Live Birth or More		
	%	95% CI	N	%	95% CI	N	%	95% CI	N
Service Utilization and Health Insurance Coverage									
Had a usual source of pre-pregnancy care	75.4	70.9 - 79.8	6,600	73.5	67.6 - 79.4	3,600	77.7	71.1 - 84.3	3,000
Initiated prenatal care in 1st trimester	94.5	92.6 - 96.4	8,200	95.5	93.3 - 97.8	4,700	93.1	89.8 - 96.3	3,500
Mom or infant needed but couldn't afford care postpartum	7.0	4.8 - 9.1	600	8.0	4.7 - 11.3	400	5.7	3.1 - 8.3	200
Participated in WIC during pregnancy†	31.3	27.6 - 35.1	2,700	22.4	17.5 - 27.4	1,100	42.8	35.9 - 49.6	1,600
Received CalFresh (food stamps) during pregnancy	10.3	8.3 - 12.2	900	5.2	3.2 - 7.1	300	16.8	12.8 - 20.7	600
Pre-pregnancy insurance									
Medi-Cal	14.8	12.2 - 17.5	1,300	7.5	4.9 - 10.1	400	24.0	18.7 - 29.3	900
Uninsured	12.6	9.4 - 15.9	1,100	13.7	8.9 - 18.5	700	11.3	6.9 - 15.6	400
Private	69.8	66.2 - 73.4	6,100	76.9	71.6 - 82.1	3,800	60.9	54.5 - 67.2	2,400
Prenatal insurance									
Medi-Cal	31.2	27.7 - 34.8	2,700	23.0	18.0 - 28.1	1,100	41.5	35.1 - 47.9	1,600
Uninsured	1.7 *	0.3 - 3.0	100	2.2 *	0.1 - 4.3	100			
Private	65.0	61.4 - 68.6	5,700	71.8	66.3 - 77.2	3,500	56.6	50.2 - 63.0	2,200
Maternal postpartum insurance									
Medi-Cal	24.9	21.3 - 28.5	2,200	20.7	15.6 - 25.8	1,000	30.2	24.2 - 36.1	1,200
Uninsured	5.7	3.5 - 8.0	500	4.6 *	1.9 - 7.4	200	7.1	3.3 - 10.9	300
Private	65.9	62.1 - 69.6	5,800	71.8	66.2 - 77.4	3,500	58.4	52.0 - 64.9	2,300
Infant health insurance									
Medi-Cal	33.2	29.6 - 36.9	2,900	25.7	20.3 - 31.1	1,200	42.8	36.4 - 49.2	1,600
Uninsured									
Private	64.4	60.7 - 68.0	5,600	70.9	65.3 - 76.6	3,400	56.1	49.7 - 62.5	2,200
Maternal Demographics									
Age									
15-19	1.1 *	0.5 - 1.8	100	2.0 *	0.8 - 3.2	<100			
20-34	60.6	55.7 - 65.6	5,300	64.4	57.9 - 71.0	3,200	55.9	48.4 - 63.3	2,200
35+	38.2	33.3 - 43.2	3,400	33.6	27.1 - 40.1	1,600	44.0	36.6 - 51.5	1,700
Did not complete high school (or GED)	8.9	6.7 - 11.0	800	5.2	3.0 - 7.5	300	13.6	9.4 - 17.8	500
Unmarried	25.2	21.2 - 29.1	2,200	24.7	19.1 - 30.2	1,200	25.9	20.0 - 31.7	1,000
Race/Ethnicity									
White	41.7	37.0 - 46.5	3,600	46.5	39.8 - 53.3	2,300	35.6	28.5 - 42.7	1,400
Hispanic	18.8	15.2 - 22.3	1,600	12.1	8.4 - 15.8	600	27.3	20.8 - 33.9	1,000
Asian/Pacific Islander	33.8	28.9 - 38.7	3,000	36.7	30.0 - 43.5	1,800	30.1	23.1 - 37.2	1,200
Black	5.4	4.7 - 6.2	500	4.4	3.1 - 5.8	200	6.7	5.1 - 8.3	300
Born outside the U.S.	45.5	40.7 - 50.4	4,000	40.1	33.4 - 46.8	2,000	52.4	45.2 - 59.6	2,000
Speaks non-English language at home	31.0	26.7 - 35.3	2,700	23.7	18.2 - 29.3	1,100	40.3	33.3 - 47.2	1,500
Lives in a high poverty neighborhood	12.0	9.0 - 15.0	1,000	9.6	6.0 - 13.2	500	15.1	9.9 - 20.2	600
Income as a percent of the Federal Poverty Guideline									
0-100%	20.1	16.9 - 23.3	1,700	16.9	12.1 - 21.8	800	24.2	19.2 - 29.2	900
101-200%	12.1	9.1 - 15.1	1,000	7.3	4.0 - 10.7	300	18.4	12.8 - 24.1	700
> 200%	67.8	64.1 - 71.5	5,600	75.8	70.4 - 81.2	3,600	57.4	50.7 - 64.0	2,100

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of San Francisco County was 594. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

^{*} Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%)

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).