

# MIHA Data Snapshot, San Mateo County by Education, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	San Mat	eo County	/	<	High Schoo	ı	High Sch	ool Graduat	e or GED		Some College	2	Co	llege Gradua	ate
	% 95	% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N
Total	100	;	8,800	10.0	6.9 - 13.0	900	11.5	8.0 - 15.0	1,000	21.1	16.9 - 25.4	1,800	57.4	52.3 - 62.5	5,000
Health Status before Pregnancy															
In good to excellent health	97.2 95.7	- 98.7	8,600												
Chronic conditions															
Diabetes†	3.0 1.3	- 4.7	300							7.3	* 1.6 - 13.0	100	1.8 *	* 0.1 - 3.6	<100
Hypertension†	1.4 * 0.4	- 2.3	100												
Asthma†	5.2 3.0	- 7.5	500							3.5	* 0.4 - 6.6	<100	4.9	2.2 - 7.7	200
Nutrition and Weight															
Daily folic acid use, month before pregnancy	41.8 36.8	- 46.7	3,700	20.7 *	7.8 - 33.6	200	22.9 *	8.4 - 37.4	200	41.0	29.7 - 52.3	800	49.4	43.0 - 55.9	2,500
Overweight before pregnancy	20.2 16.0	- 24.3	1,700	29.0	3.5 - 44.4	200	23.1 *	7.4 - 38.9	200	20.6	11.7 - 29.4	400	18.3	13.1 - 23.6	900
Obese before pregnancy	<b>15.3</b> 11.5	- 19.1	1,300	28.1	2.8 - 43.4	200	39.0	22.2 - 55.7	300	23.5	13.0 - 34.0	400	5.9	3.1 - 8.8	300
Inadequate weight gain during pregnancy	18.2 14.0	- 22.3	1,400	27.4 * 3	0.2 - 44.6	100	28.4 *	11.4 - 45.3	200	19.6	9.9 - 29.2	300	15.3	10.5 - 20.1	700
Excessive weight gain during pregnancy	38.4 33.1	- 43.8	2,900	48.0	27.7 - 68.4	300	46.2	26.4 - 66.1	300	42.1	29.8 - 54.4	700	35.0	28.5 - 41.4	1,600
Food insecurity during pregnancy	11.1 7.7	- 14.4	1,000	25.8	1.6 - 40.1	200	26.3	11.4 - 41.2	300	17.5	8.1 - 27.0	300	2.6 *	0.9 - 4.3	100
Intimate Partner Violence (IPV) and Depressive Symptoms															
Physical or psychological IPV during pregnancy	4.6 2.5	- 6.7	400							7.6	* 1.6 - 13.6	100	1.7 *	0.4 - 3.1	<100
Prenatal depressive symptoms	-	- 11.3	700	18.4 *	6.0 - 30.9	200	7.6 *	0.5 - 14.7	<100		* 5.0 - 19.8	200		2.5 - 8.1	300
Postpartum depressive symptoms	10.4 7.3	- 13.5	900	11.2 *	3.1 - 19.4	<100					8.2 - 25.5	300		4.4 - 11.8	400
Hardships and Support during Pregnancy															
Experienced two or more hardships during childhood	15.9 12.2	- 19.6	1,400	163	80.1 - 62.5	400	20.7*	7.7 - 33.7	200	21 0	12.2 - 31.7	400	7.0	4.1 - 9.9	300
Homeless or did not have a regular place to sleep	3.0 * 1.2		300		02.5	400	20.7	7.7 33.7	200	-	* 0.5 - 7.5	<100	7.0	4.1 3.3	300
Moved due to problems paying rent or mortgage	5.7 3.1		500							-	* 1.1 - 13.5	100	2.4 *	0.5 - 4.2	100
Woman or partner lost job	_	- 12.9	900		8.2 - 35.8	200		0.6 - 23.1	100	17.7	8.4 - 26.9	300	4.2	1.9 - 6.4	200
Woman or partner had pay or hours cut back	7.1 4.4		600		4.2 - 29.5	100	_	0.3 - 15.8	<100		* 4.9 - 21.4	200		0.7 - 4.6	100
Became separated or divorced	3.1 1.4		300		4.0 - 31.1	200		0.5 15.0	-200		* 0.3 - 5.3	<100			100
Had no practical or emotional support	5.7 3.1		500		4.8 - 31.5	200				_	* 2.4 - 16.0	200			
Substance Use	3.,			10.1						J. <b>L</b>					
Any smoking, 3 months before pregnancy	8.6 5.8	115	800	60*	1.0 - 12.9	<100	1/1 2 *	3.0 - 25.3	100	2/1.2	14.2 - 34.4	400	17*	0.4 - 3.0	<100
Any smoking, 3 months belove pregnancy  Any smoking, 3rd trimester†	1.1 * 0.3		<100	0.5	1.0 - 12.9	<100	14.2	3.0 - 23.3	100	24.3	14.2 - 34.4	400		0.4 - 3.0	<100
Any smoking, postpartum	2.7 1.2		200								* 1.8 - 11.7	100			
Any binge drinking, 3 months before pregnancy	13.6 10.1		1.200								11.5 - 30.8	400	13.5	9.1 - 17.9	700
Any alcohol use, 3rd trimester†			1,000								* 1.2 - 11.6	100		11.2 - 21.3	800
Pregnancy Intention and Family Planning	11.5 0.1	14.5	1,000							0.4	1.2 11.0	100	10.5	11.2 21.3	000
Mistimed or unwanted pregnancy	22 7 40 4	27.0	2 000	F1 2 '	)F2 672	400	25.5	100 [12	200	21.1	20.6 41.6	600	11.0	7.5 16.4	600
Unsure of pregnancy intentions	22.7 18.4 7.8 5.1		2,000 700	51.3	35.3 - 67.3	400		19.8 - 51.2 1.9 - 19.4	300 100	-	20.6 - 41.6 * 4.0 - 19.1	600 200	6.4	7.5 - 16.4 3.2 - 9.6	600 300
Postpartum birth control use†	7.6 5.1		6,800		73.1 - 94.8	700	10.6	1.9 - 19.4	100	_	74.4 - 90.6	1,500	_	68.5 - 80.3	3,600
	79.1 74.9	- 03.2	0,800	64.0	3.1 - 34.0	700				62.5	74.4 - 90.0	1,500	74.4	00.3 - 00.3	3,000
Infant Sleep and Breastfeeding				04.6*											
Placed infant on back to sleep	88.6 85.3		7,700		8.8 - 94.3	700		63.8 - 92.3	800		* 81.6 - 95.9	1,600		88.3 - 95.1	4,600
Infant always or often shared bed	30.1 25.4		2,600		31.7 - 63.8	400		19.7 - 51.5	300	_	30.3 - 53.3	700		15.9 - 26.5	1,100
Intended to breastfeed, before birth	92.9 90.1		8,100		95.2 - 95.2	700		72.4 - 98.1	800		* 81.6 - 96.7	1,600		95.9 - 99.3	4,800
Intended to breastfeed exclusively, before birth	64.7 59.8	- 69.6	5,600	40.6	24.7 - 56.4	400	53.3	36.8 - 69.8	500	48.9	37.4 - 60.4	900	//.1	71.8 - 82.4	3,800
Breastfeeding practices	00.7 00.0	02.0	7 700	040*	22.7 00.0	700	70.6*	67.3 01.0	000	02.0	74.2 04.0	1.500	04.0	024 070	4 700
Any breastfeeding, 1 month after delivery	89.7 86.6		7,700	-	73.7 - 96.0	700		67.3 - 91.9	800		74.2 - 91.9	1,500		92.1 - 97.8	4,700
Exclusive breastfeeding, 1 month after delivery	44.6 39.4		3,800		25.9 - 58.2	400		20.7 - 53.8	400		27.1 - 49.4	700		42.4 - 55.6	2,400
Any breastfeeding, 3 months after delivery	78.2 73.3		5,000		19.1 - 81.9	400		42.8 - 78.7	400		50.3 - 76.5	900		86.6 - 95.0	3,200
Exclusive breastfeeding, 3 months after delivery	32.8 27.1	- 38.5	2,100	19.3 *	5.1 - 33.6	100	22.1 *	5.0 - 39.3	200	22.9	12.1 - 33.7	300	42.0	34.3 - 49.8	1,500



#### MIHA Data Snapshot, San Mateo County by Education, 2013-2015

Maternal and Infant Health Assessment (MIHA) Survey

	San	Mateo Cou	nty		<high school<="" th=""><th></th><th>High Sch</th><th>nool Graduat</th><th>e or GED</th><th></th><th>Some College</th><th>•</th><th>Co</th><th>ollege Gradua</th><th>ate</th></high>		High Sch	nool Graduat	e or GED		Some College	•	Co	ollege Gradua	ate
	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N
Service Utilization and Health Insurance Coverage															
Had a usual source of pre-pregnancy care	77.7	73.5 - 81.9	6,800	59.2	43.7 - 74.7	500	57.5	41.4 - 73.6	600	84.1	76.2 - 91.9	1,600	82.4	77.4 - 87.4	4,100
Initiated prenatal care in 1st trimester	92.4	89.6 - 95.1	7,900	77.8	* 64.4 - 91.1	700	88.5	* 78.0 - 99.1	900	94.2	* 88.7 - 99.7	1,700	94.8	91.9 - 97.8	4,600
Mom or infant needed but couldn't afford care postpartum	5.7	3.6 - 7.8	500	6.0	* 0.5 - 11.5	<100	10.4	* 2.3 - 18.5	100	8.9	* 3.1 - 14.6	200	3.1	* 1.0 - 5.2	100
Participated in WIC during pregnancy†	30.5	25.7 - 35.4	2,600	91.0	* 82.6 - 99.3	800	74.3	59.9 - 88.6	700	43.2	31.7 - 54.7	800	4.9	2.3 - 7.5	200
Received CalFresh (food stamps) during pregnancy	9.1	6.3 - 12.0	800	30.8	16.2 - 45.4	300	26.7	11.8 - 41.7	300	11.0	5.2 - 16.9	200	1.2	* 0.2 - 2.2	<100
Pre-pregnancy insurance															
Medi-Cal	15.2	11.5 - 18.9	1,300	44.6	28.6 - 60.6	400	37.6	22.0 - 53.1	400	22.5	12.8 - 32.2	400	2.4	* 0.7 - 4.2	100
Uninsured	9.6	6.5 - 12.6	800	22.8	9.7 - 35.9	200	24.3	10.3 - 38.4	200	8.1	* 2.2 - 14.0	100	5.0	* 1.9 - 8.1	300
Private	72.0	67.4 - 76.5	6,300	23.2	* 9.3 - 37.0	200	35.0	19.3 - 50.8	400	65.8	55.0 - 76.6	1,200	90.5	86.7 - 94.3	4,500
Prenatal insurance															
Medi-Cal	26.3	21.8 - 30.8	2,300	74.0	59.3 - 88.7	600	70.5	55.6 - 85.5	700	31.7	21.2 - 42.2	600	6.7	3.5 - 10.0	300
Uninsured															
Private	71.7	67.1 - 76.3	6,300	21.0	* 7.4 - 34.6	200	25.4	11.3 - 39.6	300	67.6	57.1 - 78.2	1,200	91.6	88.0 - 95.2	4,600
Maternal postpartum insurance															
Medi-Cal	17.5	13.6 - 21.4	1,500	51.7	35.3 - 68.1	400	50.6	34.3 - 66.8	500	20.2	11.6 - 28.9	400	4.0	* 1.4 - 6.6	200
Uninsured	7.1	4.4 - 9.8	600	18.8	* 5.9 - 31.7	200	16.9	* 5.5 - 28.3	200	11.0	* 3.2 - 18.8	200	1.9	* 0.2 - 3.5	<100
Private	72.4	67.8 - 77.0	6,300	17.4	* 4.3 - 30.6	100	22.3	* 8.9 - 35.8	200	66.7	56.0 - 77.4	1,200	93.8	90.7 - 96.9	4,700
Infant health insurance															
Medi-Cal	30.3	25.6 - 35.1	2,600	80.5	* 67.5 - 93.5	700	85.2	* 73.1 - 97.3	800	39.3	28.0 - 50.6	700	7.4	4.0 - 10.8	400
Uninsured	1.5 *	0.3 - 2.6	100												
Private	67.9	63.1 - 72.7	5,900	13.9	* 2.7 - 25.0	100	14.8	* 2.7 - 26.9	100	56.1	44.6 - 67.5	1,000	92.1	88.7 - 95.6	4,600
Maternal Demographics															
Unmarried	25.5	21.0 - 29.9	2,200	73.3	59.2 - 87.4	600	60.8	45.0 - 76.7	600	41.2	29.9 - 52.4	800	4.1	2.0 - 6.2	200
Race/Ethnicity															
White	32.7	27.8 - 37.6	2,800				4.5	* 0.5 - 8.5	<100	22.4	12.6 - 32.1	400	46.5	39.8 - 53.1	2,200
Hispanic	31.0	26.1 - 35.9	2,600	88.4	* 77.2 - 99.5	700	75.7	61.7 - 89.8	700	37.6	26.0 - 49.1	700	8.9	5.3 - 12.5	400
Asian/Pacific Islander	34.5	29.7 - 39.3	2,900				16.8	* 3.1 - 30.5	200	36.8	25.8 - 47.9	700	43.3	36.8 - 49.8	2,000
Black	1.6	1.3 - 1.9	100				3.0	* 1.2 - 4.7	<100	3.2	1.9 - 4.5	<100	0.9	0.6 - 1.2	<100
Born outside the U.S.	49.4	44.4 - 54.5	4,400	86.6	* 76.0 - 97.1	800	51.9	35.5 - 68.2	500	45.7	34.2 - 57.2	800	44.2	37.7 - 50.7	2,200
Speaks non-English language at home	40.7	35.7 - 45.8	3,500	79.1	* 66.4 - 91.9	700	58.6	42.6 - 74.5	600	38.9	27.7 - 50.1	700	31.3	25.2 - 37.4	1,500
Lives in a high poverty neighborhood	5.6	3.3 - 7.9	500	23.4	* 9.3 - 37.4	200				4.3	* 0.6 - 7.9	<100	2.4	* 0.4 - 4.3	100
Income as a percent of the Federal Poverty Guideline															
0-100%	21.8	17.3 - 26.3	1,800	67.9	50.6 - 85.2	400	63.9	48.1 - 79.6	600	24.9	14.9 - 34.9	400	5.1	* 2.1 - 8.2	200
101-200%	14.3	10.5 - 18.1	1,200	30.1	13.0 - 47.2	200	24.6	11.0 - 38.3	200	32.4	21.1 - 43.8	600	3.1	* 0.9 - 5.2	100
> 200%	63.9	58.7 - 69.0	5,200				11.5	* 0.6 - 22.4	100	42.7	31.1 - 54.3	800	91.8	88.1 - 95.5	4,300

<sup>--</sup> Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of San Mateo County was 548. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

<sup>\*</sup> Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

<sup>†</sup> Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



# MIHA Data Snapshot, San Mateo County by Geographical Area, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

Total  Health Status before Pregnancy	%	95% CI								
		3370 CI	N	%	95%	CI	N	%	95% CI	N
Health Status before Pregnancy	100		8,800	98.3 *	97.2 -	99.4	8,500	1.7 *	0.6 - 2.8	100
In good to excellent health	97.2	95.7 - 98.7	8,600	97.2	95.7 -	98.7	8,200			
Chronic conditions										
Diabetes†	3.0	1.3 - 4.7	300	3.2	1.4 -	4.9	300			
Hypertension†	1.4 *	0.4 - 2.3	100	1.3 *	0.3 -	2.2	100			
Asthma†	5.2	3.0 - 7.5	500	5.4	3.1 -	7.8	500			
Nutrition and Weight										
Daily folic acid use, month before pregnancy	41.8	36.8 - 46.7	3,700	41.9	36.8 -	47.0	3,500	47.2 *	15.5 - 78.8	<100
Overweight before pregnancy	20.2	16.0 - 24.3	1,700	20.4	16.1 -	24.7	1,700			
Obese before pregnancy	15.3	11.5 - 19.1	1,300	14.8	10.9 -	18.6	1,200	59.8 *	29.3 - 90.2	<100
Inadequate weight gain during pregnancy	18.2	14.0 - 22.3	1,400	18.1	13.9 -	22.3	1,300			
Excessive weight gain during pregnancy	38.4	33.1 - 43.8	2,900	38.3	32.9 -	43.8	2,800			
Food insecurity during pregnancy	11.1	7.7 - 14.4	1,000	11.2	7.7 -	14.6	900			
Intimate Partner Violence (IPV) and Depressive Symptoms										
Physical or psychological IPV during pregnancy	4.6	2.5 - 6.7	400	4.5	2.4 -	6.6	400			
Prenatal depressive symptoms	8.5	5.8 - 11.3	700	8.8	6.0 -	11.7	700			
Postpartum depressive symptoms	10.4	7.3 - 13.5	900	10.5	7.4 -	13.7	900			
Hardships and Support during Pregnancy										
Experienced two or more hardships during childhood	15.9	12.2 - 19.6	1,400	15.7	12.0 -	19.5	1,300			
Homeless or did not have a regular place to sleep	3.0 *	1.2 - 4.8	300	3.1 *	1.2 -	4.9	300			
Moved due to problems paying rent or mortgage	5.7	3.1 - 8.3	500	5.2	2.6 -	7.7	400			
Woman or partner lost job	9.8	6.7 - 12.9	900	9.8	6.6 -	13.0	800			
Woman or partner had pay or hours cut back	7.1	4.4 - 9.8	600	7.0	4.3 -	9.7	600			
Became separated or divorced	3.1	1.4 - 4.9	300	2.8	1.2 -	4.5	200			
Had no practical or emotional support	5.7	3.1 - 8.2	500	5.9	3.3 -	8.6	500			
Substance Use										
Any smoking, 3 months before pregnancy	8.6	5.8 - 11.5	800	8.5	5.6 -	11.4	700			
Any smoking, 3rd trimester†	1.1 *	0.3 - 2.0	<100	0.9 *	0.2 -	1.6	<100			
Any smoking, postpartum	2.7	1.2 - 4.3	200	2.9	1.2 -	4.5	200			
Any binge drinking, 3 months before pregnancy	13.6	10.1 - 17.1	1,200	14.0	10.4 -	17.6	1,200			
Any alcohol use, 3rd trimester†	11.3	8.1 - 14.5	1,000	11.7	8.4 -	15.1	1,000			
Pregnancy Intention and Family Planning										
Mistimed or unwanted pregnancy	22.7	18.4 - 27.0	2,000	22.4	18.0 -	26.8	1,900			
Unsure of pregnancy intentions	7.8	5.1 - 10.6	700	8.0	5.2 -	10.8	700			
Postpartum birth control use†	79.1	74.9 - 83.2	6,800	78.4	74.2 -	82.7	6,500			
Infant Sleep and Breastfeeding										
Placed infant on back to sleep	88.6	85.3 - 91.8	7,700	88.2	84.8 -	91.5	7,300			
Infant always or often shared bed	30.1	25.4 - 34.8	2,600	30.3	25.4 -	35.1	2,500			
Intended to breastfeed, before birth	92.9	90.1 - 95.7	8,100	93.0	90.2 -	95.9	7,800			
Intended to breastfeed exclusively, before birth	64.7	59.8 - 69.6	5,600	64.8	59.8 -	69.8	5,400	49.6 *	16.5 - 82.7	<100
Breastfeeding practices	5-1.7	33.3	2,230	04.0	0	0	-,0	75.0		.200
Any breastfeeding, 1 month after delivery	89.7	86.6 - 92.8	7,700	89.7	86.6 -	92.9	7,500			
Exclusive breastfeeding, 1 month after delivery	44.6	39.4 - 49.7	3,800	44.4	39.2 -	49.7	3,600			
Any breastfeeding, 3 months after delivery	78.2	73.3 - 83.0	5,000	78.1	73.2 -	83.1	4,800			
Exclusive breastfeeding, 3 months after delivery	32.8	27.1 - 38.5	2,100	32.6	26.8 -	38.4	2,000			



# MIHA Data Snapshot, San Mateo County by Geographical Area, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		San Mateo Count	:у		Urban			Rural/Frontier	
	%	95% CI	N	%	95% CI	N	%	95% CI	N
Service Utilization and Health Insurance Coverage									
Had a usual source of pre-pregnancy care	77.7	73.5 - 81.9	6,800	78.1	73.8 - 82.4	6,600			
Initiated prenatal care in 1st trimester	92.4	89.6 - 95.1	7,900	92.3	89.4 - 95.1	7,600			
Mom or infant needed but couldn't afford care postpartum	5.7	3.6 - 7.8	500	5.9	3.8 - 8.1	500			
Participated in WIC during pregnancy†	30.5	25.7 - 35.4	2,600	29.7	24.8 - 34.6	2,400			
Received CalFresh (food stamps) during pregnancy	9.1	6.3 - 12.0	800	8.7	5.9 - 11.6	700			
Pre-pregnancy insurance									
Medi-Cal	15.2	11.5 - 18.9	1,300	15.1	11.3 - 18.8	1,300			
Uninsured	9.6	6.5 - 12.6	800	9.3	6.3 - 12.4	800			
Private	72.0	67.4 - 76.5	6,300	72.8	68.2 - 77.5	6,200			
Prenatal insurance									
Medi-Cal	26.3	21.8 - 30.8	2,300	26.2	21.6 - 30.8	2,200			
Uninsured									
Private	71.7	67.1 - 76.3	6,300	72.5	67.8 - 77.1	6,100			
Maternal postpartum insurance									
Medi-Cal	17.5	13.6 - 21.4	1,500	17.2	13.2 - 21.2	1,400			
Uninsured	7.1	4.4 - 9.8	600	6.8	4.1 - 9.4	600			
Private	72.4	67.8 - 77.0	6,300	73.3	68.7 - 78.0	6,200	67.1 *	35.2 - 99.1	<100
Infant health insurance									
Medi-Cal	30.3	25.6 - 35.1	2,600	29.4	24.5 - 34.2	2,500			
Uninsured	1.5 *	0.3 - 2.6	100	1.5 *	0.3 - 2.7	100			
Private	67.9	63.1 - 72.7	5,900	68.8	63.9 - 73.7	5,800	56.9 *	22.9 - 91.0	<100
Maternal Demographics									
Age									
15-19	2.0 *	0.6 - 3.4	200	2.0 *	0.6 - 3.5	200			
20-34	66.6	61.7 - 71.4	5,900	66.2	61.2 - 71.2	5,600	60.2 *	30.0 - 90.4	<100
35+	31.4	26.6 - 36.2	2,800	31.7	26.8 - 36.7	2,700			
Did not complete high school (or GED)	10.0	6.9 - 13.0	900	9.2	6.2 - 12.2	800			
Unmarried	25.5	21.0 - 29.9	2,200	24.3	19.8 - 28.8	2,100	47.8 *	15.5 - 80.0	<100
Race/Ethnicity									
White	32.7	27.8 - 37.6	2,800	32.9	27.8 - 37.9	2,700			
Hispanic	31.0	26.1 - 35.9	2,600	30.3	25.3 - 35.3	2,500	67.3 *	40.2 - 94.4	<100
Asian/Pacific Islander	34.5	29.7 - 39.3	2,900	35.1	30.1 - 40.0	2,800			
Black	1.6	1.3 - 1.9	100	1.6	1.4 - 1.9	100			
Born outside the U.S.	49.4	44.4 - 54.5	4,400	49.0	43.8 - 54.1	4,200			
Speaks non-English language at home	40.7	35.7 - 45.8	3,500	40.1	35.0 - 45.2	3,300			
Lives in a high poverty neighborhood	5.6	3.3 - 7.9	500	5.7	3.3 - 8.0	500			
Income as a percent of the Federal Poverty Guideline									
0-100%	21.8	17.3 - 26.3	1,800	21.6	17.0 - 26.2	1,700			
101-200%	14.3	10.5 - 18.1	1,200	14.1	10.2 - 18.0	1,100			
> 200%	63.9	58.7 - 69.0	5,200	64.3	59.0 - 69.5	5,000	48.1 *	14.9 - 81.4	<100

<sup>--</sup> Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of San Mateo County was 548. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. Urban and Rural/Frontier designations are based on the population size or densities of Medical Service Study Areas. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

<sup>\*</sup> Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

<sup>†</sup> Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



# MIHA Data Snapshot, San Mateo County by Income, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	San Mateo Coun	ity	0-100% Federal Poverty	Guideline	101-200% Federal Poverty	Guideline	> 200% Federal Poverty	Guideline
	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N
Total	100	8,800	21.8 17.3 - 26.3	1,800	14.3 10.5 - 18.1	1,200	63.9 58.7 - 69.0	5,200
Health Status before Pregnancy								
In good to excellent health	<b>97.2</b> 95.7 - 98.7	8,600	<b>92.5</b> * 87.5 - 97.4	1,600				
Chronic conditions								
Diabetes†	3.0 1.3 - 4.7	300			9.9 * 1.3 - 18.5	100		
Hypertension†	1.4 * 0.4 - 2.3	100						
Asthma†	<b>5.2</b> 3.0 - 7.5	500	9.5 * 2.0 - 17.0	200			5.4 2.7 - 8.2	300
Nutrition and Weight								
Daily folic acid use, month before pregnancy	41.8 36.8 - 46.7	3,700	<b>26.8</b> 15.8 - 37.8	500	27.9 15.0 - 40.8	300	<b>51.2</b> 44.8 - 57.6	2,700
Overweight before pregnancy	20.2 16.0 - 24.3	1,700	<b>20.5</b> 10.5 - 30.5	300	<b>25.1</b> 12.3 - 37.9	300	20.2 14.8 - 25.5	1,000
Obese before pregnancy	<b>15.3</b> 11.5 - 19.1	1,300	34.7 22.4 - 47.1	600	20.3 9.0 - 31.6	200	7.7 4.4 - 11.1	400
Inadequate weight gain during pregnancy	18.2 14.0 - 22.3	1,400	<b>21.8</b> 12.0 - 31.7	300	39.0 22.1 - 55.8	300	<b>11.7</b> 7.9 - <b>1</b> 5.4	500
Excessive weight gain during pregnancy	38.4 33.1 - 43.8	2,900	43.8 30.4 - 57.1	600	<b>32.1</b> 15.6 - 48.7	300	38.5 32.0 - 45.1	1,800
Food insecurity during pregnancy	11.1 7.7 - 14.4	1,000	36.0 24.1 - 47.9	600	<b>16.9</b> * 6.0 - 27.7	200	2.3 * 0.7 - 3.8	100
Intimate Partner Violence (IPV) and Depressive Symptoms								
Physical or psychological IPV during pregnancy	4.6 2.5 - 6.7	400	15.3 6.7 - 23.9	300			1.7 * 0.3 - 3.0	<100
Prenatal depressive symptoms	8.5 5.8 - 11.3	700	15.9 7.8 - 23.9	300			5.3 2.5 - 8.0	300
Postpartum depressive symptoms	10.4 7.3 - 13.5	900	15.5 6.7 - 24.3	300	<b>15.5</b> * 5.2 - 25.9	200	7.9 4.4 - 11.3	400
Hardships and Support during Pregnancy								
Experienced two or more hardships during childhood	<b>15.9</b> 12.2 - 19.6	1,400	31.8 20.4 - 43.2	600	<b>19.3</b> 8.3 - 30.3	200	<b>9.2</b> 5.8 - 12.5	500
Homeless or did not have a regular place to sleep	3.0 * 1.2 - 4.8	300	8.1 * 1.7 - 14.5	100				
Moved due to problems paying rent or mortgage	5.7 3.1 - 8.3	500	<b>19.5</b> 9.3 - 29.7	300				
Woman or partner lost job	9.8 6.7 - 12.9	900	23.5 12.8 - 34.1	400	7.8 * 0.6 - 15.0	<100	4.5 2.0 - 7.1	200
Woman or partner had pay or hours cut back	<b>7.1</b> 4.4 - 9.8	600	12.7 * 4.4 - 21.0	200	17.0 * 6.0 - 28.0	200	2.4 * 0.6 - 4.1	100
Became separated or divorced	3.1 1.4 - 4.9	300	<b>12.9</b> * 5.2 - 20.5	200				
Had no practical or emotional support	5.7 3.1 - 8.2	500	16.2 6.9 - 25.6	300				
Substance Use								
Any smoking, 3 months before pregnancy	8.6 5.8 - 11.5	800	<b>15.5</b> 7.2 - 23.8	300	<b>15.4</b> * 5.4 - 25.4	200	5.0 * 2.0 - 8.1	300
Any smoking, 3rd trimester†	1.1 * 0.3 - 2.0	<100						
Any smoking, postpartum	2.7 1.2 - 4.3	200	6.3 * 1.0 - 11.7	100				
Any binge drinking, 3 months before pregnancy	13.6 10.1 - 17.1	1,200	8.3 * 1.3 - 15.3	100	<b>22.3</b> 9.9 - 34.6	300	<b>14.2</b> 9.8 - 18.6	700
Any alcohol use, 3rd trimester†	11.3 8.1 - 14.5	1,000					16.1 11.4 - 20.7	800
Pregnancy Intention and Family Planning								
Mistimed or unwanted pregnancy	22.7 18.4 - 27.0	2,000	46.2 34.3 - 58.0	800	27.6 15.1 - 40.1	300	11.9 7.6 - 16.3	600
Unsure of pregnancy intentions	7.8 5.1 - 10.6	700	7.3 * 1.9 - 12.6	100	17.7 * 6.4 - 28.9	200	<b>6.2</b> 3.0 - 9.4	300
Postpartum birth control use†	<b>79.1</b> 74.9 - 83.2	6,800	88.0 * 80.5 - 95.5	1,500	91.0 * 84.0 - 98.1	1,100	<b>74.4</b> 68.8 - 80.0	3,800
Infant Sleep and Breastfeeding								
Placed infant on back to sleep	88.6 85.3 - 91.8	7,700	<b>79.2</b> 69.0 - 89.4	1,400	<b>85.6</b> * 75.2 - 96.0	900	92.7 89.5 - 95.8	4,800
Infant always or often shared bed	30.1 25.4 - 34.8	2,600	38.7 27.0 - 50.3	700	35.9 21.5 - 50.4	400	<b>25.2</b> 19.8 - 30.6	1,300
Intended to breastfeed, before birth	92.9 90.1 - 95.7	8,100	88.5 * 81.6 - 95.4	1,600	<b>88.9</b> * 78.2 - 99.7	1,000	<b>95.5</b> * 92.6 - 98.4	4,900
Intended to breastfeed exclusively, before birth	64.7 59.8 - 69.6	5,600	48.4 36.6 - 60.1	900	54.8 40.1 - 69.5	600	73.1 67.3 - 78.9	3,800
Breastfeeding practices								
Any breastfeeding, 1 month after delivery	89.7 86.6 - 92.8	7,700	80.8 72.0 - 89.6	1,400	<b>81.3</b> * 69.2 - 93.5	900	93.8 90.8 - 96.8	4,800
Exclusive breastfeeding, 1 month after delivery	<b>44.6</b> 39.4 - 49.7	3,800	<b>39.5</b> 27.6 - 51.4	700	<b>37.9</b> 23.6 - 52.2	400	48.0 41.6 - 54.5	2,400
Any breastfeeding, 3 months after delivery	<b>78.2</b> 73.3 - 83.0	5,000	56.5 43.1 - 70.0	700	<b>68.9</b> 53.0 - 84.9	600	<b>87.5</b> 82.5 - 92.4	3,100
Exclusive breastfeeding, 3 months after delivery	<b>32.8</b> 27.1 - 38.5	2,100	<b>24.4</b> 11.3 - 37.5	300	<b>16.5</b> * 5.8 - 27.3	200	41.7 34.0 - 49.4	1,500



## MIHA Data Snapshot, San Mateo County by Income, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	S	an Mateo Coun	ity	0-100% Fe	deral Poverty	Guideline	101-200% F	ederal Povert	y Guideline	> 200% Fe	ederal Poverty	Guideline
	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N
Service Utilization and Health Insurance Coverage												
Had a usual source of pre-pregnancy care	77.7	73.5 - 81.9	6,800	64.7	53.5 - 75.8	1,200	<b>72.5</b> 5	9.3 - 85.6	800	84.9	80.2 - 89.5	4,400
Initiated prenatal care in 1st trimester	92.4	89.6 - 95.1	7,900	87.7	80.9 - 94.5	1,500	88.3 * 7	7.5 - 99.1	1,000	96.3 *	93.7 - 98.8	4,900
Mom or infant needed but couldn't afford care postpartum	5.7	3.6 - 7.8	500	11.3	5.0 - 17.6	200	9.6 *	1.9 - 17.3	100	3.0 *	1.0 - 5.0	200
Participated in WIC during pregnancy†	30.5	25.7 - 35.4	2,600	82.3	73.4 - 91.1	1,500	63.1 4	8.9 - 77.3	700	1.7 *	0.3 - 3.1	<100
Received CalFresh (food stamps) during pregnancy	9.1	6.3 - 12.0	800	31.2	20.6 - 41.9	600	15.7 *	6.0 - 25.5	200			
Pre-pregnancy insurance												
Medi-Cal	15.2	11.5 - 18.9	1,300	44.4	32.7 - 56.1	800	26.1 1	.3.4 - 38.9	300			
Uninsured	9.6	6.5 - 12.6	800	29.2	18.2 - 40.2	500	11.7 *	3.1 - 20.3	100	2.0 *	0.2 - 3.8	100
Private	72.0	67.4 - 76.5	6,300	22.8	12.8 - 32.7	400	54.4 4	0.1 - 68.8	600	95.9 *	93.4 - 98.3	5,000
Prenatal insurance												
Medi-Cal	26.3	21.8 - 30.8	2,300	77.8	68.1 - 87.6	1,400	47.8 3	3.4 - 62.2	600	1.2 *	0.1 - 2.4	<100
Uninsured												
Private	71.7	67.1 - 76.3	6,300	17.9	9.2 - 26.6	300	<b>52.2</b> 3	7.8 - 66.6	600	97.3 *	95.4 - 99.2	5,100
Maternal postpartum insurance												
Medi-Cal	17.5	13.6 - 21.4	1,500	52.7	41.0 - 64.5	900	30.7 1	.7.3 - 44.1	300	0.9 *	0.0 - 1.8	<100
Uninsured	7.1	4.4 - 9.8	600	24.3	13.8 - 34.8	400	10.1 *	2.6 - 17.6	100			
Private	72.4	67.8 - 77.0	6,300	16.3	8.3 - 24.3	300	55.3 4	0.8 - 69.8	600	98.0 *	96.6 - 99.5	5,100
Infant health insurance												
Medi-Cal	30.3	25.6 - 35.1	2,600	86.3	78.7 - 93.8	1,500	60.2 4	5.4 - 74.9	700	2.4 *	0.7 - 4.0	100
Uninsured	1.5	* 0.3 - 2.6	100									
Private	67.9	63.1 - 72.7	5,900	10.6 *	3.9 - 17.2	200	35.6 2	1.0 - 50.3	400	96.9 *	95.0 - 98.8	5,000
Maternal Demographics												
Did not complete high school (or GED)	10.0	6.9 - 13.0	900	24.8	14.7 - 34.9	400	16.6 *	6.2 - 26.9	200			
Unmarried	25.5	21.0 - 29.9	2,200	66.4	55.5 - 77.2	1,200	49.4 3	4.9 - 64.0	600	5.6	3.0 - 8.3	300
Race/Ethnicity												
White	32.7	27.8 - 37.6	2,800	10.6 *	3.3 - 17.9	200				44.8	38.2 - 51.4	2,200
Hispanic	31.0	26.1 - 35.9	2,600	70.2	59.6 - 80.8	1,200	57.6 4	2.8 - 72.4	600	10.7	6.6 - 14.7	500
Asian/Pacific Islander	34.5	29.7 - 39.3	2,900	15.2	6.5 - 23.8	300	36.7 2	2.1 - 51.3	400	43.1	36.8 - 49.5	2,100
Black	1.6	1.3 - 1.9	100	4.1	2.5 - 5.7	<100	0.7 *	0.2 - 1.2	<100	1.0	0.7 - 1.3	<100
Born outside the U.S.	49.4	44.4 - 54.5	4,400	62.2	50.8 - 73.5	1,100	64.7 5	0.7 - 78.8	800	38.6	32.4 - 44.9	2,000
Speaks non-English language at home	40.7	35.7 - 45.8	3,500	55.5	43.7 - 67.2	1,000	66.9 5	3.2 - 80.7	800	26.1	20.5 - 31.8	1,300
Lives in a high poverty neighborhood	5.6	3.3 - 7.9	500	13.9	5.8 - 21.9	200	10.5 *	2.3 - 18.6	100			

<sup>--</sup> Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of San Mateo County was 548. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

<sup>\*</sup> Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

<sup>†</sup> Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



# MIHA Data Snapshot, San Mateo County by Maternal Age, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	San Mateo Cour	nty	15-19 Years		20-34 Years		35 Years or Olde	er
	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N
Total	100	8,800	<b>2.0</b> * 0.6 - 3.4	200	66.6 61.7 - 71.4	5,900	31.4 26.6 - 36.2	2,800
Health Status before Pregnancy								
In good to excellent health	97.2 95.7 - 98.7	8,600			97.1 * 95.2 - 99.0	5,700		
Chronic conditions								
Diabetes†	3.0 1.3 - 4.7	300			<b>3.1</b> * 1.0 - 5.3	200	3.1 * 0.1 - 6.0	<100
Hypertension†	1.4 * 0.4 - 2.3	100					<b>3.2</b> * 0.6 - 5.8	<100
Asthma†	5.2 3.0 - 7.5	500			4.8 2.1 - 7.6	300	5.0 * 1.3 - 8.7	100
Nutrition and Weight								
Daily folic acid use, month before pregnancy	<b>41.8</b> 36.8 - 46.7	3,700			<b>37.</b> 5 31.7 - 43.4	2,200	<b>52.1</b> 42.7 - 61.6	1,400
Overweight before pregnancy	20.2 16.0 - 24.3	1,700			16.5 12.1 - 20.8	900	<b>26.7</b> 17.8 - 35.5	700
Obese before pregnancy	<b>15.3</b> 11.5 - 19.1	1,300			17.4 12.4 - 22.4	1,000	<b>12.1</b> 6.4 - 17.9	300
Inadequate weight gain during pregnancy	18.2 14.0 - 22.3	1,400			18.7 13.6 - 23.7	900	<b>17.5</b> 9.9 - <b>2</b> 5.0	400
Excessive weight gain during pregnancy	38.4 33.1 - 43.8	2,900			39.7 33.3 - 46.2	2,000	<b>33.1</b> 23.5 - 42.7	800
Food insecurity during pregnancy	11.1 7.7 - 14.4	1,000			13.2 8.7 - 17.7	800	6.2 * 1.9 - 10.4	200
Intimate Partner Violence (IPV) and Depressive Symptoms								
Physical or psychological IPV during pregnancy	4.6 2.5 - 6.7	400			5.3 2.4 - 8.1	300	3.4 * 0.6 - 6.3	<100
Prenatal depressive symptoms	8.5 5.8 - 11.3	700			9.9 6.2 - 13.7	600	<b>5.1</b> * 1.6 - 8.5	100
Postpartum depressive symptoms	10.4 7.3 - 13.5	900			12.5 8.4 - 16.7	700	6.2 * 1.7 - 10.6	200
Hardships and Support during Pregnancy								
Experienced two or more hardships during childhood	<b>15.9</b> 12.2 - 19.6	1,400	37.6 * 5.8 - 69.4	<100	18.1 13.3 - 22.9	1,000	10.0 4.7 - 15.4	300
Homeless or did not have a regular place to sleep	3.0 * 1.2 - 4.8	300			3.1 * 0.8 - 5.4	200		
Moved due to problems paying rent or mortgage	5.7 3.1 - 8.3	500			7.9 4.1 - 11.6	500		
Woman or partner lost job	9.8 6.7 - 12.9	900			12.0 7.8 - 16.3	700	5.5 * 1.7 - 9.2	100
Woman or partner had pay or hours cut back	<b>7.1</b> 4.4 - 9.8	600			8.8 5.2 - 12.5	500	3.9 * 0.7 - 7.1	100
Became separated or divorced	3.1 1.4 - 4.9	300			<b>3.1</b> * 1.0 - 5.3	200		
Had no practical or emotional support	5.7 3.1 - 8.2	500			<b>7.2</b> 3.6 - 10.7	400	3.0 * 0.1 - 5.8	<100
Substance Use								
Any smoking, 3 months before pregnancy	8.6 5.8 - 11.5	800	22.9 * 0.9 - 45.0	<100	9.0 5.4 - 12.7	500	6.8 * 2.0 - 11.7	200
Any smoking, 3rd trimester†	1.1 * 0.3 - 2.0	<100						
Any smoking, postpartum	2.7 1.2 - 4.3	200			<b>3.2</b> * 1.0 - 5.4	200		
Any binge drinking, 3 months before pregnancy	13.6 10.1 - 17.1	1,200			14.6 10.1 - 19.0	800	<b>11.1</b> 5.7 - 16.6	300
Any alcohol use, 3rd trimester†	11.3 8.1 - 14.5	1,000			7.3 4.4 - 10.3	400	19.4 11.7 - 27.1	500
Pregnancy Intention and Family Planning								
Mistimed or unwanted pregnancy	22.7 18.4 - 27.0	2,000			25.6 20.2 - 31.1	1,500	13.2 6.5 - 19.8	400
Unsure of pregnancy intentions	<b>7.8</b> 5.1 - 10.6	700			7.9 4.5 - 11.3	500	<b>7.2</b> * 2.7 - 11.7	200
Postpartum birth control use†	79.1 74.9 - 83.2	6,800			81.3 76.7 - 85.8	4,600	75.1 66.5 - 83.8	2,100
Infant Sleep and Breastfeeding								
Placed infant on back to sleep	88.6 85.3 - 91.8	7,700			86.5 82.1 - 90.9	5,000	93.0 * 88.6 - 97.4	2,500
Infant always or often shared bed	30.1 25.4 - 34.8	2,600	64.8 * 37.0 - 92.6	100	29.5 23.7 - 35.2	1,700	29.1 20.7 - 37.5	800
Intended to breastfeed, before birth	92.9 90.1 - 95.7	8,100			92.7 89.2 - 96.2	5,300	94.8 * 90.3 - 99.3	2,600
Intended to breastfeed exclusively, before birth	64.7 59.8 - 69.6	5,600	50.5 * 16.3 - 84.7	<100	68.0 62.1 - 73.8	3,900	58.9 49.6 - 68.2	1,600
Breastfeeding practices		,				•		
Any breastfeeding, 1 month after delivery	89.7 86.6 - 92.8	7,700	66.5 * 35.9 - 97.0	100	91.1 87.7 - 94.6	5,200	88.3 82.1 - 94.5	2,400
Exclusive breastfeeding, 1 month after delivery	44.6 39.4 - 49.7	3,800	<del></del>		<b>50.8</b> 44.6 - 57.0	2,900	<b>34.0</b> 25.0 - 42.9	900
Any breastfeeding, 3 months after delivery	<b>78.2 73.3</b> - <b>83.0</b>	5,000			77.3 71.2 - 83.3	3,200	82.4 74.4 - 90.5	1,700
Exclusive breastfeeding, 3 months after delivery	<b>32.8</b> 27.1 - 38.5	2,100			35.6 28.6 - 42.6	1,500	<b>29.6</b> 19.6 - 39.7	600



# MIHA Data Snapshot, San Mateo County by Maternal Age, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	San Mateo Cou	inty	15-19 Years		20-34 Years		35 Years or Olde	er
	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N
Service Utilization and Health Insurance Coverage								
Had a usual source of pre-pregnancy care	77.7 73.5 - 81.9	6,800	<b>59.6</b> * 24.3 - 94.9	100	73.8 68.4 - 79.2	4,300	87.0 80.8 - 93.1	2,400
Initiated prenatal care in 1st trimester	92.4 89.6 - 95.1	7,900	<b>44.4</b> * 11.8 - 77.0	<100	93.3 90.2 - 96.4	5,400	93.6 * 89.1 - 98.1	2,500
Mom or infant needed but couldn't afford care postpartum	5.7 3.6 - 7.8	500			<b>6.2</b> 3.6 - 8.7	300	<b>5.0</b> * 1.2 - 8.9	100
Participated in WIC during pregnancy†	30.5 25.7 - 35.4	2,600	<b>41.7</b> * 9.8 - 73.6	<100	<b>35.3</b> 29.1 - 41.5	2,000	<b>19.6</b> 12.4 - 26.8	500
Received CalFresh (food stamps) during pregnancy	9.1 6.3 - 12.0	800			<b>11.2</b> 7.3 - 15.0	700	<b>4.1</b> * 0.7 - 7.5	100
Pre-pregnancy insurance								
Medi-Cal	<b>15.2</b> 11.5 - 18.9	1,300	41.3 * 6.1 - 76.5	<100	<b>18.4</b> 13.6 - 23.3	1,100	6.8 * 2.3 - 11.3	200
Uninsured	9.6 6.5 - 12.6	800			<b>12.6</b> 8.3 - 16.8	700		
Private	<b>72.0</b> 67.4 - 76.5	6,300	37.6 * 4.1 - 71.1	<100	66.2 60.4 - 72.1	3,900	<b>86.4</b> 80.5 - 92.3	2,400
Prenatal insurance								
Medi-Cal	26.3 21.8 - 30.8	2,300			<b>31.5</b> 25.7 - 37.3	1,800	<b>12.1</b> 6.5 - 17.6	300
Uninsured								
Private	71.7 67.1 - 76.3	6,300			66.2 60.3 - 72.1	3,900	86.4 80.4 - 92.3	2,400
Maternal postpartum insurance								
Medi-Cal	17.5 13.6 - 21.4	1,500			<b>20.9</b> 15.7 - 26.0	1,200	<b>5.9</b> * 1.9 - 9.8	200
Uninsured	7.1 4.4 - 9.8	600			9.0 5.3 - 12.7	500		
Private	<b>72.4</b> 67.8 - 77.0	6,300			67.0 61.1 - 72.9	3,900	<b>88.2</b> 82.7 - 93.7	2,400
Infant health insurance								
Medi-Cal	30.3 25.6 - 35.1	2,600			36.3 30.3 - 42.4	2,100	<b>13.3</b> 7.4 - 19.3	400
Uninsured	1.5 * 0.3 - 2.6	100			<b>1.9</b> * 0.2 - 3.6	100		
Private	67.9 63.1 - 72.7	5,900			<b>61.2</b> 55.1 - 67.3	3,500	<b>86.1</b> 80.1 - 92.1	2,400
Maternal Demographics								
Did not complete high school (or GED)	10.0 6.9 - 13.0	900	39.9 * 7.4 - 72.4	<100	9.8 6.0 - 13.5	600	8.5 * 3.4 - 13.6	200
Unmarried	<b>25.5</b> 21.0 - 29.9	2,200			30.5 24.7 - 36.3	1,800	10.2 5.1 - 15.3	300
Race/Ethnicity								
White	<b>32.7</b> 27.8 - 37.6	2,800			30.5 24.8 - 36.2	1,700	38.6 29.0 - 48.2	1,000
Hispanic	31.0 26.1 - 35.9	2,600	<b>83.3</b> * 67.0 - 99.6	100	34.8 28.7 - 40.9	1,900	20.0 12.5 - 27.5	500
Asian/Pacific Islander	34.5 29.7 - 39.3	2,900			33.0 27.3 - 38.7	1,800	39.5 30.3 - 48.6	1,100
Black	1.6 1.3 - 1.9	100			1.7 1.3 - 2.1	<100	1.3 0.6 - 1.9	<100
Born outside the U.S.	49.4 44.4 - 54.5	4,400			49.3 43.2 - 55.4	2,900	51.7 42.3 - 61.1	1,400
Speaks non-English language at home	40.7 35.7 - 45.8	3,500			42.3 36.2 - 48.4	2,400	37.7 28.4 - 47.0	1,000
Lives in a high poverty neighborhood	5.6 3.3 - 7.9	500			7.0 3.9 - 10.2	400		

<sup>--</sup> Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of San Mateo County was 548. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

<sup>\*</sup> Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

<sup>†</sup> Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



# MIHA Data Snapshot, San Mateo County by Neighborhood Poverty, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	San Mateo (	County	0-4.9% of Reside		5-9.9% of Reside below Povert		10-19.9% of Residual Below Povert		≥ 20% of Reside below Poverty	
	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N
Total	100	8,800	35.9 31.1 - 40.7	3,100	37.1 32.1 - 42.0	3,200	21.5 17.1 - 25.9	1,900	5.6 3.3 - 7.9	500
Health Status before Pregnancy										
In good to excellent health Chronic conditions	<b>97.2</b> 95.7 - 98	.7 8,600			96.2 * 93.1 - 99.3	3,100	95.5 * 91.8 - 99.2	1,800		
Diabetes†	3.0 1.3 - 4	.7 300			4.1 * 0.8 - 7.4	100				
Hypertension†	1.4 * 0.4 - 2	.3 100			2.3 * 0.1 - 4.5	<100				
Asthma†	5.2 3.0 - 7	.5 500	5.3 * 1.6 - 9.0	200	7.1 3.0 - 11.3	200				
Nutrition and Weight										
Daily folic acid use, month before pregnancy	41.8 36.8 - 46	.7 3,700	49.6 41.4 - 57.9	1,500	40.1 31.9 - 48.3	1,300	39.0 27.5 - 50.6	700	16.8 * 1.3 - 32.2	<100
Overweight before pregnancy	20.2 16.0 - 24	.3 1,700	22.1 14.8 - 29.4	700	15.9 9.7 - 22.0	500	23.8 13.5 - 34.1	400	24.9 * 7.4 - 42.4	100
Obese before pregnancy	<b>15.3</b> 11.5 - 19	.1 1,300	11.9 6.8 - 17.1	400	13.2 7.0 - 19.4	400	<b>19.3</b> 9.6 - 28.9	300	41.7 19.4 - 64.1	200
Inadequate weight gain during pregnancy	18.2 14.0 - 22		11.5 6.4 - 16.6	300	21.7 14.2 - 29.2	600	19.7 10.1 - 29.2	300	36.9 * 13.4 - 60.3	100
Excessive weight gain during pregnancy	<b>38.4</b> 33.1 - 43		42.9 34.2 - 51.6	1,200	30.7 22.3 - 39.0	900	44.2 31.3 - 57.1	700	37.3 * 14.8 - 59.8	100
Food insecurity during pregnancy	11.1 7.7 - 14	.4 1,000	3.0 * 0.4 - 5.6	<100	11.8 6.3 - 17.3	400	19.9 9.8 - 30.0	400	27.8 * 8.3 - 47.2	100
Intimate Partner Violence (IPV) and Depressive Symptoms										
Physical or psychological IPV during pregnancy	4.6 2.5 - 6		3.6 * 1.2 - 6.1	100	4.8 * 1.5 - 8.1	200				
Prenatal depressive symptoms	8.5 5.8 - 11		6.0 2.5 - 9.6	200	6.4 2.7 - 10.1	200	14.9 * 6.0 - 23.7	300	<b>17.4</b> * 3.9 - 30.9	<100
Postpartum depressive symptoms	10.4 7.3 - 13	.5 900	8.7 4.5 - 12.8	300	7.3 * 2.9 - 11.7	200	18.6 8.9 - 28.2	300	14.8 * 1.3 - 28.3	<10
Hardships and Support during Pregnancy										
Experienced two or more hardships during childhood	<b>15.9</b> 12.2 - 19	,	8.0 4.3 - 11.8	200	15.0 8.8 - 21.1	500	28.5 17.8 - 39.3	500	<b>25.4</b> * 6.9 - 43.9	100
Homeless or did not have a regular place to sleep	3.0 * 1.2 - 4						5.7 * 0.5 - 10.9	100		
Moved due to problems paying rent or mortgage	5.7 3.1 - 8				7.2 * 2.7 - 11.7	200	7.7 * 0.3 - 15.1	100		
Woman or partner lost job	9.8 6.7 - 12		4.9 * 1.8 - 8.0	200	9.3 4.4 - 14.3	300	<b>13.4</b> * 5.0 - 21.9	200	29.4 * 8.7 - 50.1	10
Woman or partner had pay or hours cut back	7.1 4.4 - 9		3.6 * 1.0 - 6.3	100	9.4 4.3 - 14.6	300			<b>18.7</b> * 1.6 - 35.7	<10
Became separated or divorced Had no practical or emotional support	3.1 1.4 - 4 5.7 3.1 - 8			-100			5.5 * 0.6 - 10.4	100		
	5.7 3.1 - 8	.2 500	2.8 * 0.4 - 5.2	<100			15.4 * 6.0 - 24.8	300		
Substance Use	0.6 50 44	- 000	C 0 * 22 44 5	200	0.2 40 420	200	44.0* 24.406	200		
Any smoking, 3 months before pregnancy Any smoking, 3rd trimester†	8.6 5.8 - 11		6.8 * 2.2 - 11.5	200	9.3 4.8 - 13.8	300	11.0 * 3.4 - 18.6	200		
Any smoking, 5rd trimester i	1.1 * 0.3 - 2 2.7 1.2 - 4				3.9 * 0.7 - 7.1	100				
Any binge drinking, 3 months before pregnancy	13.6 10.1 - 17		14.9 9.1 - 20.7	500	11.2 5.9 - 16.5	400	14.7 * 6.0 - 23.4	300	<b>22.3</b> * 4.9 - 39.7	<10
Any alcohol use, 3rd trimester†	11.3 8.1 - 14		17.4 11.0 - 23.8	500	11.8 6.1 - 17.4	400	14.7 0.0 - 23.4	300	22.3 4.9 - 33.7	100
Pregnancy Intention and Family Planning	11.5 0.1 14	.5 1,000	17.4 11.0 25.0	300	11.0 0.1 17.4	400				
Mistimed or unwanted pregnancy	22.7 18.4 - 27	.0 2,000	11.6 6.4 - 16.9	400	22.6 15.4 - 29.8	700	35.4 24.1 - 46.7	700	41.7 20.8 - 62.6	200
Unsure of pregnancy intentions	7.8 5.1 - 10	,	9.1 4.2 - 13.9	300	6.7 * 2.2 - 11.2	200	4.1 * 0.2 - 7.9	<100	22.7 * 4.9 - 40.6	100
Postpartum birth control use†	79.1 74.9 - 83		76.9 69.9 - 84.0	2,300	73.6 65.9 - 81.3	2,300	90.2 * 84.1 - 96.3	1,600	81.6 * 65.5 - 97.8	400
Infant Sleep and Breastfeeding	7512	,,,,,,	70.5	,	75.5	,	30.2	,	02.0	
Placed infant on back to sleep	88.6 85.3 - 91	.8 7,700	88.9 84.3 - 93.6	2,600	89.4 84.1 - 94.8	2,800	<b>89.6</b> * 82.3 - 96.9	1,700	<b>72.3</b> * 51.1 - 93.6	300
Infant always or often shared bed	30.1 25.4 - 34	•	26.0 18.9 - 33.1	800	36.8 28.5 - 45.0	1,200	27.2 16.3 - 38.2	500	27.0 * 7.5 - 46.5	100
Intended to breastfeed, before birth	92.9 90.1 - 95	,	97.2 * 94.6 - 99.8	2,900	92.9 88.8 - 97.0	2,900	85.0 * 75.6 - 94.4	1,600		_0
Intended to breastfeed exclusively, before birth	64.7 59.8 - 69		69.9 62.2 - 77.6	2,100	67.6 59.8 - 75.4	2,100	54.3 42.5 - 66.1	1,000	50.6 29.5 - 71.7	200
Breastfeeding practices										
Any breastfeeding, 1 month after delivery	89.7 86.6 - 92	.8 7,700	90.3 85.5 - 95.1	2,700	90.7 85.8 - 95.5	2,800	86.1 * 77.8 - 94.4	1,600		
Exclusive breastfeeding, 1 month after delivery	44.6 39.4 - 49		42.6 34.3 - 51.0	1,300	48.7 40.1 - 57.2	1,500	34.4 23.2 - 45.6	600	62.4 42.8 - 82.0	300
Any breastfeeding, 3 months after delivery	<b>78.2</b> 73.3 - 83		87.5 81.5 - 93.6	1,900	80.8 73.3 - 88.2	1,900	64.7 51.4 - 78.0	900	60.7 38.5 - 83.0	200
Exclusive breastfeeding, 3 months after delivery	<b>32.8</b> 27.1 - 38	.5 2,100	36.8 27.2 - 46.5	800	35.5 26.1 - 44.8	800	23.4 10.6 - 36.2	300	<b>24.0</b> * 5.1 - 42.8	<100



### MIHA Data Snapshot, San Mateo County by Neighborhood Poverty, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	San	Mateo Cou	nty	0-4.9% of Reside		5-9.9% of Reside		10-19.9% of Robelow Pov		≥ 20% of Reside	
	%	95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N
Service Utilization and Health Insurance Coverage											
Had a usual source of pre-pregnancy care	77.7	73.5 - 81.9	6,800	84.3 78.5 - 90.1	2,600	77.5 70.3 - 84.8	2,500	<b>74.8</b> 65.0 - 84	.7 1,400	60.2 39.7 - 80.7	300
Initiated prenatal care in 1st trimester	92.4	89.6 - 95.1	7,900	96.4 * 93.2 - 99.7	2,900	91.2 86.5 - 96.0	2,800	90.6 * 83.3 - 97	.9 1,600	79.9 * 64.5 - 95.3	400
Mom or infant needed but couldn't afford care postpartum	5.7	3.6 - 7.8	500	4.3 * 1.2 - 7.3	100	7.7 3.7 - 11.7	200			<b>15.2</b> * 2.8 - 27.7	<100
Participated in WIC during pregnancy†		25.7 - 35.4	2,600	<b>12.3</b> 7.0 - 17.5	400	<b>25.1</b> 17.6 - 32.7	800	<b>52.8</b> 40.9 - 64	.7 1,000	86.1 * 72.8 - 99.5	400
Received CalFresh (food stamps) during pregnancy	9.1	6.3 - 12.0	800	3.9 * 1.2 - 6.7	100	7.3 3.0 - 11.6	200	16.7 7.9 - 25	.5 300	20.9 * 4.9 - 36.8	100
Pre-pregnancy insurance											
Medi-Cal	15.2	11.5 - 18.9	1,300	<b>5.2</b> * 1.8 - 8.5	200	<b>16.4</b> 9.8 - 22.9	500	23.7 14.0 - 33	.5 400	37.8 16.7 - 58.9	200
Uninsured	9.6	6.5 - 12.6	800	<b>2.7</b> * 0.1 - 5.4	<100	8.5 3.9 - 13.2	300	<b>18.9</b> 9.3 - 28	.6 300	<b>17.2</b> * 4.0 - 30.5	<100
Private	72.0	67.4 - 76.5	6,300	89.8 85.2 - 94.5	2,800	71.0 63.3 - 78.7	2,300	55.5 43.8 - 67	.1 1,000	41.5 20.5 - 62.4	200
Prenatal insurance											
Medi-Cal	26.3	21.8 - 30.8	2,300	9.0 4.8 - 13.2	300	26.7 19.2 - 34.2	900	46.6 34.9 - 58	.3 900	54.9 33.6 - 76.3	300
Uninsured											
Private	71.7	67.1 - 76.3	6,300	89.0 84.2 - 93.7	2,800	<b>72.9</b> 65.4 - 80.4	2,300	53.4 41.7 - 65	.1 1,000	<b>37.2</b> 16.5 - 58.0	200
Maternal postpartum insurance											
Medi-Cal	17.5	13.6 - 21.4	1,500	6.5 3.0 - 10.1	200	17.0 10.7 - 23.2	500	<b>28.9</b> 18.0 - 39	.8 500	46.9 25.1 - 68.7	200
Uninsured	7.1	4.4 - 9.8	600			8.1 * 3.0 - 13.2	300	<b>11.9</b> * 4.6 - 19	.2 200	16.4 * 3.3 - 29.5	<100
Private	72.4	67.8 - 77.0	6,300	90.4 86.1 - 94.8	2,800	73.3 65.7 - 80.8	2,300	53.8 42.1 - 65	.5 1,000	36.7 15.8 - 57.6	200
Infant health insurance											
Medi-Cal	30.3	25.6 - 35.1	2,600	12.1 7.0 - 17.3	400	29.1 21.3 - 36.9	900	48.5 36.7 - 60	.3 900	<b>68.8</b> * 49.0 - 88.6	300
Uninsured	1.5 *	0.3 - 2.6	100								
Private	67.9	63.1 - 72.7	5,900	86.6 81.3 - 92.0	2,600	67.9 59.8 - 75.9	2,100	50.2 38.5 - 62	.0 900	31.2 * 11.4 - 51.0	200
Maternal Demographics											
Age											
15-19	2.0 *	0.6 - 3.4	200			<b>3.6</b> * 0.3 - 6.9	100				
20-34	66.6	61.7 - 71.4	5,900	56.0 47.7 - 64.3	1,700	66.3 58.1 - 74.5	2,100	<b>78.3</b> 68.8 - 87	.8 1,400	83.3 * 68.2 - 98.4	400
35+	31.4	26.6 - 36.2	2,800	43.7 35.4 - 52.1	1,400	30.1 22.1 - 38.0	1,000	19.0 9.8 - 28	.2 400		
Did not complete high school (or GED)	10.0	6.9 - 13.0	900			9.5 4.9 - 14.2	300	<b>13.8</b> 5.8 - 21	.9 300	40.7 19.5 - 61.9	200
Unmarried	25.5	21.0 - 29.9	2,200	10.4 5.8 - 15.0	300	<b>25.4</b> 18.1 - 32.8	800	<b>39.8</b> 28.3 - 51	.4 700	53.0 32.0 - 73.9	300
Race/Ethnicity											
White	32.7	27.8 - 37.6	2,800	46.6 38.1 - 55.1	1,400	33.2 25.0 - 41.4	1,000	13.7 6.2 - 21	.2 200		
Hispanic	31.0	26.1 - 35.9	2,600	13.6 8.2 - 19.0	400	26.4 18.6 - 34.1	800	56.1 44.6 - 67	.6 1,000	<b>74.5</b> * 55.3 - 93.6	300
Asian/Pacific Islander	34.5	29.7 - 39.3	2,900	38.6 30.6 - 46.6	1,100	<b>39.3</b> 30.9 - 47.6	1,200	26.7 16.6 - 36	.8 500		
Black	1.6	1.3 - 1.9	100	1.0 0.4 - 1.5	<100	1.2 0.7 - 1.6	<100	3.5 2.1 - 4	.9 <100		
Born outside the U.S.	49.4	44.4 - 54.5	4,400	<b>42.1</b> 34.0 - 50.3	1,300	49.5 41.1 - 57.9	1,600	54.0 42.3 - 65	.7 1,000	66.9 48.0 - 85.7	300
Speaks non-English language at home	40.7	35.7 - 45.8	3,500	31.9 24.3 - 39.6	1,000	40.8 32.3 - 49.2	1,300	49.6 37.8 - 61	.4 900	47.1 26.1 - 68.1	200
Income as a percent of the Federal Poverty Guideline	-			-	•		•				
0-100%	21.8	17.3 - 26.3	1,800	7.4 3.2 - 11.6	200	19.8 12.6 - 27.0	600	40.8 28.6 - 52	.9 700	54.8 32.8 - 76.9	200
101-200%	_	10.5 - 18.1	1,200	8.7 3.8 - 13.6	300	15.8 9.1 - 22.5	500	18.1 8.3 - 27		27.3 * 8.1 - 46.6	100
> 200%	63.9	58.7 - 69.0	5,200	83.9 77.7 - 90.0	2,500	64.4 55.8 - 73.0	1,900	41.2 29.3 - 53	.1 700	17.8 * 1.0 - 34.7	<100

<sup>--</sup> Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of San Mateo County was 548. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. Neighborhood poverty is the percentage of residents with incomes below poverty in a woman's census tract. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

<sup>\*</sup> Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

<sup>†</sup> Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



### MIHA Data Snapshot, San Mateo County by Prenatal Health Insurance, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		San Mateo Count	ty		Medi-Cal			Private	
	%	95% CI	N	%	95% CI	N	%	95% CI	N
Total	100		8,800	26.3	21.8 - 30.8	2,300	71.7	67.1 - 76.3	6,300
Health Status before Pregnancy									
In good to excellent health	97.2	95.7 - 98.7	8,600	92.7 *	88.4 - 97.0	2,100			
Chronic conditions									
Diabetes†	3.0	1.3 - 4.7	300	5.9 *	1.6 - 10.1	100	1.9 *	0.1 - 3.6	100
Hypertension†	1.4 *	0.4 - 2.3	100						
Asthma†	5.2	3.0 - 7.5	500	6.2 *	0.9 - 11.6	100	5.0	2.6 - 7.5	300
Nutrition and Weight									
Daily folic acid use, month before pregnancy	41.8	36.8 - 46.7	3,700	24.0	15.1 - 33.0	600	48.2	42.3 - 54.0	3,000
Overweight before pregnancy	20.2	16.0 - 24.3	1,700	27.0	17.6 - 36.3	600	18.4	13.6 - 23.1	1,200
Obese before pregnancy	15.3	11.5 - 19.1	1,300	30.2	20.0 - 40.3	600	10.8	7.2 - 14.4	700
Inadequate weight gain during pregnancy	18.2	14.0 - 22.3	1,400	27.9	17.8 - 38.1	500	15.3	10.9 - 19.8	900
Excessive weight gain during pregnancy	38.4	33.1 - 43.8	2,900	40.0	28.2 - 51.7	700	38.1	32.1 - 44.2	2,200
Food insecurity during pregnancy	11.1	7.7 - 14.4	1,000	31.4	21.6 - 41.2	700	4.1	1.8 - 6.5	300
Intimate Partner Violence (IPV) and Depressive Symptoms									
Physical or psychological IPV during pregnancy	4.6	2.5 - 6.7	400	10.8 *	4.4 - 17.2	200	2.2 *	0.7 - 3.8	100
Prenatal depressive symptoms	8.5	5.8 - 11.3	700	15.8	8.7 - 22.9	400	5.8	3.1 - 8.5	400
Postpartum depressive symptoms	10.4	7.3 - 13.5	900	14.8	7.6 - 22.1	300	9.1	5.7 - 12.5	600
Hardships and Support during Pregnancy									
Experienced two or more hardships during childhood	15.9	12.2 - 19.6	1,400	31.4	21.8 - 40.9	700	10.8	7.3 - 14.3	700
Homeless or did not have a regular place to sleep	3.0 *	1.2 - 4.8	300	8.4 *	2.9 - 13.9	200			
Moved due to problems paying rent or mortgage	5.7	3.1 - 8.3	500	15.4	7.5 - 23.3	400	1.7 *	0.2 - 3.3	100
Woman or partner lost job	9.8	6.7 - 12.9	900	24.2	15.4 - 33.1	500	4.3	2.0 - 6.6	300
Woman or partner had pay or hours cut back	7.1	4.4 - 9.8	600	14.8	7.6 - 21.9	300	4.5	1.9 - 7.1	300
Became separated or divorced	3.1	1.4 - 4.9	300	9.6	4.0 - 15.2	200			
Had no practical or emotional support	5.7	3.1 - 8.2	500	16.2	8.1 - 24.3	400	2.0 *	0.4 - 3.6	100
Substance Use									
Any smoking, 3 months before pregnancy	8.6	5.8 - 11.5	800	13.0	6.4 - 19.6	300	7.3	4.1 - 10.4	500
Any smoking, 3rd trimester†	1.1 *	0.3 - 2.0	<100	1.9 *	0.1 - 3.7	<100			
Any smoking, postpartum	2.7	1.2 - 4.3	200	5.6 *	1.3 - 9.9	100	1.8 *	0.3 - 3.3	100
Any binge drinking, 3 months before pregnancy	13.6	10.1 - 17.1	1,200	7.6 *	1.9 - 13.3	200	16.1	11.7 - 20.4	1,000
Any alcohol use, 3rd trimester†	11.3	8.1 - 14.5	1,000				15.3	11.0 - 19.7	1,000
Pregnancy Intention and Family Planning									
Mistimed or unwanted pregnancy	22.7	18.4 - 27.0	2,000	48.0	37.9 - 58.0	1,100	13.6	9.4 - 17.8	900
Unsure of pregnancy intentions	7.8	5.1 - 10.6	700	9.4	4.0 - 14.8	200	7.3	4.1 - 10.5	500
Postpartum birth control use†	79.1	74.9 - 83.2	6,800	87.6	81.5 - 93.6	1,900	76.2	71.1 - 81.4	4,700
Infant Sleep and Breastfeeding									
Placed infant on back to sleep	88.6	85.3 - 91.8	7,700	81.7	73.5 - 89.9	1,900	91.8	88.7 - 94.9	5,700
Infant always or often shared bed	30.1	25.4 - 34.8	2,600	37.2	27.3 - 47.2	800	28.2	22.8 - 33.6	1,700
Intended to breastfeed, before birth	92.9	90.1 - 95.7	8,100	90.0	84.1 - 95.9	2,100	93.8	90.5 - 97.0	5,800
Intended to breastfeed exclusively, before birth	64.7	59.8 - 69.6	5,600	52.2	42.2 - 62.3	1,200	69.2	63.6 - 74.9	4,300
Breastfeeding practices									
Any breastfeeding, 1 month after delivery	89.7	86.6 - 92.8	7,700	86.6	80.2 - 93.0	2,000	90.6	87.0 - 94.2	5,600
Exclusive breastfeeding, 1 month after delivery	44.6	39.4 - 49.7	3,800	41.9	31.7 - 52.0	900	45.5	39.5 - 51.5	2,800
Any breastfeeding, 3 months after delivery	78.2	73.3 - 83.0	5,000	68.8	59.0 - 78.6	1,200	82.6	77.1 - 88.1	3,700
Exclusive breastfeeding, 3 months after delivery	32.8	27.1 - 38.5	2,100	25.2	14.8 - 35.6	500	36.5	29.6 - 43.3	1,600



### MIHA Data Snapshot, San Mateo County by Prenatal Health Insurance, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		San Mateo Count	у		Medi-Cal			Private	
	%	95% CI	N	%	95% CI	N	%	95% CI	N
Service Utilization and Health Insurance Coverage									
Had a usual source of pre-pregnancy care	77.7	73.5 - 81.9	6,800	61.0	51.5 - 70.6	1,400	84.8	80.5 - 89.1	5,400
Initiated prenatal care in 1st trimester	92.4	89.6 - 95.1	7,900	86.7	80.0 - 93.4	2,000	94.5	91.6 - 97.4	5,800
Mom or infant needed but couldn't afford care postpartum	5.7	3.6 - 7.8	500	13.5	7.5 - 19.6	300	2.9 *	1.1 - 4.7	200
Participated in WIC during pregnancy†	30.5	25.7 - 35.4	2,600	84.1	76.7 - 91.6	1,900	9.8	5.9 - 13.6	600
Received CalFresh (food stamps) during pregnancy	9.1	6.3 - 12.0	800	32.8	23.6 - 42.1	800			
Pre-pregnancy insurance									
Medi-Cal	15.2	11.5 - 18.9	1,300	53.4	43.5 - 63.4	1,200			
Uninsured	9.6	6.5 - 12.6	800	31.2	21.9 - 40.5	700			
Private	72.0	67.4 - 76.5	6,300	7.8 *	2.4 - 13.1	200	97.3 *	95.5 - 99.1	6,100
Maternal postpartum insurance									
Medi-Cal	17.5	13.6 - 21.4	1,500	61.8	52.0 - 71.6	1,400			
Uninsured	7.1	4.4 - 9.8	600	23.7	15.0 - 32.4	500			
Private	72.4	67.8 - 77.0	6,300	4.4 *	0.9 - 7.8	<100	98.1 *	96.5 - 99.7	6,200
Infant health insurance									
Medi-Cal	30.3	25.6 - 35.1	2,600	95.2 *	91.6 - 98.8	2,200	5.6	2.7 - 8.4	300
Uninsured	1.5 *	0.3 - 2.6	100						
Private	67.9	63.1 - 72.7	5,900	3.3 *	0.2 - 6.5	<100	92.7	89.5 - 95.9	5,800
Maternal Demographics									
Age									
15-19	2.0 *	0.6 - 3.4	200	5.7 *	1.2 - 10.2	100			
20-34	66.6	61.7 - 71.4	5,900	79.9	72.3 - 87.5	1,800	61.5	55.5 - 67.4	3,900
35+	31.4	26.6 - 36.2	2,800	14.4	7.9 - 21.0	300	37.8	31.9 - 43.8	2,400
Did not complete high school (or GED)	10.0	6.9 - 13.0	900	28.3	19.5 - 37.2	600	2.9 *	0.8 - 5.0	200
Unmarried	25.5	21.0 - 29.9	2,200	64.8	55.4 - 74.1	1,500	10.5	6.9 - 14.1	700
Race/Ethnicity									
White	32.7	27.8 - 37.6	2,800	7.3 *	2.3 - 12.4	200	41.9	35.9 - 48.0	2,500
Hispanic	31.0	26.1 - 35.9	2,600	75.2	67.0 - 83.4	1,600	15.2	10.7 - 19.6	900
Asian/Pacific Islander	34.5	29.7 - 39.3	2,900	14.1	7.3 - 21.0	300	41.6	35.7 - 47.5	2,500
Black	1.6	1.3 - 1.9	100	3.3	2.1 - 4.4	<100	1.0	0.7 - 1.3	<100
Born outside the U.S.	49.4	44.4 - 54.5	4,400	69.4	60.2 - 78.6	1,600	40.8	34.9 - 46.6	2,600
Speaks non-English language at home	40.7	35.7 - 45.8	3,500	68.0	58.5 - 77.4	1,500	30.2	24.6 - 35.8	1,900
Lives in a high poverty neighborhood	5.6	3.3 - 7.9	500	11.7	5.8 - 17.6	300	2.9 *	0.8 - 4.9	200
Income as a percent of the Federal Poverty Guideline							_		
0-100%	21.8	17.3 - 26.3	1,800	69.1	59.5 - 78.6	1,400	5.3	2.6 - 8.1	300
101-200%	14.3	10.5 - 18.1	1,200	27.8	18.5 - 37.1	600	10.2	6.1 - 14.3	600
> 200%	63.9	58.7 - 69.0	5,200	3.2 *	0.2 - 6.1	<100	84.5	79.8 - 89.2	5,100

<sup>--</sup> Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of San Mateo County was 548. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

<sup>\*</sup> Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%)

<sup>†</sup> Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



# MIHA Data Snapshot, San Mateo County by Race/Ethnicity, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	San Mateo County		Hispanic		Black		White	Asian/Pacific Islander		
	% 95%	CI N	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N
Total	100	8,800	31.0 26.1 - 35.9	2,600	1.6 1.3 - 1.9	100	32.7 27.8 - 37.6	2,800	34.5 29.7 - 39.3	2,900
Health Status before Pregnancy										
In good to excellent health	97.2 95.7 -	98.7 8,600	95.4 * 92.0 - 98.7	2,500	91.3 * 85.3 - 97.4	100				
Chronic conditions										
Diabetes†	3.0 1.3 -	4.7 300							6.0 * 1.7 - 10.3	200
Hypertension†	1.4 * 0.4 -	2.3 100			11.2 * 1.7 - 20.7	<100				
Asthma†	5.2 3.0 -	7.5 500	6.5 * 1.2 - 11.7	200	16.3 7.2 - 25.4	<100	7.4 3.1 - 11.6	200		
Nutrition and Weight										
Daily folic acid use, month before pregnancy	41.8 36.8 -	46.7 3,700	31.2 21.8 - 40.6	800	<b>25.4</b> 15.2 - 35.6	<100	46.7 37.6 - 55.9	1,300	49.0 40.7 - 57.4	1,400
Overweight before pregnancy	20.2 16.0 -	24.3 1,700	24.8 15.9 - 33.7	600	22.2 12.1 - 32.3	<100	23.9 15.8 - 32.0	700	14.5 8.5 - 20.5	400
Obese before pregnancy	15.3 11.5 -	19.1 1,300	32.6 22.8 - 42.5	800	29.0 15.8 - 42.3	<100	9.3 4.5 - 14.2	300	8.0 * 3.1 - 12.9	200
Inadequate weight gain during pregnancy	18.2 14.0 -	22.3 1,400	25.7 16.5 - 35.0	500	<b>12.3</b> * 3.5 - 21.1	<100	13.4 6.5 - 20.3	300	19.5 12.7 - 26.4	500
Excessive weight gain during pregnancy	38.4 33.1 -	43.8 2,900	<b>42.0</b> 30.7 - 53.2	800	52.5 37.8 - 67.3	<100	46.8 37.1 - 56.4	1,200	<b>25.2</b> 17.5 - 32.9	700
Food insecurity during pregnancy	11.1 7.7 -	14.4 1,000	23.5 14.9 - 32.1	600	17.0 * 6.9 - 27.0	<100			8.8 * 3.6 - 13.9	300
<b>Intimate Partner Violence (IPV) and Depressive Symptoms</b>										
Physical or psychological IPV during pregnancy	4.6 2.5 -	6.7 400	9.0 * 3.2 - 14.8	200					3.4 * 0.7 - 6.0	<100
Prenatal depressive symptoms	8.5 5.8 -	11.3 700	15.0 8.1 - 21.9	400	<b>15.8</b> * 5.9 - 25.8	<100	6.0 * 2.0 - 9.9	200	5.2 * 1.6 - 8.8	200
Postpartum depressive symptoms	10.4 7.3 -	13.5 900	14.5 7.5 - 21.4	400	11.4 * 4.3 - 18.6	<100	5.6 * 1.1 - 10.1	200	11.9 6.5 - 17.2	300
Hardships and Support during Pregnancy										
Experienced two or more hardships during childhood	15.9 12.2 -	19.6 1,400	30.4 21.3 - 39.6	800	36.4 24.0 - 48.9	<100	8.4 3.8 - 13.0	200	11.1 5.9 - 16.3	300
Homeless or did not have a regular place to sleep	3.0 * 1.2 -	4.8 300	6.2 * 1.2 - 11.3	200	18.7 * 7.6 - 29.7	<100				
Moved due to problems paying rent or mortgage	5.7 3.1 -	8.3 500	8.8 * 2.7 - 14.9	200	11.0 * 1.9 - 20.2	<100	4.7 * 0.7 - 8.8	100	4.4 * 0.3 - 8.4	100
Woman or partner lost job	9.8 6.7 -	12.9 900	19.2 10.9 - 27.5	500	14.3 6.6 - 21.9	<100	6.0 * 2.1 - 9.8	200	5.8 * 2.2 - 9.4	200
Woman or partner had pay or hours cut back	7.1 4.4 -	9.8 600	13.8 6.9 - 20.6	400	9.3 * 2.1 - 16.6	<100	6.5 * 1.9 - 11.1	200	2.6 * 0.2 - 5.1	<100
Became separated or divorced	3.1 1.4 -	4.9 300	6.5 * 1.8 - 11.2	200	22.7 10.7 - 34.6	<100				
Had no practical or emotional support	5.7 3.1 -	8.2 500	10.3 * 4.1 - 16.5	300					7.2 * 2.2 - 12.3	200
Substance Use										
Any smoking, 3 months before pregnancy	8.6 5.8 -	11.5 800	10.9 4.9 - 16.9	300	13.5 * 4.2 - 22.8	<100	8.1 * 2.5 - 13.7	200	7.4 3.3 - 11.6	200
Any smoking, 3rd trimester†	1.1 * 0.3 -	2.0 <100								
Any smoking, postpartum	2.7 1.2 -	4.3 200	3.8 * 0.2 - 7.5	<100	11.1 * 2.3 - 20.0	<100			2.7 * 0.1 - 5.4	<100
Any binge drinking, 3 months before pregnancy	13.6 10.1 -	17.1 1,200	16.0 8.6 - 23.3	400	4.5 * 0.8 - 8.2	<100	18.4 11.5 - 25.3	500	6.7 * 2.7 - 10.8	200
Any alcohol use, 3rd trimester†	11.3 8.1 -	14.5 1,000			<b>2.9</b> * 0.5 - 5.3	<100	<b>24.5</b> 16.3 - 32.6	700	4.9 * 1.7 - 8.0	100
Pregnancy Intention and Family Planning										
Mistimed or unwanted pregnancy	22.7 18.4 -	27.0 2,000	35.6 26.2 - 44.9	900	32.7 20.9 - 44.6	<100	18.2 10.6 - 25.9	500	15.5 9.4 - 21.6	400
Unsure of pregnancy intentions	7.8 5.1 -	10.6 700	9.2 3.8 - 14.6	200	17.3 8.6 - 26.0	<100	4.2 * 0.4 - 8.1	100	10.2 4.8 - 15.7	300
Postpartum birth control use†	79.1 74.9 -	83.2 6,800	86.3 80.0 - 92.7	2,200	80.7 71.0 - 90.3	100	76.1 67.5 - 84.7	2,000	75.8 68.8 - 82.9	2,200
Infant Sleep and Breastfeeding										
Placed infant on back to sleep	88.6 85.3 -	91.8 7,700	86.1 79.0 - 93.2	2,200	65.4 53.0 - 77.8	<100	90.4 85.5 - 95.3	2,400	91.2 86.2 - 96.2	2,600
Infant always or often shared bed	30.1 25.4 -		38.9 29.3 - 48.5	1,000	46.6 33.9 - 59.3	<100	14.2 7.6 - 20.7	400	36.8 28.6 - 45.1	1,100
Intended to breastfeed, before birth	92.9 90.1 -	95.7 8,100	86.9 79.6 - 94.2	2,200	92.0 * 85.9 - 98.2	100	95.6 * 92.1 - 99.0	2,500	94.9 * 91.0 - 98.7	2,800
Intended to breastfeed exclusively, before birth	64.7 59.8 -	69.6 5,600	52.0 42.3 - 61.8	1,300	75.0 64.3 - 85.8	100	80.0 72.3 - 87.7	2,100	59.1 50.9 - 67.4	1,700
Breastfeeding practices										
Any breastfeeding, 1 month after delivery	89.7 86.6 -	92.8 7,700	86.6 80.0 - 93.1	2,200	87.7 * 79.9 - 95.5	100	93.6 * 89.5 - 97.8	2,500	87.8 81.9 - 93.8	2,500
Exclusive breastfeeding, 1 month after delivery	44.6 39.4 -	49.7 3,800	40.0 30.3 - 49.8	1,000	59.4 46.5 - 72.4	<100	52.5 43.1 - 62.0	1,400	38.7 30.5 - 46.9	1,100
Any breastfeeding, 3 months after delivery	78.2 73.3 -	83.0 5,000	67.2 57.0 - 77.3	1,400	54.8 41.2 - 68.4	<100	85.8 78.4 - 93.1	1,600	81.2 73.0 - 89.5	1,600
Exclusive breastfeeding, 3 months after delivery	32.8 27.1 -	38.5 2,100	20.7 11.8 - 29.7	400	26.9 16.1 - 37.7	<100	46.3 34.6 - 57.9	900	32.6 23.1 - 42.2	600



#### MIHA Data Snapshot, San Mateo County by Race/Ethnicity, 2013-2015

Maternal and Infant Health Assessment (MIHA) Survey

	San Mateo County		Hispanic			Black			White			Asian/Pacific Islander			
	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N	%	95% CI	N
Service Utilization and Health Insurance Coverage															
Had a usual source of pre-pregnancy care	77.7	73.5 - 81.9	6,800	71.3	62.7 - 79.9	1,900	83.3	74.7 - 91.8	100	83.1	76.3 - 89.8	2,300	78.9	71.9 - 86.0	2,300
Initiated prenatal care in 1st trimester	92.4	89.6 - 95.1	7,900	87.7	81.0 - 94.4	2,200	88.8	* 81.7 - 96.0	100				93.0	88.9 - 97.1	2,600
Mom or infant needed but couldn't afford care postpartum	5.7	3.6 - 7.8	500	9.6	4.8 - 14.4	200	9.9 *	* 2.2 - 17.6	<100	5.1	* 1.3 - 8.9	100	3.2	* 0.3 - 6.0	<100
Participated in WIC during pregnancy†	30.5	25.7 - 35.4	2,600	70.3	61.4 - 79.2	1,800	57.2	44.8 - 69.6	<100	6.5	* 1.6 - 11.5	200	15.3	8.8 - 21.7	400
Received CalFresh (food stamps) during pregnancy	9.1	6.3 - 12.0	800	19.6	12.1 - 27.0	500	43.1	30.6 - 55.6	<100				5.2	* 0.9 - 9.5	200
Pre-pregnancy insurance															
Medi-Cal	15.2	11.5 - 18.9	1,300	36.1	26.6 - 45.6	900	39.0	26.1 - 51.8	<100	3.5	* 0.3 - 6.7	<100	6.9	* 2.8 - 11.0	200
Uninsured	9.6	6.5 - 12.6	800	19.3	11.7 - 26.9	500	10.2 *	* 3.1 - 17.3	<100	4.0	* 0.2 - 7.7	100	6.3	* 1.6 - 11.0	200
Private	72.0	67.4 - 76.5	6,300	37.8	28.3 - 47.2	1,000	47.8	35.2 - 60.5	<100	90.8	85.7 - 95.9	2,500	85.0	78.8 - 91.1	2,500
Prenatal insurance															
Medi-Cal	26.3	21.8 - 30.8	2,300	63.3	53.9 - 72.6	1,600	53.4	41.0 - 65.8	<100	5.9	* 1.8 - 9.9	200	10.7	5.5 - 15.9	300
Uninsured															
Private	71.7	67.1 - 76.3	6,300	35.3	26.0 - 44.5	900	43.8	31.4 - 56.1	<100	92.3	* 87.5 - 97.1	2,500	86.7	80.9 - 92.4	2,500
Maternal postpartum insurance															
Medi-Cal	17.5	13.6 - 21.4	1,500	38.5	28.9 - 48.2	1,000	43.2	30.3 - 56.1	<100	3.7	* 0.5 - 6.9	100	10.3	5.0 - 15.6	300
Uninsured	7.1	4.4 - 9.8	600	17.9	10.4 - 25.5	500	4.1 *	* 0.9 - 7.2	<100						
Private	72.4	67.8 - 77.0	6,300	34.6	25.4 - 43.9	900	51.8	38.9 - 64.6	<100	92.7	* 88.0 - 97.4	2,500	87.4	81.7 - 93.0	2,500
Infant health insurance															
Medi-Cal	30.3	25.6 - 35.1	2,600	67.4	58.3 - 76.5	1,700	47.6	34.9 - 60.2	<100	7.5	* 2.6 - 12.3	200	18.4	11.5 - 25.2	500
Uninsured	1.5 *	0.3 - 2.6	100												
Private	67.9	63.1 - 72.7	5,900	31.3	22.2 - 40.3	800	48.6	36.3 - 61.0	<100	89.9	84.3 - 95.5	2,400	80.0	73.0 - 87.0	2,300
Maternal Demographics															
Age															
15-19		0.6 - 3.4	200	4.9 *	0.8 - 9.0	100									
20-34	66.6	61.7 - 71.4	5,900	74.6	66.3 - 82.9	1,900	70.6	58.9 - 82.3	<100	61.9	52.5 - 71.3	1,700	63.5	55.3 - 71.6	1,800
35+	31.4	26.6 - 36.2	2,800		12.9 - 28.2	500	24.9	13.7 - 36.1	<100	37.5	28.1 - 47.0	1,000	36.4	28.3 - 44.5	1,100
Did not complete high school (or GED)	10.0	6.9 - 13.0	900	28.9	20.3 - 37.6	700									
Unmarried	25.5	21.0 - 29.9	2,200	58.7	49.3 - 68.2	1,500	63.2	51.7 - 74.6	<100	7.8	* 2.9 - 12.8	200	10.6	5.4 - 15.8	300
Born outside the U.S.	49.4	44.4 - 54.5	4,400	61.9	52.3 - 71.4	1,600	13.4 *	* 3.5 - 23.2	<100	22.5	14.2 - 30.8	600	66.4	58.4 - 74.3	1,900
Speaks non-English language at home	40.7	35.7 - 45.8	3,500	56.5	46.8 - 66.1	1,500	9.6 *	* 0.5 - 18.7	<100	14.7	7.3 - 22.2	400	53.3	44.9 - 61.7	1,500
Lives in a high poverty neighborhood	5.6	3.3 - 7.9	500	13.1	6.7 - 19.5	300									
Income as a percent of the Federal Poverty Guideline															
0-100%	21.8	17.3 - 26.3	1,800	50.7	40.5 - 61.0	1,200	54.5	41.9 - 67.0	<100	7.4	* 2.2 - 12.5	200	9.3	3.8 - 14.7	300
101-200%		10.5 - 18.1	1,200	26.9	17.8 - 36.0	600	6.0 *	* 2.1 - 9.8	<100				14.5	7.8 - 21.2	400
> 200%	63.9	58.7 - 69.0	5,200	22.3	14.2 - 30.5	500	39.6	27.4 - 51.7	<100	90.4	84.8 - 96.0	2,200	76.3	68.4 - 84.1	2,100

<sup>--</sup> Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of San Mateo County was 548. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

<sup>\*</sup> Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

<sup>†</sup> Measure/definition may not be comparable to previous Snapshots (see Technical Notes).



# MIHA Data Snapshot, San Mateo County by Total Live Births, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	San Mateo County				First Liv	e Birth	Second Live Birth or More			
	%	95% CI	N	%	95%	CI	N	%	95% CI	N
Total	100		8,800	45.4	40.4 -	50.5	4,000	54.6	49.5 - 59	.6 4,800
Health Status before Pregnancy										
In good to excellent health Chronic conditions	97.2	95.7 - 98.7	8,600	98.3 *	96.9 -	99.7	3,900	96.3 *	93.8 - 98	4,600
Diabetes†	3.0	1.3 - 4.7	300					4.2 *		5.8 200
Hypertension†	1.4 *	0.4 - 2.3	100					1.5 *		.9 <100
Asthma†	5.2	3.0 - 7.5	500	5.9 *	2.4 -	9.4	200	4.7 *	1.7 - 7	.6 200
Nutrition and Weight										
Daily folic acid use, month before pregnancy	41.8	36.8 - 46.7	3,700	48.0	40.5 -	55.4	1,900	36.7	30.1 - 43	,
Overweight before pregnancy	20.2	16.0 - 24.3	1,700	16.9	11.0 -	22.8	700	23.0	17.0 - 28	,
Obese before pregnancy	15.3	11.5 - 19.1	1,300	7.9	3.7 -	12.1	300	21.7	15.8 - 27	
Inadequate weight gain during pregnancy	18.2	14.0 - 22.3	1,400	13.7	8.4 -	18.9	500	22.2	16.0 - 28	
Excessive weight gain during pregnancy	38.4	33.1 - 43.8	2,900	42.5	34.7 -	50.4	1,500	34.8	27.5 - 42	
Food insecurity during pregnancy	11.1	7.7 - 14.4	1,000	7.0	3.4 -	10.5	300	14.5	9.2 - 19	.8 700
Intimate Partner Violence (IPV) and Depressive Symptoms										
Physical or psychological IPV during pregnancy	4.6	2.5 - 6.7	400	3.4 *	1.3 -	5.6	100	5.6 *		.0 300
Prenatal depressive symptoms	8.5	5.8 - 11.3	700	6.3	3.1 -	9.4	300	10.4	6.1 - 14	.7 500
Postpartum depressive symptoms	10.4	7.3 - 13.5	900	12.8	7.6 -	17.9	500	8.5	4.7 - 12	2 400
Hardships and Support during Pregnancy										
Experienced two or more hardships during childhood	15.9	12.2 - 19.6	1,400	12.4	7.7 -	17.1	500	18.9	13.4 - 24	.4 900
Homeless or did not have a regular place to sleep	3.0 *	1.2 - 4.8	300	2.8 *	0.5 -	5.2	100	3.1 *	0.4 - 5	.7 100
Moved due to problems paying rent or mortgage	5.7	3.1 - 8.3	500	5.3 *	1.9 -	8.7	200	6.0 *	2.2 - 9	.8 300
Woman or partner lost job	9.8	6.7 - 12.9	900	10.3	5.9 -	14.7	400	9.4	5.1 - 13	.7 400
Woman or partner had pay or hours cut back	7.1	4.4 - 9.8	600	6.3	2.7 -	10.0	300	7.7	3.9 - 11	6 400
Became separated or divorced	3.1	1.4 - 4.9	300	4.6 *	1.3 -	7.8	200	2.0 *	0.3 - 3	.6 <100
Had no practical or emotional support	5.7	3.1 - 8.2	500	2.3 *	0.1 -	4.6	<100	8.5	4.3 - 12	7 400
Substance Use										
Any smoking, 3 months before pregnancy	8.6	5.8 - 11.5	800	7.2	3.6 -	10.8	300	9.8	5.5 - 14	.1 500
Any smoking, 3rd trimester†	1.1 *	0.3 - 2.0	<100							
Any smoking, postpartum	2.7	1.2 - 4.3	200	2.1 *	0.4 -	3.9	<100	3.3 *	0.8 - 5	.8 200
Any binge drinking, 3 months before pregnancy	13.6	10.1 - 17.1	1,200	15.6	10.3 -	21.0	600	11.9	7.3 - 16	600
Any alcohol use, 3rd trimester†	11.3	8.1 - 14.5	1,000	10.4	5.9 -	14.9	400	12.1	7.5 - 16	6.6 600
Pregnancy Intention and Family Planning										
Mistimed or unwanted pregnancy	22.7	18.4 - 27.0	2,000	21.9	15.8 -	28.1	900	23.4	17.3 - 29	.4 1,100
Unsure of pregnancy intentions	7.8	5.1 - 10.6	700	8.2	3.9 -	12.5	300	7.6	4.1 - 11	.0 400
Postpartum birth control use†	79.1	74.9 - 83.2	6,800	75.7	69.2 -	82.3	2,900	81.8	76.6 - 87	3,800
Infant Sleep and Breastfeeding										
Placed infant on back to sleep	88.6	85.3 - 91.8	7,700	91.1	87.0 -	95.2	3,600	86.4	81.6 - 91	3 4,100
Infant always or often shared bed	30.1	25.4 - 34.8	2,600	27.9	21.0 -	34.7	1,100	32.0	25.4 - 38	1,500
Intended to breastfeed, before birth	92.9	90.1 - 95.7	8,100	95.3 *	91.7 -	98.9	3,800	90.9	86.8 - 95	4,300
Intended to breastfeed exclusively, before birth	64.7	59.8 - 69.6	5,600	77.8	71.7 -	83.9	3,100	53.8	46.8 - 60	2,500
Breastfeeding practices										
Any breastfeeding, 1 month after delivery	89.7	86.6 - 92.8	7,700	94.2	90.9 -	97.4	3,700	86.0	81.1 - 90	4,000
Exclusive breastfeeding, 1 month after delivery	44.6	39.4 - 49.7	3,800	49.0	41.4 -	56.6	1,900	40.8	33.9 - 47	1,900
Any breastfeeding, 3 months after delivery	78.2	73.3 - 83.0	5,000	83.0	76.9 -	89.2	2,300	74.3	67.2 - 81	4 2,600
Exclusive breastfeeding, 3 months after delivery	32.8	27.1 - 38.5	2,100	35.8	27.2 -	44.5	1,000	30.4	23.0 - 37	.9 1,100



# MIHA Data Snapshot, San Mateo County by Total Live Births, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		San Mateo Count	y		First Live Birth		Second Live Birth or More			
	%	95% CI	N	%	95% CI	N	%	95% CI	N	
Service Utilization and Health Insurance Coverage										
Had a usual source of pre-pregnancy care	77.7	73.5 - 81.9	6,800	72.9	66.3 - 79.6	2,900	81.6	76.3 - 86.9	3,900	
Initiated prenatal care in 1st trimester	92.4	89.6 - 95.1	7,900	93.1	89.5 - 96.8	3,700	91.7	87.7 - 95.7	4,300	
Mom or infant needed but couldn't afford care postpartum	5.7	3.6 - 7.8	500	4.3	1.8 - 6.8	200	6.9	3.7 - 10.1	300	
Participated in WIC during pregnancy†	30.5	25.7 - 35.4	2,600	19.3	13.6 - 25.0	700	39.7	32.7 - 46.7	1,800	
Received CalFresh (food stamps) during pregnancy	9.1	6.3 - 12.0	800	3.7 *	1.3 - 6.1	100	13.6	8.8 - 18.3	700	
Pre-pregnancy insurance										
Medi-Cal	15.2	11.5 - 18.9	1,300	7.2	3.5 - 11.0	300	21.9	16.1 - 27.7	1,100	
Uninsured	9.6	6.5 - 12.6	800	10.9	6.3 - 15.6	400	8.4	4.4 - 12.5	400	
Private	72.0	67.4 - 76.5	6,300	79.0	73.2 - 84.9	3,200	66.1	59.5 - 72.7	3,200	
Prenatal insurance										
Medi-Cal	26.3	21.8 - 30.8	2,300	18.1	12.6 - 23.5	700	33.1	26.5 - 39.7	1,600	
Uninsured										
Private	71.7	67.1 - 76.3	6,300	79.0	73.1 - 84.9	3,200	65.6	58.9 - 72.3	3,200	
Maternal postpartum insurance										
Medi-Cal	17.5	13.6 - 21.4	1,500	12.2	7.4 - 17.0	500	21.9	16.1 - 27.8	1,000	
Uninsured	7.1	4.4 - 9.8	600	7.0	3.3 - 10.6	300	7.2	3.3 - 11.1	300	
Private	72.4	67.8 - 77.0	6,300	79.8	74.0 - 85.6	3,200	66.3	59.6 - 73.0	3,200	
Infant health insurance										
Medi-Cal	30.3	25.6 - 35.1	2,600	23.1	17.0 - 29.3	900	36.3	29.5 - 43.1	1,700	
Uninsured	1.5 *	0.3 - 2.6	100							
Private	67.9	63.1 - 72.7	5,900	74.6	68.2 - 81.0	2,900	62.3	55.5 - 69.1	2,900	
Maternal Demographics										
Age										
15-19	2.0 *	0.6 - 3.4	200	4.2 *	1.3 - 7.2	200				
20-34	66.6	61.7 - 71.4	5,900	76.6	70.0 - 83.3	3,100	58.2	51.4 - 65.0	2,800	
35+	31.4	26.6 - 36.2	2,800	19.1	12.8 - 25.5	800	41.6	34.8 - 48.4	2,000	
Did not complete high school (or GED)	10.0	6.9 - 13.0	900	6.7	2.9 - 10.6	300	12.7	8.1 - 17.2	600	
Unmarried	25.5	21.0 - 29.9	2,200	23.9	17.7 - 30.1	900	26.8	20.6 - 33.1	1,300	
Race/Ethnicity										
White	32.7	27.8 - 37.6	2,800	34.9	27.4 - 42.4	1,300	30.9	24.4 - 37.5	1,500	
Hispanic	31.0	26.1 - 35.9	2,600	24.4	17.6 - 31.2	900	36.2	29.4 - 43.0	1,700	
Asian/Pacific Islander	34.5	29.7 - 39.3	2,900	39.0	31.6 - 46.4	1,500	30.9	24.7 - 37.2	1,500	
Black	1.6	1.3 - 1.9	100	1.3	0.9 - 1.8	<100	1.8	1.3 - 2.3	<100	
Born outside the U.S.	49.4	44.4 - 54.5	4,400	43.4	36.0 - 50.8	1,700	54.5	47.6 - 61.3	2,600	
Speaks non-English language at home	40.7	35.7 - 45.8	3,500	38.5	31.2 - 45.9	1,500	42.5	35.7 - 49.4	2,000	
Lives in a high poverty neighborhood	5.6	3.3 - 7.9	500	5.5 *	2.2 - 8.8	200	5.6	2.5 - 8.8	300	
Income as a percent of the Federal Poverty Guideline										
0-100%	21.8	17.3 - 26.3	1,800	16.9	11.2 - 22.6	600	26.0	19.4 - 32.6	1,200	
101-200%	14.3	10.5 - 18.1	1,200	9.1	4.4 - 13.9	300	18.6	12.9 - 24.3	800	
> 200%	63.9	58.7 - 69.0	5,200	74.0	67.1 - 80.9	2,700	55.4	48.3 - 62.6	2,500	

<sup>--</sup> Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of San Mateo County was 548. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

<sup>\*</sup> Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%)

<sup>†</sup> Measure/definition may not be comparable to previous Snapshots (see Technical Notes).