

MIHA Data Snapshot, Sonoma County by Education, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Sonon	na Count	y		<high sch<="" th=""><th>lool</th><th>High</th><th>School</th><th>Graduat</th><th>e or GED</th><th></th><th>Some College</th><th>e</th><th> Co</th><th>ollege Gradu</th><th>ate</th></high>	lool	High	School	Graduat	e or GED		Some College	e	Co	ollege Gradu	ate
	% 95	% CI	Ν	%	95% CI	I N	%	9	5% CI	Ν	%	95% CI	N	%	95% CI	Ν
Total	100		4,900	14.9	11.6 - 18	3.3 70	0 19	9.0 15.3	3 - 22.8	900	34.9	30.2 - 39.5	1,700	31.1	26.8 - 35.5	1,500
Health Status before Pregnancy																
In good to excellent health	93.5 91.2	- 95.8	4,600	75.6	64.7 - 86	5.5 50	o 94	1.2 * 89.9	9 - 98.4	900	95.2	* 92.1 - 98.4	1,600			
Chronic conditions																
Diabetes ⁺	2.2 1.0	- 3.4	100	5.7 *	1.3 - 10).1 <10	0				2.9	* 0.4 - 5.3	<100			
Hypertension ⁺	3.3 1.9	- 4.8	200								5.4	* 2.1 - 8.8	<100			
Asthma ⁺	10.7 7.9	- 13.5	500	15.9	6.9 - 24	1.9 10	0 9).3 * 3.4	4 - 15.2	<100	8.7	4.3 - 13.2	100	11.9	6.9 - 16.9	200
Nutrition and Weight																
Daily folic acid use, month before pregnancy	38.7 34.0	- 43.4	1,900	25.1	14.0 - 36	5.2 20	o 31	L.2 20.5	5 - 41.8	300	41.8	33.4 - 50.2	700	48.9	40.7 - 57.1	700
Overweight before pregnancy	24.0 19.9	- 28.1	1,100	42.4	28.8 - 56	5.0 20	D 26	5.0 16.7	7 - 35.3	200	23.5	16.5 - 30.6	400	17.0	11.1 - 23.0	300
Obese before pregnancy	21.7 17.5	- 25.9	1,000	25.4	14.0 - 36	5.8 10	0 25	5.9 14.5	5 - 37.4	200	27.1	19.1 - 35.1	400	13.1	8.1 - 18.1	200
Inadequate weight gain during pregnancy	14.6 11.1	- 18.0	600	21.4 *	8.6 - 34	1.3 <10	o 2 3	3.4 13.9	9 - 32.9	200	11.3	6.0 - 16.6	200	10.8	6.2 - 15.4	200
Excessive weight gain during pregnancy	48.5 43.3	- 53.7	2,000	51.3	36.3 - 66	5.3 20	0 40).2 28.5	5 - 51.8	300	53.6	44.4 - 62.8	800	47.9	39.5 - 56.2	700
Food insecurity during pregnancy	14.7 11.2	- 18.2	700	29.1	18.0 - 40).2 20	0 19	9.0 10.5	5 - 27.6	200	14.9	8.6 - 21.2	300	2.0	* 0.1 - 3.8	<100
Intimate Partner Violence (IPV) and Depressive Symptoms																
Physical or psychological IPV during pregnancy	6.0 3.5	- 8.6	300	14.3 *	5.4 - 23	8.3 10	D 5	5.8 * 1.5	5 - 10.1	<100	5.2	* 0.8 - 9.6	<100			
Prenatal depressive symptoms	14.1 10.8	- 17.5	700	28.9	17.7 - 40	0.1 20	D 11	L.2 5.0) - 17.5	100	12.1	6.5 - 17.7	200	8.6	4.3 - 12.8	100
Postpartum depressive symptoms	13.7 10.5	- 16.9	700	23.3	12.8 - 33	8.9 20	0 9	9.9* 4.0) - 15.8	<100	12.1	7.4 - 16.8	200	11.2	6.0 - 16.5	200
Hardships and Support during Pregnancy																
Experienced two or more hardships during childhood	27.2 23.0	- 31.4	1,300	40.1	27.8 - 52	2.5 30	0 28	3.2 18.3	3 - 38.1	300	29.0	21.8 - 36.2	500	17.7	11.9 - 23.5	300
Homeless or did not have a regular place to sleep	3.7 1.6	- 5.7	200	7.2 *	1.5 - 12	2.9 <10	0				5.0	* 0.5 - 9.6	<100			
Moved due to problems paying rent or mortgage	6.5 4.0	- 8.9	300	7.2 *	1.0 - 13	3.4 <10	0 5	5.5 * 0.4	4 - 10.6	<100	7.3	* 2.5 - 12.1	100	5.8	* 1.7 - 9.9	<100
Woman or partner lost job	13.0 9.6	- 16.4	600	19.5	9.6 - 29	9.4 10	D 13	3.8 * 5.7	7 - 22.0	100	14.4	8.6 - 20.2	200	5.2	* 1.3 - 9.1	<100
Woman or partner had pay or hours cut back	14.4 10.7	- 18.0	700	17.0	7.2 - 26	5.8 10	0 10). 7 * 3.3	3 - 18.1	<100	17.8	11.1 - 24.6	300	9.9	4.6 - 15.1	100
Became separated or divorced	7.1 4.4	- 9.7	300	13.5 *	4.4 - 22	2.6 <10	0 7	7.3* 2.3	3 - 12.4	<100	7.0	* 2.3 - 11.7	100	1.6	* 0.1 - 3.1	<100
Had no practical or emotional support	4.5 1.9	- 7.0	200	9.2 *	0.7 - 17	7.7 <10	0									
Substance Use																
Any smoking, 3 months before pregnancy	13.7 10.4	- 17.1	700	20.2	10.0 - 30).4 10	D 7	7.2 * 2.9	9 - 11.5	<100	17.8	11.2 - 24.3	300	9.0	4.5 - 13.5	100
Any smoking, 3rd trimester ⁺	3.1 * 1.2	- 5.0	200	9.5 *	1.6 - 17	7.3 <10	0									
Any smoking, postpartum	5.3 3.0	- 7.5	300	11.9 *	3.9 - 19	9.8 <10	0				8.0	* 3.0 - 13.1	100			
Any binge drinking, 3 months before pregnancy	18.4 14.6	- 22.2	900	10.1 *	3.2 - 17	/.0 <10	0 14	1.1 6.5	5 - 21.7	100	19.6	12.8 - 26.4	300	22.8	15.7 - 30.0	300
Any alcohol use, 3rd trimester ⁺	10.0 7.3	- 12.7	500								5.7	* 2.1 - 9.4	<100	22.4	15.7 - 29.0	300
Pregnancy Intention and Family Planning																
Mistimed or unwanted pregnancy	26.1 21.9	- 30.4	1,300	35.1	23.0 - 47	7.2 20	0 18	3.6 11.0) - 26.2	200	34.4	26.2 - 42.6	600	16.7	10.5 - 22.9	300
Unsure of pregnancy intentions	11.0 8.3	- 13.8	500	11.3 *	4.5 - 18	3.1 <10	0 16	5.0 8.7	7 - 23.4	100	9.0	5.0 - 13.0	100	9.1	4.0 - 14.2	100
Postpartum birth control use ⁺	87.1 84.0	- 90.2	4,300	85.9*	75.9 - 95	5.9 60	D 88	3.7 82.5	5 - 94.8	800	87.1	81.9 - 92.4	1,400	87.1	81.6 - 92.7	1,300
Infant Sleep and Breastfeeding																
Placed infant on back to sleep	83.7 80.1	- 87.3	4,000	73.8	62.0 - 85	5.5 50	0 81	L.O 72.4	4 - 89.6	700	88.1	83.0 - 93.3	1,500	86.8	81.8 - 91.9	1,300
Infant always or often shared bed	39.7 35.0	- 44.4	1,900	51.2	38.8 - 63	3.6 30	0 42	2.3 31.5	5 - 53.2	400	37.7	29.7 - 45.8	600	34.3	26.5 - 42.0	500
Intended to breastfeed, before birth	95.4 93.5	- 97.4	4,600	92.1 *	84.9 - 99	9.2 60	o 93	3.3 * 87.8	3 - 98.9	800	96.6	* 93.9 - 99.3	1,600	96.6	* 93.6 - 99.6	1,500
Intended to breastfeed exclusively, before birth	81.0 77.3	- 84.7	3,900	67.9	56.3 - 79	9.5 50	66	5.5 56.3	3 - 76.8	600	87.5	82.5 - 92.4	1,500	87.3	81.5 - 93.0	1,300
Breastfeeding practices																
Any breastfeeding, 1 month after delivery	92.3 89.9	- 94.7	4,500	86.3 *	78.0 - 94	1.6 60	o 91	L .3 * 85.5	5 - 97.2	800	92.7	88.9 - 96.5	1,600	94.9	* 90.8 - 99.0	1,400
Exclusive breastfeeding, 1 month after delivery	63.5 58.9		3,000	52.2	39.5 - 64	1.9 30	0 60).2 49.3	3 - 71.1	500	62.6	54.5 - 70.6	1,000	70.1	62.2 - 78.0	1,000
Any breastfeeding, 3 months after delivery	82.8 78.6	- 87.1	3,000	78.3	67.1 - 89	9.5 50	o 7 4	1.9 63.5	5 - 86.3	500	79.1	71.2 - 87.1	1,000	93.2	* 88.8 - 97.7	1,000
Exclusive breastfeeding, 3 months after delivery	49.0 43.4	- 54.6	1,800	38.2	24.1 - 52	2.2 20	<u>44</u>	1.2 31.6	5 - 56.7	300	<u>47.7</u>	37.8 - 57.5	600	56.0	46.1 - 65.9	600



MIHA Data Snapshot, Sonoma County by Education, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Sonoma Co	ounty	<high schoo<="" th=""><th>1</th><th>High School Graduate</th><th>e or GED</th><th>Son</th><th>ne College</th><th>2</th><th>College Grad</th><th>uate</th></high>	1	High School Graduate	e or GED	Son	ne College	2	College Grad	uate
	% 95% C	N	% 95% CI	Ν	% 95% CI	N	% 9	95% CI	N	% 95% CI	N
Service Utilization and Health Insurance Coverage											
Had a usual source of pre-pregnancy care	80.9 77.1 - 84	.6 4,000	85.0 76.6 - 93.3	600	81.0 73.2 - 88.8	700	75.8 68	.6 - 82.9	1,300	88.0 82.9 - 93.1	1,300
Initiated prenatal care in 1st trimester	92.3 89.9 - 94	.7 4,500	79.6 68.7 - 90.4	500	89.0 82.7 - 95.2	800	94.4 91	.2 - 97.6	1,600	97.3 * 95.2 - 99.4	1,500
Mom or infant needed but couldn't afford care postpartum	8.5 5.6 - 12	.5 400	13.3 * 4.4 - 22.3	<100	5.1 [*] 1.1 - 9.0	<100	10.4 4	.6 - 16.3	200	4.0 * 1.4 - 6.5	5 <100
Participated in WIC during pregnancy ⁺	41.9 37.2 - 46	.6 2,100	81.6 71.5 - 91.8	600	63.9 53.5 - 74.3	600	39.4 31	.3 - 47.6	700	10.0 5.0 - 14.9	200
Received CalFresh (food stamps) during pregnancy	19.3 15.6 - 23	.0 1,000	48.6 36.5 - 60.8	400	23.0 14.1 - 31.8	200	16.8 10	.9 - 22.8	300	3.7 * 1.1 - 6.4	× 100
Pre-pregnancy insurance											
Medi-Cal	17.4 14.0 - 20	.9 900	43.5 31.5 - 55.4	300	29.5 19.4 - 39.6	300	11.7 6	.8 - 16.5	200	4.7 * 0.6 - 8.7	<100
Uninsured	24.4 20.4 - 28	.5 1,200	40.2 28.1 - 52.3	300	38.1 27.5 - 48.6	300	23.0 16	.4 - 29.6	400	7.2 3.5 - 10.8	3 100
Private	55.3 50.6 - 60	.0 2,700	13.4 * 5.2 - 21.5	<100	27.9 18.1 - 37.7	300	61.2 53	.1 - 69.2	1,000	87.8 82.6 - 93.1	1,300
Prenatal insurance											
Medi-Cal	47.1 42.4 - 52	.9 2,300	91.3 * 85.0 - 97.7	700	76.1 67.3 - 84.9	700	41.5 33	.3 - 49.8	700	12.3 7.0 - 17.6	5 200
Uninsured	1.1 * 0.3 - 2	.8 <100									
Private	48.5 43.8 - 53	.3 2,400	8.0 * 1.8 - 14.3	<100	18.1 10.3 - 25.8	200	52.5 44	.0 - 60.9	900	84.1 78.3 - 89.8	3 1,300
Maternal postpartum insurance											
Medi-Cal	30.6 26.1 - 3	.0 1,500	58.2 46.2 - 70.3	400	46.7 35.8 - 57.6	400	31.0 23	.1 - 38.9	500	5.9 * 1.9 - 9.9	9 <100
Uninsured	15.8 12.5 - 19	.1 800	36.1 24.4 - 47.9	300	27.6 17.7 - 37.4	300	9.5 5	.4 - 13.7	200	5.1 2.1 - 8.0) <100
Private	51.0 46.2 - 55	.8 2,500			22.9 13.9 - 31.8	200	54.4 45	.9 - 62.8	900	87.9 82.9 - 92.8	3 1,300
Infant health insurance											
Medi-Cal	48.2 43.5 - 53	.0 2,300	90.0 * 81.8 - 98.2	600	78.1 69.6 - 86.6	700	43.6 35	.3 - 52.0	700	14.5 8.9 - 20.0	200
Uninsured	1.1 * 0.2 - 2	.9 <100									
Private	47.1 42.3 - 52	.9 2,300			17.6 10.0 - 25.3	200	49.3 40	.8 - 57.8	800	83.8 78.1 - 89.6	5 1,300
Maternal Demographics											
Unmarried	33.2 28.9 - 3	.5 1,600	57.8 45.9 - 69.7	400	40.6 30.3 - 50.8	400	35.7 27	.9 - 43.5	600	13.0 8.1 - 18.0) 200
Race/Ethnicity											
White	51.7 46.8 - 56	.5 2,300	12.6 * 3.4 - 21.7	<100	25.4 16.0 - 34.8	200	60.2 51	.9 - 68.5	900	78.7 71.3 - 86.1	1,100
Hispanic	40.8 36.0 - 4	.5 1,800	79.5 68.8 - 90.1	500	68.6 58.6 - 78.7	600	32.9 25	.0 - 40.7	500	11.8 7.0 - 16.6	5 200
Asian/Pacific Islander	5.1 2.4 -	.7 200								8.2 * 2.0 - 14.5	5 100
Black	1.4 0.6 - 2	.2 <100									
Born outside the U.S.	30.6 26.2 - 35	.0 1,500	63.7 51.8 - 75.5	500	44.0 33.1 - 54.9	400	20.4 13	.8 - 27.1	300	16.8 9.5 - 24.0	300
Speaks non-English language at home	24.8 20.8 - 28	.8 1,200	59.9 47.9 - 71.9	400	41.8 31.2 - 52.4	400	14.8 8	.8 - 20.9	300	9.0 3.9 - 14.1	100
Lives in a high poverty neighborhood	17.0 13.6 - 20	.3 800	25.3 14.8 - 35.8	200	24.0 15.2 - 32.8	200	15.1 9	.7 - 20.6	200	11.8 6.7 - 16.9	200
Income as a percent of the Federal Poverty Guideline											
0-100%	27.2 22.8 - 32	.5 1,300	75.3 64.2 - 86.4	500	39.8 28.5 - 51.2	300	24.4 17	.0 - 31.8	400	3.9 * 1.3 - 6.5	5 <100
101-200%	24.0 19.9 - 28	.2 1,100	20.3 10.2 - 30.4	100	42.2 30.9 - 53.5	400	26.4 19	.2 - 33.6	400	10.4 5.3 - 15.6	5 200
> 200%	48.8 44.0 - 53	.7 2,300			18.0 9.4 - 26.6	200	49.2 40	.7 - 57.8	800	85.7 80.1 - 91.4	1,300

-- Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

* Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

+ Measure/definition may not be comparable to previous Snapshots (see Technical Notes).

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Sonoma County was 557. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.



MIHA Data Snapshot, Sonoma County by Geographical Area, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		Sonoma County	,			Urban			Rural/Frontier	
	%	95% CI	N	-	%	95% CI	N	%	95% CI	Ν
Total	100		4,900		85.5	82.1 - 88.9	4,100	14.5	11.1 - 17.9	700
Health Status before Pregnancy										
In good to excellent health	93.5	91.2 - 95.8	4,600		94.4	92.1 - 96.7	3,900	91.8 *	85.4 - 98.2	600
Chronic conditions										
Diabetes ⁺	2.2	1.0 - 3.4	100		2.3	1.0 - 3.6	<100			
Hypertension ⁺	3.3	1.9 - 4.8	200		2.5	1.2 - 3.8	<100			
Asthma ⁺	10.7	7.9 - 13.5	500		11.1	8.0 - 14.2	400	7.8 *	1.6 - 14.0	<100
Nutrition and Weight										
Daily folic acid use, month before pregnancy	38.7	34.0 - 43.4	1,900		40.6	35.4 - 45.8	1,700	28.4	17.6 - 39.2	200
Overweight before pregnancy	24.0	19.9 - 28.1	1,100		22.9	18.7 - 27.2	900	27.3	14.1 - 40.4	200
Obese before pregnancy	21.7	17.5 - 25.9	1,000		21.6	17.0 - 26.2	800	25.2	12.9 - 37.6	200
Inadequate weight gain during pregnancy	14.6	11.1 - 18.0	600		15.8	11.9 - 19.6	500	9.2 *	0.7 - 17.7	<100
Excessive weight gain during pregnancy	48.5	43.3 - 53.7	2,000		45.9	40.4 - 51.5	1,600	63.9	50.0 - 77.8	400
Food insecurity during pregnancy	14.7	11.2 - 18.2	700		14.1	10.3 - 17.9	600	17.3	7.7 - 26.9	100
Intimate Partner Violence (IPV) and Depressive Symptoms										
Physical or psychological IPV during pregnancy	6.0	3.5 - 8.6	300		5.5	2.6 - 8.3	200	7.3 *		<100
Prenatal depressive symptoms	14.1	10.8 - 17.5	700		15.4	11.5 - 19.2	600	4.7 *	0.1 - 9.2	<100
Postpartum depressive symptoms	13.7	10.5 - 16.9	700		14.2	10.6 - 17.8	600	6.8 *	1.6 - 12.0	<100
Hardships and Support during Pregnancy										
Experienced two or more hardships during childhood	27.2	23.0 - 31.4	1,300		26.6	22.0 - 31.2	1,100	25.4	15.0 - 35.9	200
Homeless or did not have a regular place to sleep	3.7	1.6 - 5.7	200		3.4 *	1.1 - 5.7	100			
Moved due to problems paying rent or mortgage	6.5	4.0 - 8.9	300		6.2	3.4 - 9.0	300	8.1 *	2.3 - 13.9	<100
Woman or partner lost job	13.0	9.6 - 16.4	600		12.5	8.8 - 16.1	500	18.1	7.6 - 28.5	100
Woman or partner had pay or hours cut back	14.4	10.7 - 18.0	700		14.6	10.5 - 18.7	600	14.3	5.9 - 22.7	<100
Became separated or divorced	7.1	4.4 - 9.7	300		5.8	3.1 - 8.5	200	11.2 *	2.9 - 19.4	<10
Had no practical or emotional support	4.5	1.9 - 7.0	200		4.4 *	1.6 - 7.2	200			
Substance Use										
Any smoking, 3 months before pregnancy	13.7	10.4 - 17.1	700		14.3	10.5 - 18.1	600	5.1 *	0.3 - 9.8	<100
Any smoking, 3rd trimester ⁺	3.1 *	1.2 - 5.0	200		2.6 *	0.7 - 4.6	100			
Any smoking, postpartum	5.3	3.0 - 7.5	300		5.3	2.8 - 7.8	200			
Any binge drinking, 3 months before pregnancy	18.4	14.6 - 22.2	900		18.9	14.7 - 23.1	800	17.9	7.4 - 28.3	100
Any alcohol use, 3rd trimester†	10.0	7.3 - 12.7	500		9.9	6.9 - 12.9	400	10.8 *	4.2 - 17.4	<100
Pregnancy Intention and Family Planning										
Mistimed or unwanted pregnancy	26.1	21.9 - 30.4	1,300		24.3	19.8 - 28.9	1,000	36.8	24.3 - 49.3	300
Unsure of pregnancy intentions	11.0	8.3 - 13.8	500		11.4	8.4 - 14.5	500	9.6 *		<10
Postpartum birth control use [†]	87.1	84.0 - 90.2	4,300		87.7	84.3 - 91.0	3,600	85.7 *	76.7 - 94.8	600
Infant Sleep and Breastfeeding										
Placed infant on back to sleep	83.7	80.1 - 87.3	4,000		83.5	79.5 - 87.4	3,400	85.7 *		600
Infant always or often shared bed	39.7	35.0 - 44.4	1,900		38.8	33.7 - 43.9	1,600	46.5	33.6 - 59.5	300
Intended to breastfeed, before birth	95.4	93.5 - 97.4	4,600		95.0	92.8 - 97.2	3,900			
Intended to breastfeed exclusively, before birth	81.0	77.3 - 84.7	3,900		79.8	75.7 - 83.9	3,200	85.1 *	76.0 - 94.2	600
Breastfeeding practices	02.2		4 500		01 7	00.0	2 700			
Any breastfeeding, 1 month after delivery	92.3	89.9 - 94.7	4,500		91.7	88.9 - 94.5	3,700		F0 7 00 0	
Exclusive breastfeeding, 1 month after delivery	63.5	58.9 - 68.2	3,000		62.4	57.4 - 67.5	2,500	70.4	58.7 - 82.0	500
Any breastfeeding, 3 months after delivery	82.8	78.6 - 87.1	3,000		82.5	77.8 - 87.2	2,500	82.1	71.8 - 92.4	500
Exclusive breastfeeding, 3 months after delivery	49.0	43.4 - 54.6	1,800		49.4	43.2 - 55.6	1,500	46.9	33.2 - 60.6	300



MIHA Data Snapshot, Sonoma County by Geographical Area, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		Sonoma County			Urban			Rural/Frontier	
	%	95% CI	N	%	95% CI	N	%	95% CI	N
Service Utilization and Health Insurance Coverage									
Had a usual source of pre-pregnancy care	80.9	77.1 - 84.6	4,000	82.5	78.5 - 86.5	3,400	72.1	60.9 - 83.4	500
Initiated prenatal care in 1st trimester	92.3	89.9 - 94.7	4,500	92.2	89.6 - 94.9	3,700			
Mom or infant needed but couldn't afford care postpartum	8.5	5.6 - 11.5	400	8.8	5.5 - 12.2	400	7.3 *	1.3 - 13.2	<100
Participated in WIC during pregnancy ⁺	41.9	37.2 - 46.6	2,100	41.2	36.1 - 46.3	1,700	49.9	37.0 - 62.9	300
Received CalFresh (food stamps) during pregnancy	19.3	15.6 - 23.0	1,000	20.0	15.8 - 24.1	800	12.5	5.2 - 19.8	<100
Pre-pregnancy insurance									
Medi-Cal	17.4	14.0 - 20.9	900	17.2	13.4 - 21.0	700	14.9	6.4 - 23.4	100
Uninsured	24.4	20.4 - 28.5	1,200	23.1	18.8 - 27.4	900	33.4	21.5 - 45.3	200
Private	55.3	50.6 - 60.0	2,700	57.6	52.5 - 62.6	2,400	44.4	31.6 - 57.2	300
Prenatal insurance									
Medi-Cal	47.1	42.4 - 51.9	2,300	46.4	41.2 - 51.5	1,900	49.7	36.8 - 62.6	300
Uninsured	1.1 *	0.3 - 1.8	<100	1.0 *	0.2 - 1.8	<100			
Private	48.5	43.8 - 53.3	2,400	50.8	45.6 - 56.0	2,100	36.2	24.1 - 48.4	300
Maternal postpartum insurance									
Medi-Cal	30.6	26.1 - 35.0	1,500	30.6	25.8 - 35.5	1,200	27.3	16.2 - 38.4	200
Uninsured	15.8	12.5 - 19.1	800	15.5	11.9 - 19.2	600	18.4	8.7 - 28.1	100
Private	51.0	46.2 - 55.8	2,500	52.5	47.4 - 57.7	2,100	43.1	30.4 - 55.9	300
Infant health insurance									
Medi-Cal	48.2	43.5 - 53.0	2,300	47.0	41.8 - 52.2	1,900	56.8	44.0 - 69.7	400
Uninsured	1.1 *	0.2 - 1.9	<100	1.3 *	0.3 - 2.3	<100			
Private	47.1	42.3 - 51.9	2,300	49.2	44.0 - 54.4	2,000	32.5	20.8 - 44.1	200
Maternal Demographics									
Age									
15-19	3.8	2.1 - 5.5	200	4.1	2.2 - 6.0	200			
20-34	74.0	69.5 - 78.5	3,700	76.1	71.4 - 80.9	3,100	64.8	51.7 - 77.8	400
35+	22.2	17.8 - 26.5	1,100	19.8	15.2 - 24.4	800	32.1	19.3 - 44.9	200
Did not complete high school (or GED)	14.9	11.6 - 18.3	700	14.6	11.0 - 18.2	600	12.7 *	3.7 - 21.7	<100
Unmarried	33.2	28.9 - 37.5	1,600	32.7	28.1 - 37.4	1,300	33.6	21.9 - 45.4	200
Race/Ethnicity									
White	51.7	46.8 - 56.5	2,300	51.3	46.0 - 56.5	1,900	53.1	39.6 - 66.6	300
Hispanic	40.8	36.0 - 45.5	1,800	40.9	35.8 - 46.0	1,500	45.2	31.7 - 58.7	300
Asian/Pacific Islander	5.1	2.4 - 7.7	200	5.3	2.5 - 8.0	200			
Black	1.4	0.6 - 2.2	<100	1.4 *	0.4 - 2.3	<100			
Born outside the U.S.	30.6	26.2 - 35.0	1,500	32.0	27.1 - 36.8	1,300	23.2	12.5 - 33.9	200
Speaks non-English language at home	24.8	20.8 - 28.8	1,200	25.4	20.9 - 29.8	1,000	25.8	14.2 - 37.4	200
Lives in a high poverty neighborhood	17.0	13.6 - 20.3	800	19.0	15.2 - 22.8	800			
Income as a percent of the Federal Poverty Guideline									
0-100%	27.2	22.8 - 31.5	1,300	27.0	22.2 - 31.8	1,100	27.1	15.9 - 38.3	200
101-200%	24.0	19.9 - 28.2	1,100	23.4	18.9 - 28.0	900	29.1	17.0 - 41.3	200
> 200%	48.8	44.0 - 53.7	2,300	49.5	44.2 - 54.8	1,900	43.8	30.6 - 57.1	300

-- Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

* Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Sonoma County was 557. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. Urban and Rural/Frontier designations are based on the population size or densities of Medical Service Study Areas. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.



MIHA Data Snapshot, Sonoma County by Income, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Sonoma Count	Y	0-100% Federal Poverty	Guideline	101-200% Federal Poverty	Guideline	> 200% Federal Poverty	Guideline
	% 95% CI	Ν	% 95% CI	Ν	% 95% CI	N	% 95% CI	Ν
Total	100	4,900	27.2 22.8 - 31.5	1,300	24.0 19.9 - 28.2	1,100	48.8 44.0 - 53.7	2,300
Health Status before Pregnancy								
In good to excellent health	93.5 91.2 - 95.8	4,600	83.9 77.2 - 90.7	1,100	93.6 * 88.9 - 98.2	1,100		
Chronic conditions								
Diabetes ⁺	2.2 1.0 - 3.4	100	4.3 * 1.3 - 7.3	<100				
Hypertension [†]	3.3 1.9 - 4.8	200	2.9 * 0.6 - 5.1	<100			3.3 * 1.0 - 5.5	<100
Asthma†	10.7 7.9 - 13.5	500	14.7 8.0 - 21.3	200	7.5 * 3.1 - 12.0	<100	10.3 6.3 - 14.3	200
Nutrition and Weight					A - - - - - - - - - -			
Daily folic acid use, month before pregnancy	38.7 34.0 - 43.4	1,900	29.0 20.3 - 37.8	400	27.7 19.0 - 36.4	300	51.1 44.1 - 58.1	1,200
Overweight before pregnancy	24.0 19.9 - 28.1	1,100	32.5 23.1 - 41.8	400	23.1 15.1 - 31.2	200	20.4 14.9 - 25.9	500
Obese before pregnancy Inadequate weight gain during pregnancy	21.7 17.5 - 25.9	1,000 600	25.4 16.4 - 34.4	300 200	22.4 14.2 - 30.7	200	20.7 14.5 - 26.8 7.8 4.6 - 11.0	500 200
Excessive weight gain during pregnancy	14.6 11.1 - 18.0 48.5 43.3 - 53.7	2,000	23.4 14.3 - 32.4 40.4 29.8 - 51.0	400	18.2 10.1 - 26.3 53.1 42.1 - 64.0	200 500	7.8 4.6 - 11.0 50.2 42.9 - 57.5	1,100
Food insecurity during pregnancy	14.7 11.2 - 18.2	2,000	32.3 23.3 - 41.3	400	16.4 8.8 - 24.1	200	2.0 * 0.4 - 3.6	<100
Intimate Partner Violence (IPV) and Depressive Symptoms	14.7 11.2 10.2	700	52.5 25.5 41.5	400	10.4 0.0 24.1	200	2.0 0.4 5.0	<100
Physical or psychological IPV during pregnancy	6.0 3.5 - 8.6	300	13.1 5.8 - 20.3	200	8.4 * 2.1 - 14.7	<100	1.4 * 0.2 - 2.6	<100
Prenatal depressive symptoms	14.1 10.8 - 17.5	300 700	13.1 5.8 - 20.3 23.1 15.0 - 31.2	300	14.9 7.3 - 22.5	200	1.4 0.2 - 2.6 9.0 5.2 - 12.7	200
Postpartum depressive symptoms	14.1 10.8 - 17.3 13.7 10.5 - 16.9	700	19.5 12.4 - 26.6	200	14.9 7.3 - 22.3 16.9 9.0 - 24.7	200	9.0 5.2 - 12.7 9.5 5.6 - 13.4	200
Hardships and Support during Pregnancy	13.7 10.5 10.5	700	19.5 12.4 20.0	200	10.5 5.0 24.7	200	9.9 3 . 0 13. 4	200
Experienced two or more hardships during childhood	27.2 23.0 - 31.4	1,300	40.5 31.1 - 49.8	500	32 .9 23.5 - 42.3	400	18.4 13.5 - 23.3	400
Homeless or did not have a regular place to sleep	3.7 1.6 - 5.7	200	40.5 31.1 - 49.8 7.4 * 1.7 - 13.1	<100	$52.9 \ 23.5 \ - \ 42.3 \ 5.6^{*} \ 0.2 \ - \ 11.0$	400 <100	18.4 13.5 - 23.3 	400
Moved due to problems paying rent or mortgage	6.5 4.0 - 8.9	300	12.4 5.4 - 19.3	200	10.1 * 4.0 - 16.1	100	 1.9 * 0.3 - 3.4	<100
Woman or partner lost job	13.0 9.6 - 16.4	600	24.8 16.2 - 33.5	300	15.1 7.3 - 22.9	200	4.7 2.1 - 7.2	100
Woman or partner had pay or hours cut back	14.4 10.7 - 18.0	700	23.6 14.7 - 32.4	300	17.3 8.8 - 25.8	200	8.0 4.2 - 11.7	200
Became separated or divorced	7.1 4.4 - 9.7	300	14.5 7.4 - 21.6	200	9.0 * 2.5 - 15.6	100	1.5 * 0.3 - 2.7	<100
Had no practical or emotional support	4.5 1.9 - 7.0	200	8.2 * 1.8 - 14.7	100				
Substance Use								
Any smoking, 3 months before pregnancy	13.7 10.4 - 17.1	700	19.5 11.5 - 27.5	200	13.4 6.3 - 20.6	200	11.0 6.7 - 15.2	300
Any smoking, 3rd trimester ⁺	3.1 * 1.2 - 5.0	200	7.9 * 1.7 - 14.2	100				
Any smoking, postpartum	5.3 3.0 - 7.5	300	10.1 * 3.5 - 16.7	100	3.9 * 0.8 - 7.1	<100	3.0 * 0.7 - 5.3	<100
Any binge drinking, 3 months before pregnancy	18.4 14.6 - 22.2	900	15.8 8.5 - 23.0	200	17.1 8.8 - 25.4	200	21.1 15.4 - 26.7	500
Any alcohol use, 3rd trimester [†]	10.0 7.3 - 12.7	500	4.8 * 1.3 - 8.3	<100	2.4 * 0.4 - 4.4	<100	17.1 12.0 - 22.1	400
Pregnancy Intention and Family Planning								
Mistimed or unwanted pregnancy	26.1 21.9 - 30.4	1,300	39.3 29.8 - 48.7	500	32.0 22.5 - 41.6	400	15.5 10.6 - 20.4	400
Unsure of pregnancy intentions	11.0 8.3 - 13.8	500	14.2 8.6 - 19.9	200	11.9 6.3 - 17.6	100	8.0 4.0 - 12.0	200
Postpartum birth control use ⁺	87.1 84.0 - 90.2	4,300	87.6 81.8 - 93.4	1,100	90.2 85.2 - 95.2	1,000	86.3 81.5 - 91.1	2,000
Infant Sleep and Breastfeeding								
Placed infant on back to sleep	83.7 80.1 - 87.3	4,000	85.8 79.3 - 92.3	1,000	81.0 72.8 - 89.2	900	86.7 82.1 - 91.2	2,000
Infant always or often shared bed	39.7 35.0 - 44.4	1,900	45.9 36.3 - 55.5	600	49.7 39.8 - 59.6	600	30.8 24.4 - 37.1	700
Intended to breastfeed, before birth	95.4 93.5 - 97.4	4,600	95.1 * 91.0 - 99.2	1,200			95.1 * 92.0 - 98.1	2,200
Intended to breastfeed exclusively, before birth	81.0 77.3 - 84.7	3,900	76.2 68.4 - 84.1	900	77.6 69.6 - 85.6	900	86.3 81.4 - 91.1	2,000
Breastfeeding practices								
Any breastfeeding, 1 month after delivery	92.3 89.9 - 94.7	4,500	89.5 84.2 - 94.7	1,100	92.1 * 87.2 - 97.1	1,000	94.5 * 91.2 - 97.8	2,200
Exclusive breastfeeding, 1 month after delivery	63.5 58.9 - 68.2	3,000	59.9 50.4 - 69.3	700	64.3 54.8 - 73.8	700	66.7 60.0 - 73.3	1,500
Any breastfeeding, 3 months after delivery	82.8 78.6 - 87.1	3,000	71.4 61.4 - 81.4	700	84.3 76.2 - 92.3	700	89.2 84.2 - 94.2	1,400
Exclusive breastfeeding, 3 months after delivery	49.0 43.4 - 54.6	1,800	40.6 30.1 - 51.0	400	52.4 40.6 - 64.2	400	54.3 45.8 - 62.7	900



MIHA Data Snapshot, Sonoma County by Income, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Sonoma Count	у	0-100% Federal Poverty	Guideline	101-200% Federal Povert	y Guideline	> 200% Federal Poverty	Guideline
	% 95% CI	Ν	% 95% CI	Ν	% 95% CI	N	% 95% CI	Ν
Service Utilization and Health Insurance Coverage								
Had a usual source of pre-pregnancy care	80.9 77.1 - 84.6	4,000	70.7 62.1 - 79.4	900	75.6 66.9 - 84.3	900	90.1 86.3 - 94.0	2,100
Initiated prenatal care in 1st trimester	92.3 89.9 - 94.7	4,500	85.9 79.3 - 92.6	1,100	92.7 * 87.9 - 97.6	1,000	97.3 * 95.5 - 99.1	2,200
Mom or infant needed but couldn't afford care postpartum	8.5 5.6 - 11.5	400	14.0 6.5 - 21.6	200	9.2 * 2.4 - 16.1	100	5.6 2.5 - 8.7	100
Participated in WIC during pregnancy ⁺	41.9 37.2 - 46.6	2,100	81.5 73.7 - 89.3	1,000	66.1 56.9 - 75.2	700	5.6 2.6 - 8.6	100
Received CalFresh (food stamps) during pregnancy	19.3 15.6 - 23.0	1,000	51.6 42.2 - 61.1	700	20.1 11.7 - 28.5	200		
Pre-pregnancy insurance								
Medi-Cal	17.4 14.0 - 20.9	900	43.5 34.2 - 52.7	600	20.9 13.1 - 28.7	200		
Uninsured	24.4 20.4 - 28.5	1,200	40.4 31.3 - 49.6	500	37.0 27.4 - 46.6	400	5.3 2.8 - 7.8	100
Private	55.3 50.6 - 60.0	2,700	14.0 6.8 - 21.2	200	39.0 29.3 - 48.7	400	90.9 86.8 - 95.1	2,100
Prenatal insurance								
Medi-Cal	47.1 42.4 - 51.9	2,300	93.4 * 88.7 - 98.0	1,200	71.4 62.9 - 79.9	800	5.0 2.4 - 7.7	100
Uninsured	1.1 * 0.3 - 1.8	<100						
Private	48.5 43.8 - 53.3	2,400	5.8 * 1.3 - 10.4	<100	22.6 14.8 - 30.5	300	89.5 85.1 - 93.8	2,000
Maternal postpartum insurance								
Medi-Cal	30.6 26.1 - 35.0	1,500	66.6 58.2 - 75.1	800	42.2 32.3 - 52.1	500	3.6 * 1.4 - 5.9	<100
Uninsured	15.8 12.5 - 19.1	800	28.2 20.4 - 36.1	400	21.1 12.9 - 29.3	200	2.8 * 0.8 - 4.7	<100
Private	51.0 46.2 - 55.8	2,500	4.7 * 0.4 - 9.0	<100	33.0 23.8 - 42.2	400	89.9 85.6 - 94.2	2,100
Infant health insurance								
Medi-Cal	48.2 43.5 - 53.0	2,300	92.0 * 86.0 - 98.0	1,100	74.0 65.8 - 82.3	800	8.3 4.9 - 11.7	200
Uninsured	1.1 [*] 0.2 - 1.9	<100						
Private	47.1 42.3 - 51.9	2,300			18.8 11.6 - 26.0	200	87.3 82.6 - 92.0	2,000
Maternal Demographics								
Did not complete high school (or GED)	14.9 11.6 - 18.3	700	37.1 28.1 - 46.1	500	11.8 5.6 - 18.0	100		
Unmarried	33.2 28.9 - 37.5	1,600	57.6 48.3 - 66.9	700	34.6 25.9 - 43.3	400	15.6 11.1 - 20.1	400
Race/Ethnicity								
White	51.7 46.8 - 56.5	2,300	25.5 16.4 - 34.6	300	35.9 25.8 - 45.9	400	78.9 72.7 - 85.0	1,600
Hispanic	40.8 36.0 - 45.5	1,800	65.4 55.7 - 75.1	800	59.4 49.2 - 69.6	600	12.3 7.8 - 16.8	300
Asian/Pacific Islander	5.1 2.4 - 7.7	200	5.5 * 0.3 - 10.7	<100			7.5 * 2.8 - 12.2	200
Black	1.4 0.6 - 2.2	<100						
Born outside the U.S.	30.6 26.2 - 35.0	1,500	48.5 39.1 - 57.8	600	41.3 31.5 - 51.0	500	11.1 6.2 - 16.0	300
Speaks non-English language at home	24.8 20.8 - 28.8	1,200	43.3 34.3 - 52.4	600	35.2 25.5 - 44.8	400	5.4 * 2.2 - 8.7	100
Lives in a high poverty neighborhood	17.0 13.6 - 20.3	800	26.2 18.4 - 33.9	300	19.6 12.2 - 27.0	200	10.4 6.3 - 14.5	200

-- Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

* Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

⁺ Measure/definition may not be comparable to previous Snapshots (see Technical Notes).

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Sonoma County was 557. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.



MIHA Data Snapshot, Sonoma County by Maternal Age, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Sonoma Count	y	15-19 Years		20-34 Years		35 Years or Olde	er
	% 95% CI	Ν	% 95% CI	Ν	% 95% CI	N	% 95% CI	Ν
Total	100	4,900	3.8 2.1 - 5.5	200	74.0 69.5 - 78.5	3,700	22.2 17.8 - 26.5	1,100
Health Status before Pregnancy								
In good to excellent health	93.5 91.2 - 95.8	4,600	78.2 * 59.7 - 96.7	100	93.8 91.4 - 96.3	3,400	94.9 * 89.9 - 99.8	1,000
Chronic conditions								
Diabetes [†]	2.2 1.0 - 3.4	100			2.0 * 0.7 - 3.3	<100		
Hypertension ⁺	3.3 1.9 - 4.8	200			3.2 1.5 - 4.9	100	4.3 * 0.6 - 8.0	<100
Asthma ⁺	10.7 7.9 - 13.5	500	39.5 * 15.8 - 63.3	<100	10.0 7.0 - 13.1	400	8.1 * 2.7 - 13.5	<100
Nutrition and Weight								
Daily folic acid use, month before pregnancy	38.7 34.0 - 43.4	1,900			38.1 33.0 - 43.3	1,400	45.9 34.4 - 57.5	500
Overweight before pregnancy	24.0 19.9 - 28.1	1,100			26.1 21.5 - 30.8	900	18.5 9.3 - 27.7	200
Obese before pregnancy	21.7 17.5 - 25.9	1,000			21.2 16.8 - 25.6	700	25.1 13.5 - 36.7	300
Inadequate weight gain during pregnancy	14.6 11.1 - 18.0	600			15.3 11.3 - 19.3	500	10.2 4.3 - 16.1	<100
Excessive weight gain during pregnancy	48.5 43.3 - 53.7	2,000	35.8 * 13.4 - 58.3	<100	50.7 44.9 - 56.4	1,500	43.8 31.4 - 56.2	400
Food insecurity during pregnancy	14.7 11.2 - 18.2	700			15.6 11.4 - 19.8	600	11.4 5.0 - 17.8	100
Intimate Partner Violence (IPV) and Depressive Symptoms								
Physical or psychological IPV during pregnancy	6.0 3.5 - 8.6	300			6.1 3.0 - 9.2	200	3.1 [*] 0.3 - 5.9	<100
Prenatal depressive symptoms	14.1 10.8 - 17.5	700	23.9 * 5.2 - 42.6	<100	15.6 11.5 - 19.7	600	7.3 * 2.5 - 12.1	<100
Postpartum depressive symptoms	13.7 10.5 - 16.9	700			14.9 11.1 - 18.7	500	10.5 * 4.0 - 17.1	100
Hardships and Support during Pregnancy								
Experienced two or more hardships during childhood	27.2 23.0 - 31.4	1,300	43.8 19.6 - 68.0	<100	27.6 22.8 - 32.4	1,000	23.2 14.6 - 31.8	300
Homeless or did not have a regular place to sleep	3.7 1.6 - 5.7	200			3.6 * 1.3 - 5.9	100		
Moved due to problems paying rent or mortgage	6.5 4.0 - 8.9	300			6.3 3.6 - 9.1	200	5.5 * 0.4 - 10.6	<100
Woman or partner lost job	13.0 9.6 - 16.4	600	39.5 * 15.2 - 63.9	<100	12.5 8.7 - 16.3	500	10.6 * 3.6 - 17.6	100
Woman or partner had pay or hours cut back	14.4 10.7 - 18.0	700			14.2 10.0 - 18.5	500	15.2 7.2 - 23.2	200
Became separated or divorced	7.1 4.4 - 9.7	300			7.7 4.5 - 10.9	300	5.0 * 0.2 - 9.9	<100
Had no practical or emotional support	4.5 1.9 - 7.0	200			3.5 * 1.2 - 5.8	100		
Substance Use			10.1.*				0.0.÷	
Any smoking, 3 months before pregnancy	13.7 10.4 - 17.1	700	16.1 * 1.6 - 30.5	<100	14.8 10.8 - 18.9	500	9.8 * 3.4 - 16.1	100
Any smoking, 3rd trimester [†]	3.1 * 1.2 - 5.0	200			3.8 * 1.3 - 6.2	100		
Any smoking, postpartum	5.3 3.0 - 7.5	300			6.4 3.5 - 9.3	200		200
Any binge drinking, 3 months before pregnancy Any alcohol use, 3rd trimester ⁺	18.4 14.6 - 22.2	900 500	24.6 * 4.6 - 44.5	<100	19.4 14.9 - 23.8 9.0 6.1 - 11.9	700 300	14.2 6.6 - 21.9 14.2 7.2 - 21.3	200 200
	10.0 7.3 - 12.7	500			9.0 6.1 - 11.9	300	14.2 7.2 - 21.3	200
Pregnancy Intention and Family Planning Mistimed or unwanted pregnancy	26.4 24.0 20.4	4 200	10 6 * 46 2 64 2	.400	27.0 22.0 22.0	4 000	47.2 04 255	200
1 0 1	26.1 21.9 - 30.4	1,300	40.6 * 16.3 - 64.9	<100	27.9 23.0 - 32.9	1,000	17.3 9.1 - 25.5	200
Unsure of pregnancy intentions Postpartum birth control use [†]	11.0 8.3 - 13.8 87.1 84.0 - 90.2	500 4,300	20.1 * 3.6 - 36.6	<100	8.5 6.0 - 11.1 87.1 83.7 - 90.6	300 3,100	18.0 9.6 - 26.4 86.1 78.2 - 94.0	200 900
•	87.1 84.0 - 90.2	4,300			87.1 83.7 - 90.6	3,100	80.1 /8.2 - 94.0	900
Infant Sleep and Breastfeeding Placed infant on back to sleep	02 7 00 4 07 0	4.000	00.4 * ca a	200	04.4 00.4 00.4	2 000	02.2 70.6 00.7	000
Infant always or often shared bed	83.7 80.1 - 87.3	4,000	80.1 * 63.3 - 96.8	200	84.4 80.4 - 88.4	3,000	82.2 73.6 - 90.7	900
Intended to breastfeed, before birth	39.7 35.0 - 44.4	1,900	55.0 32.0 - 77.9	100	37.8 32.7 - 43.0	1,400	43.2 31.8 - 54.6	500
Intended to breastfeed exclusively, before birth	95.4 93.5 - 97.4 81.0 77.3 - 84.7	4,600	 74.6 * 55.7 - 93.6	100	95.6 93.4 - 97.8 80.2 76.0 - 84.4	3,400	 84.6 76.7 - 92.4	000
Breastfeeding practices	ol.U //.3 - 84./	3,900	/4.0 >>./ - 93.6	100	00.2 /0.0 - 84.4	2,900	04.0 /0./ - 92.4	900
Any breastfeeding, 1 month after delivery	92.3 89.9 - 94.7	4,500	82.2 * 66.3 - 98.0	200	93.2 90.7 - 95.7	3,300	91.1 * 84.5 - 97.6	1,000
Exclusive breastfeeding, 1 month after delivery	63.5 58.9 - 68.2	3,000	56.9 33.6 - 80.2	<100	63.1 57.9 - 68.3	2,200	66.1 55.3 - 76.8	700
Any breastfeeding, 3 months after delivery	82.8 78.6 - 87.1	3,000	75.8 * 55.4 - 96.3	100	80.7 75.5 - 85.9	2,200	91.4 * 85.6 - 97.2	700
Exclusive breastfeeding, 3 months after delivery	49.0 43.4 - 54.6	1,800	43.4 * 15.9 - 70.9	<100	48.1 41.8 - 54.4	1,300	53.0 39.9 - 66.1	400
Exclusive breastreeding, 5 months after delivery	+3.0 +3.4 - 34.0	1,000	43.4 13.3 - 70.3	~100	40.1 71.0 - 34.4	1,500	JJ. <i>JJ. J</i> J J J J J J J J J J	400

MIHA Data Snapshot, Sonoma County by Maternal Age, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Sonoma Count	У	15-19 Years		20-34 Years		35 Years or Olde	er
	% 95% CI	N	% 95% CI	N	% 95% CI	N	% 95% CI	N
Service Utilization and Health Insurance Coverage								
Had a usual source of pre-pregnancy care	80.9 77.1 - 84.6	4,000	49.9 26.6 - 73.3	<100	80.2 75.9 - 84.5	2,900	88.5 82.1 - 94.9	1,000
Initiated prenatal care in 1st trimester	92.3 89.9 - 94.7	4,500	69.0 * 45.5 - 92.5	100	92.3 89.6 - 95.0	3,300	96.6 * 93.5 - 99.6	1,000
Mom or infant needed but couldn't afford care postpartum	8.5 5.6 - 11.5	400			9.1 5.5 - 12.8	300	5.6 * 0.7 - 10.4	<100
Participated in WIC during pregnancy ⁺	41.9 37.2 - 46.6	2,100			42.8 37.5 - 48.0	1,600	31.9 21.6 - 42.2	300
Received CalFresh (food stamps) during pregnancy	19.3 15.6 - 23.0	1,000	49.6 25.8 - 73.3	<100	19.7 15.3 - 24.1	700	13.4 7.5 - 19.2	100
Pre-pregnancy insurance								
Medi-Cal	17.4 14.0 - 20.9	900	48.2 25.3 - 71.0	<100	15.9 12.3 - 19.6	600	17.2 8.1 - 26.3	200
Uninsured	24.4 20.4 - 28.5	1,200	40.6 * 16.5 - 64.7	<100	25.2 20.5 - 29.8	900	19.2 11.5 - 27.0	200
Private	55.3 50.6 - 60.0	2,700			57.1 51.9 - 62.3	2,100	57.6 46.3 - 68.9	600
Prenatal insurance								
Medi-Cal	47.1 42.4 - 51.9	2,300			48.0 42.7 - 53.2	1,700	36.6 25.8 - 47.3	400
Uninsured	1.1 * 0.3 - 1.8	<100						
Private	48.5 43.8 - 53.3	2,400			49.3 44.0 - 54.6	1,800	53.8 42.4 - 65.3	600
Maternal postpartum insurance								
Medi-Cal	30.6 26.1 - 35.0	1,500	78.2 * 59.8 - 96.7	100	30.8 25.8 - 35.8	1,100	21.9 12.5 - 31.3	200
Uninsured	15.8 12.5 - 19.1	800			15.9 12.0 - 19.8	600	15.3 8.2 - 22.3	200
Private	51.0 46.2 - 55.8	2,500			51.7 46.4 - 57.0	1,900	56.3 44.9 - 67.6	600
Infant health insurance								
Medi-Cal	48.2 43.5 - 53.0	2,300			48.5 43.2 - 53.8	1,700	39.0 28.0 - 49.9	400
Uninsured	1.1 * 0.2 - 1.9	<100			1.5 * 0.3 - 2.6	<100		
Private	47.1 42.3 - 51.9	2,300			47.4 42.1 - 52.7	1,700	54.1 42.6 - 65.5	600
Maternal Demographics								
Did not complete high school (or GED)	14.9 11.6 - 18.3	700	55.3 31.3 - 79.3	100	12.4 8.8 - 16.0	400	16.5 8.8 - 24.3	200
Unmarried	33.2 28.9 - 37.5	1,600	77.9 * 57.1 - 98.7	100	34.9 29.9 - 39.9	1,300	20.1 12.2 - 28.0	200
Race/Ethnicity								
White	51.7 46.8 - 56.5	2,300	19.0 * 3.8 - 34.3	<100	51.5 46.1 - 57.0	1,700	58.3 46.9 - 69.8	600
Hispanic	40.8 36.0 - 45.5	1,800	81.0 * 65.7 - 96.2	200	40.7 35.3 - 46.0	1,400	33.5 23.0 - 44.1	300
Asian/Pacific Islander	5.1 2.4 - 7.7	200			4.8 2.1 - 7.5	200		
Black	1.4 0.6 - 2.2	<100			1.7 * 0.5 - 2.9	<100		
Born outside the U.S.	30.6 26.2 - 35.0	1,500	29.2 * 8.6 - 49.9	<100	29.8 24.9 - 34.7	1,100	33.4 22.5 - 44.3	400
Speaks non-English language at home	24.8 20.8 - 28.8	1,200	25.9 * 5.5 - 46.3	<100	25.2 20.5 - 29.9	900	23.3 14.7 - 32.0	300
Lives in a high poverty neighborhood	17.0 13.6 - 20.3	800	27.4 * 6.3 - 48.4	<100	19.1 15.0 - 23.2	700	7.8 3.5 - 12.2	<100

- - Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

* Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

[†] Measure/definition may not be comparable to previous Snapshots (see Technical Notes).

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Sonoma County was 557. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.



MIHA Data Snapshot, Sonoma County by Neighborhood Poverty, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Son	ioma Cour	nty		9% of Reside		5-9.9% of Resi below Pove		10-19.9% of Res below Pover		≥ 20% of Reside	
	%	95% CI	N	%	95% CI	Ν	% 95% CI	N	% 95% CI	N	% 95% CI	Ν
Total	100		4,900	8.9	6.0 - 11.9	400	27.6 23.2 - 32.2	1,300	46.4 41.6 - 51.2	2,200	17.0 13.6 - 20.3	800
Health Status before Pregnancy												
In good to excellent health Chronic conditions	93.5	91.2 - 95.8	4,600				96.3 * 93.0 - 99.0	5 1,300	92.2 88.7 - 95.8	2,100	93.3 * 87.8 - 98.8	800
Diabetes ⁺	2.2	1.0 - 3.4	100						3.0 * 1.1 - 5.0	<100		
Hypertension ⁺	3.3	1.9 - 4.8	200						3.8 1.7 - 6.0	<100		
Asthma ⁺	10.7	7.9 - 13.5	500				10.6 5.2 - 16.3	. 100	12.5 8.0 - 16.9	300	7.8 * 2.6 - 12.9	<100
Nutrition and Weight												
Daily folic acid use, month before pregnancy	38.7 3	84.0 - 43.4	1,900	24.1	9.7 - 38.5	100	41.4 31.8 - 50.9	500	39.6 32.7 - 46.4	900	40.2 29.8 - 50.7	300
Overweight before pregnancy	24.0 1	9.9 - 28.1	1,100	17.3 *	6.5 - 28.1	<100	26.0 17.4 - 34.6	300	22.3 16.5 - 28.1	500	26.3 16.5 - 36.0	200
Obese before pregnancy	21.7 1	7.5 - 25.9	1,000	27.2	10.7 - 43.8	100	17.2 8.6 - 25.9	200	24.1 17.9 - 30.3	500	22.0 13.2 - 30.7	200
Inadequate weight gain during pregnancy	14.6 1	1.1 - 18.0	600				10.1 5.0 - 15.2	100	17.8 12.0 - 23.5	300	20.5 11.1 - 29.9	100
Excessive weight gain during pregnancy	48.5 4	3.3 - 53.7	2,000	38.6	21.3 - 56.0	100	4 7.9 37.7 - 58.2	600	51.6 44.0 - 59.2	900	45.8 34.0 - 57.6	300
Food insecurity during pregnancy	14.7 1	1.2 - 18.2	700	20.4 '	2.9 - 37.9	<100	9.9 * 3.4 - 16.4	100	14.6 10.0 - 19.2	300	19.0 10.5 - 27.6	200
Intimate Partner Violence (IPV) and Depressive Symptoms												
Physical or psychological IPV during pregnancy	6.0	3.5 - 8.6	300				5.9 * 0.3 - 11.0	<100	4.3 [*] 1.8 - 6.8	<100	8.5 * 2.1 - 14.8	<100
Prenatal depressive symptoms		0.8 - 17.5	700				16.1 8.7 - 23.5		10.6 6.7 - 14.4	200	19.4 11.0 - 27.7	200
Postpartum depressive symptoms	13.7 1	0.5 - 16.9	700				10.2 5.0 - 15.5	100	10.3 6.6 - 14.1	200	22.6 13.7 - 31.5	200
Hardships and Support during Pregnancy												
Experienced two or more hardships during childhood		23.0 - 31.4	1,300	22.6 *	5.2 - 39.9	<100	22.3 14.7 - 29.9	300	28.2 22.1 - 34.3	600	30.9 20.9 - 40.8	200
Homeless or did not have a regular place to sleep	-	1.6 - 5.7	200						3.0 * 0.5 - 5.5	<100		
Moved due to problems paying rent or mortgage		4.0 - 8.9	300				6.0 * 0.6 - 11.3		6.9 3.4 - 10.4	200	7.6 * 1.2 - 14.1	<100
Woman or partner lost job		9.6 - 16.4	600	17.3 '	0.5 - 34.1	<100	12.6 5.7 - 19.5		13.4 8.9 - 17.9	300	12.1 * 4.7 - 19.5	<100
Woman or partner had pay or hours cut back		0.7 - 18.0	700				12.6 5.5 - 19.3		16.1 10.8 - 21.4	400	13.6 6.2 - 21.0	100
Became separated or divorced		4.4 - 9.7	300				10.5 * 3.8 - 17.3	. 100	3.2 1.4 - 5.0	<100	9.0 * 3.1 - 15.0	<100
Had no practical or emotional support	4.5	1.9 - 7.0	200						6.2 * 1.9 - 10.5	100		
Substance Use												
Any smoking, 3 months before pregnancy		0.4 - 17.1	700				14.7 7.6 - 21.9		13.4 8.9 - 17.9	300	10.1 * 3.7 - 16.5	<100
Any smoking, 3rd trimester ⁺	-	1.2 - 5.0	200				6.2 * 0.6 - 11.8					
Any smoking, postpartum		3.0 - 7.5	300				7.4 * 1.7 - 13.		4.1 * 1.6 - 6.7	<100		
Any binge drinking, 3 months before pregnancy	-	4.6 - 22.2	900	18.5 '	2.0 - 35.0	<100	17.4 9.8 - 24.9		20.0 14.4 - 25.6	400	17.9 9.5 - 26.2	100
Any alcohol use, 3rd trimester [†]	10.0	7.3 - 12.7	500				14.1 7.9 - 20.2	200	7.6 4.1 - 11.1	200	10.4 * 4.0 - 16.9	<100
Pregnancy Intention and Family Planning	26.4			20.4			04 E		04.4		20.0	
Mistimed or unwanted pregnancy	-	21.9 - 30.4	1,300		12.7 - 48.0	100	31.5 22.2 - 40.3		21.1 15.5 - 26.8	500	29.0 19.2 - 38.8	200
Unsure of pregnancy intentions Postpartum birth control use [†]	-	8.3 - 13.8	500	-	1.6 - 27.3	<100	9.3 4.4 - 14.2		11.6 7.5 - 15.6	300	11.4 5.2 - 17.7	<100
	87.1 8	34.0 - 90.2	4,300	87.9	78.0 - 97.8	400	88.2 82.5 - 93.8	3 1,200	85.7 80.7 - 90.8	1,900	90.3 * 84.5 - 96.1	700
Infant Sleep and Breastfeeding												
Placed infant on back to sleep		80.1 - 87.3	4,000				84.5 77.8 - 91.2		80.8 75.2 - 86.3	1,700	88.7 82.3 - 95.2	700
Infant always or often shared bed Intended to breastfeed, before birth		35.0 - 44.4	1,900	-	16.8 - 51.6	100	33.2 24.1 - 42.4		42.9 36.0 - 49.8	900	45.7 35.0 - 56.3	400
Intended to breastfeed exclusively, before birth		93.5 - 97.4 77.3 - 84.7	4,600 3,900		54.2 - 85.9	300	94.0 * 89.6 - 98.9 81.1 73.6 - 88.0		96.7 * 94.2 - 99.1 85.3 80.7 - 90.0	2,100 1,900	 72.2 62.6 - 81.8	600
Breastfeeding practices	01.0 /	7.3 - 84./	5,900	70.0	J4.Z - 83.9	300	OI.I /3.0 - 88.0	, 1,100	00.0 00.7 - 90.0	1,900	12.2 02.0 - 81.8	000
Any breastfeeding, 1 month after delivery	973 9	89.9 - 94.7	4,500	86.2 *	73.5 - 98.9	400	93.6 * 89.2 - 98.0	1,200	94.0 91.2 - 96.8	2,100	87.9 * 80.5 - 95.2	700
Exclusive breastfeeding, 1 month after delivery		58.9 - 68.2	3,000		36.5 - 71.6	200	60.6 51.0 - 70.2		68.9 62.6 - 75.1	1,500	59.3 48.6 - 70.0	500
Any breastfeeding, 3 months after delivery		78.6 - 87.1	3,000		55.7 - 93.3	200	83.2 73.8 - 92.0		83.8 78.4 - 89.2	1,400	81.7 72.2 - 91.1	500
Exclusive breastfeeding, 3 months after delivery		3.4 - 54.6	1,800	-	25.9 - 68.5	100	50.1 38.0 - 62.1		49.8 42.0 - 57.5	800	46.2 33.7 - 58.7	300



MIHA Data Snapshot, Sonoma County by Neighborhood Poverty, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Sc	noma Coun	ty		.9% of Reside below Povert			.9% of Reside			9.9% of Reside			0% of Reside elow Povert	
	%	95% CI	N	%	95% CI	Ν	%	95% CI	N	%	95% CI	N	%	95% CI	Ν
Service Utilization and Health Insurance Coverage															
Had a usual source of pre-pregnancy care	80.9	77.1 - 84.6	4,000	80.8	* 65.5 - 96.2	300	82.8	75.6 - 90.0	1,100	79.2	73.7 - 84.7	1,800	83.0	75.1 - 90.9	700
Initiated prenatal care in 1st trimester	92.3	89.9 - 94.7	4,500				95.9	* 92.9 - 98.9	1,200	90.7	86.8 - 94.6	2,000	90.0	* 82.9 - 97.2	700
Mom or infant needed but couldn't afford care postpartum	8.5	5.6 - 11.5	400				8.7	* 2.5 - 14.9	100	9.4	5.2 - 13.6	200	6.7	* 1.7 - 11.8	<100
Participated in WIC during pregnancy [†]	41.9	37.2 - 46.6	2,100	33.0	15.8 - 50.1	100	26.3	17.6 - 35.0	300	46.4	39.5 - 53.3	1,000	63.5	53.1 - 73.9	500
Received CalFresh (food stamps) during pregnancy	19.3	15.6 - 23.0	1,000	20.6	* 3.4 - 37.8	<100	13.9	7.1 - 20.6	200	16.8	12.1 - 21.5	400	32.4	22.4 - 42.3	300
Pre-pregnancy insurance															
Medi-Cal	17.4	14.0 - 20.9	900	14.1	* 3.0 - 25.2	<100	11.5	5.9 - 17.0	200	19.9	14.2 - 25.6	400	18.7	10.5 - 26.9	200
Uninsured	24.4	20.4 - 28.5	1,200	24.5	* 7.1 - 41.9	100	11.6	5.6 - 17.5	200	29.0	22.8 - 35.1	600	33.9	23.8 - 44.0	300
Private	55.3	50.6 - 60.0	2,700	61.4	43.5 - 79.2	300	70.1	61.2 - 79.0	900	50.0	43.1 - 56.8	1,100	44.5	33.9 - 55.2	400
Prenatal insurance															
Medi-Cal	47.1	42.4 - 51.9	2,300	41.3	23.5 - 59.2	200	32.9	23.8 - 42.0	400	51.8	44.9 - 58.7	1,100	59.0	48.5 - 69.5	500
Uninsured	1.1 *	0.3 - 1.8	<100												
Private	48.5	43.8 - 53.3	2,400	56.3	38.5 - 74.1	200	57.2	47.6 - 66.9	800	46.2	39.4 - 53.1	1,000	37.4	27.0 - 47.8	300
Maternal postpartum insurance															
Medi-Cal	30.6	26.1 - 35.0	1,500	28.7	* 10.9 - 46.6	100	25.6	17.1 - 34.1	300	31.6	25.0 - 38.1	700	34.5	24.4 - 44.6	300
Uninsured	15.8	12.5 - 19.1	800	9.9	* 0.7 - 19.1	<100	6.6	* 1.9 - 11.3	<100	19.2	13.8 - 24.6	400	25.9	16.6 - 35.1	200
Private	51.0	46.2 - 55.8	2,500	61.3	43.3 - 79.3	300	60.0	50.4 - 69.7	800	48.1	41.2 - 55.0	1,100	39.6	29.1 - 50.2	300
Infant health insurance															
Medi-Cal	48.2	43.5 - 53.0	2,300	41.6	22.9 - 60.3	200	30.7	21.9 - 39.4	400	55.6	48.8 - 62.4	1,200	61.3	50.8 - 71.9	500
Uninsured	1.1 *	0.2 - 1.9	<100												
Private	47.1	42.3 - 51.9	2,300	54.4	35.9 - 72.9	200	57.8	48.1 - 67.4	800	42.2	35.4 - 49.0	900	37.4	27.0 - 47.9	300
Maternal Demographics															
Age															
15-19	3.8	2.1 - 5.5	200							5.0	2.1 - 7.9	100	6.3	* 0.8 - 11.9	<100
20-34	74.0	69.5 - 78.5	3,700	73.2	* 56.9 - 89.5	300	70.7	61.0 - 80.3	900	73.6	67.2 - 80.1	1,600	83.8	76.3 - 91.2	700
35+	22.2	17.8 - 26.5	1,100	26.8	* 10.5 - 43.1	100	27.4	17.8 - 37.0	400	21.4	15.2 - 27.5	500	9.9	4.6 - 15.3	<100
Did not complete high school (or GED)	14.9	11.6 - 18.3	700				11.6	5.3 - 17.9	200	15.4	10.4 - 20.3	300	21.0	12.2 - 29.8	200
Unmarried	33.2	28.9 - 37.5	1,600	26.1	11.4 - 40.7	100	32.0	23.1 - 40.9	400	31.0	25.0 - 37.0	700	42.6	32.1 - 53.2	300
Race/Ethnicity															
White	51.7	46.8 - 56.5	2,300	70.0	55.0 - 85.0	300	68.8	60.0 - 77.6	800	44.3	37.4 - 51.3	900	34.2	23.7 - 44.7	300
Hispanic	40.8	36.0 - 45.5	1,800	28.4	13.6 - 43.2	100	24.6	16.5 - 32.6	300	47.5	40.4 - 54.5	1,000	58.9	47.8 - 69.9	400
Asian/Pacific Islander	5.1	2.4 - 7.7	200							5.5	* 1.7 - 9.3	100			
Black	1.4	0.6 - 2.2	<100												
Born outside the U.S.	30.6	26.2 - 35.0	1,500	28.9	13.6 - 44.1	100	16.0	9.4 - 22.6	200	35.6	28.8 - 42.5	800	42.1	31.7 - 52.5	300
Speaks non-English language at home	24.8	20.8 - 28.8	1,200	15.1	* 3.9 - 26.3	<100	13.8	7.6 - 20.0	200	30.0	23.5 - 36.5	700	37.0	26.9 - 47.1	300
Income as a percent of the Federal Poverty Guideline															
0-100%	27.2	22.8 - 31.5	1,300	21.4	* 6.3 - 36.5	<100	15.9	8.3 - 23.6	200	29.4	22.8 - 35.9	600	41.9	31.2 - 52.6	300
101-200%	24.0	19.9 - 28.2	1,100	19.0	* 3.4 - 34.5	<100	18.1	11.0 - 25.2	200	27.5	21.1 - 34.0	600	28.2	18.4 - 38.0	200
> 200%	48.8	44.0 - 53.7	2,300	59.6	41.6 - 77.7	300	66.0	56.7 - 75.3	800	43.1	36.2 - 50.0	900	29.9	19.7 - 40.1	200

-- Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

* Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

⁺ Measure/definition may not be comparable to previous Snapshots (see Technical Notes).

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Sonoma County was 557. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. Neighborhood poverty is the percentage of residents with incomes below poverty in a woman's census tract. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.



MIHA Data Snapshot, Sonoma County by Prenatal Health Insurance, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		Sonoma County			Medi-Cal			Private	
	%	95% CI	Ν	%	95% CI	Ν	%	95% CI	Ν
Total	100		4,900	47.1	42.4 - 51.9	2,300	48.5	43.8 - 53.3	2,400
Health Status before Pregnancy									
In good to excellent health Chronic conditions	93.5	91.2 - 95.8	4,600	88.2	83.7 - 92.6	2,000	98.3 *	96.9 - 99.7	2,400
Diabetes ⁺	2.2	1.0 - 3.4	100	2.5 *	0.8 - 4.1	<100	2.2 *	0.4 - 3.9	<100
Hypertension ⁺	3.3	1.9 - 4.8	200	4.2	1.8 - 6.6	<100	2.8 *	0.8 - 4.7	<100
Asthma ⁺	10.7	7.9 - 13.5	500	10.3	6.4 - 14.2	200	11.1	6.9 - 15.3	300
Nutrition and Weight									
Daily folic acid use, month before pregnancy	38.7	34.0 - 43.4	1,900	28.0	21.6 - 34.4	700	49.8	42.9 - 56.6	1,200
Overweight before pregnancy	24.0	19.9 - 28.1	1,100	26.6	20.2 - 32.9	500	21.3	16.0 - 26.5	500
Obese before pregnancy	21.7	17.5 - 25.9	1,000	24.6	18.1 - 31.1	500	19.9	14.0 - 25.8	500
Inadequate weight gain during pregnancy	14.6	11.1 - 18.0	600	20.9	14.6 - 27.2	400	9.9	6.2 - 13.7	200
Excessive weight gain during pregnancy	48.5	43.3 - 53.7	2,000	46.7	38.8 - 54.7	800	48.0	41.0 - 55.1	1,000
Food insecurity during pregnancy	14.7	11.2 - 18.2	700	27.0	20.6 - 33.4	600	4.3 *	1.6 - 6.9	100
Intimate Partner Violence (IPV) and Depressive Symptoms									
Physical or psychological IPV during pregnancy	6.0	3.5 - 8.6	300	11.0	5.9 - 16.0	300	1.4 *	0.2 - 2.6	<100
Prenatal depressive symptoms	14.1	10.8 - 17.5	700	20.4	14.6 - 26.2	500	8.7	5.1 - 12.4	200
Postpartum depressive symptoms	13.7	10.5 - 16.9	700	17.7	12.4 - 23.0	400	10.8	6.9 - 14.8	300
Hardships and Support during Pregnancy									
Experienced two or more hardships during childhood	27.2	23.0 - 31.4	1,300	36.5	29.7 - 43.4	800	19.1	14.2 - 23.9	500
Homeless or did not have a regular place to sleep	3.7	1.6 - 5.7	200	7.4	3.2 - 11.6	200			
Moved due to problems paying rent or mortgage	6.5	4.0 - 8.9	300	11.2	6.4 - 16.0	300	2.3 *	0.6 - 3.9	<100
Woman or partner lost job	13.0	9.6 - 16.4	600	22.8	16.4 - 29.2	500	4.5	2.1 - 6.9	100
Woman or partner had pay or hours cut back	14.4	10.7 - 18.0	700	21.2	15.0 - 27.5	500	8.4	4.4 - 12.4	200
Became separated or divorced	7.1	4.4 - 9.7	300	13.6	8.3 - 18.9	300	1.3 *	0.1 - 2.4	<100
Had no practical or emotional support	4.5	1.9 - 7.0	200	8.9	3.7 - 14.1	200			
Substance Use									
Any smoking, 3 months before pregnancy	13.7	10.4 - 17.1	700	17.5	11.9 - 23.2	400	11.0	7.0 - 15.1	300
Any smoking, 3rd trimester [†]	3.1 *	1.2 - 5.0	200	5.8 *	2.0 - 9.6	100			
Any smoking, postpartum	5.3	3.0 - 7.5	300	8.3	4.2 - 12.5	200	2.8 *	0.7 - 5.0	<100
Any binge drinking, 3 months before pregnancy	18.4	14.6 - 22.2	900	16.2	10.7 - 21.7	400	21.1	15.4 - 26.7	500
Any alcohol use, 3rd trimester ⁺	10.0	7.3 - 12.7	500	3.8 *	1.4 - 6.2	<100	16.0	11.3 - 20.8	400
Pregnancy Intention and Family Planning									
Mistimed or unwanted pregnancy	26.1	21.9 - 30.4	1,300	37.3	30.4 - 44.2	900	16.2	11.3 - 21.0	400
Unsure of pregnancy intentions	11.0	8.3 - 13.8	500	14.2	10.1 - 18.4	300	7.5	3.9 - 11.2	200
Postpartum birth control use [†]	87.1	84.0 - 90.2	4,300	87.1	82.5 - 91.7	2,000	86.7	82.1 - 91.2	2,100
Infant Sleep and Breastfeeding									
Placed infant on back to sleep	83.7	80.1 - 87.3	4,000	80.5	74.6 - 86.3	1,800	87.6	83.3 - 91.8	2,100
Infant always or often shared bed	39.7	35.0 - 44.4	1,900	47.8	40.7 - 54.8	1,100	30.2	24.2 - 36.3	700
Intended to breastfeed, before birth	95.4	93.5 - 97.4	4,600	95.4 *	92.5 - 98.3	2,100	95.8 *	93.3 - 98.4	2,300
Intended to breastfeed exclusively, before birth	81.0	77.3 - 84.7	3,900	77.3	71.7 - 83.0	1,700	85.0	80.2 - 89.8	2,000
Breastfeeding practices									
Any breastfeeding, 1 month after delivery	92.3	89.9 - 94.7	4,500	90.3	86.5 - 94.0	2,000	95.2 *	92.3 - 98.0	2,300
Exclusive breastfeeding, 1 month after delivery	63.5	58.9 - 68.2	3,000	60.5	53.5 - 67.5	1,300	67.4	61.0 - 73.9	1,600
Any breastfeeding, 3 months after delivery	82.8	78.6 - 87.1	3,000	76.6	69.8 - 83.4	1,400	88.6	83.4 - 93.7	1,500
Exclusive breastfeeding, 3 months after delivery	49.0	43.4 - 54.6	1,800	45.5	37.4 - 53.6	800	53.6	45.3 - 61.8	900



MIHA Data Snapshot, Sonoma County by Prenatal Health Insurance, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		Sonoma County			Medi-Cal			Private	
	%	95% CI	Ν	%	95% CI	Ν	%	95% CI	Ν
Service Utilization and Health Insurance Coverage									
Had a usual source of pre-pregnancy care	80.9	77.1 - 84.6	4,000	71.4	65.0 - 77.8	1,700	89.3	85.4 - 93.2	2,100
Initiated prenatal care in 1st trimester	92.3	89.9 - 94.7	4,500	86.4	81.7 - 91.0	1,900	97.6 *	95.8 - 99.3	2,300
Mom or infant needed but couldn't afford care postpartum	8.5	5.6 - 11.5	400	12.1	6.9 - 17.2	300	5.3 *	2.0 - 8.6	100
Participated in WIC during pregnancy ⁺	41.9	37.2 - 46.6	2,100	79.1	73.5 - 84.7	1,800	7.8	4.4 - 11.2	200
Received CalFresh (food stamps) during pregnancy	19.3	15.6 - 23.0	1,000	40.4	33.6 - 47.2	900			
Pre-pregnancy insurance									
Medi-Cal	17.4	14.0 - 20.9	900	37.1	30.6 - 43.7	900			
Uninsured	24.4	20.4 - 28.5	1,200	46.9	40.0 - 53.7	1,100	3.0 *	1.0 - 5.0	<100
Private	55.3	50.6 - 60.0	2,700	14.6	9.3 - 19.9	300	97.0 *	95.0 - 99.0	2,300
Maternal postpartum insurance									
Medi-Cal	30.6	26.1 - 35.0	1,500	60.9	54.3 - 67.6	1,400	2.9 *	0.8 - 5.0	<100
Uninsured	15.8	12.5 - 19.1	800	31.6	25.3 - 37.9	700			
Private	51.0	46.2 - 55.8	2,500	7.3	3.7 - 10.8	200	96.4 *	94.1 - 98.6	2,300
Infant health insurance									
Medi-Cal	48.2	43.5 - 53.0	2,300	92.9	89.2 - 96.5	2,100	8.9	5.3 - 12.5	200
Uninsured	1.1 *	0.2 - 1.9	<100						
Private	47.1	42.3 - 51.9	2,300	4.2 *	1.3 - 7.0	<100	89.4	85.5 - 93.3	2,100
Maternal Demographics									
Age									
15-19	3.8	2.1 - 5.5	200	7.5	4.0 - 11.0	200			
20-34	74.0	69.5 - 78.5	3,700	75.3	69.2 - 81.4	1,700	75.1	68.5 - 81.7	1,800
35+	22.2	17.8 - 26.5	1,100	17.2	11.8 - 22.7	400	24.6	18.0 - 31.2	600
Did not complete high school (or GED)	14.9	11.6 - 18.3	700	29.4	23.0 - 35.8	700	2.5 *	0.5 - 4.4	<100
Unmarried	33.2	28.9 - 37.5	1,600	51.7	44.8 - 58.6	1,200	16.9	12.2 - 21.5	400
Race/Ethnicity									
White	51.7	46.8 - 56.5	2,300	28.4	21.8 - 35.0	600	76.0	69.8 - 82.2	1,600
Hispanic	40.8	36.0 - 45.5	1,800	66.9	60.1 - 73.6	1,500	14.5	9.8 - 19.1	300
Asian/Pacific Islander	5.1	2.4 - 7.7	200				8.0 *	3.1 - 12.8	200
Black	1.4	0.6 - 2.2	<100						
Born outside the U.S.	30.6	26.2 - 35.0	1,500	47.9	41.0 - 54.8	1,100	15.4	10.0 - 20.7	400
Speaks non-English language at home	24.8	20.8 - 28.8	1,200	41.8	35.1 - 48.6	1,000	9.2	5.0 - 13.4	200
Lives in a high poverty neighborhood	17.0	13.6 - 20.3	800	21.5	16.1 - 26.8	500	13.1	8.7 - 17.4	300
Income as a percent of the Federal Poverty Guideline	-			-			-		
0-100%	27.2	22.8 - 31.5	1,300	56.4	49.2 - 63.7	1,200	3.1 *	0.7 - 5.6	<100
101-200%	24.0	19.9 - 28.2	1,100	38.1	30.9 - 45.3	800	10.8	6.8 - 14.7	300
> 200%	48.8	44.0 - 53.7	2,300	5.4	2.6 - 8.3	100	86.1	81.6 - 90.6	2,000

-- Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

* Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%)

⁺ Measure/definition may not be comparable to previous Snapshots (see Technical Notes).

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Sonoma County was 557. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.



MIHA Data Snapshot, Sonoma County by Race/Ethnicity, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Son	oma Coun	ty	Hispanic	Black			White			Asian/Pacific Islander			
	%	95% CI	N	% 95% CI	Ν	%	95% CI	Ν	%	95% CI	N	%	95% CI	Ν
Total	100		4,900	40.8 36.0 - 45.5	1,800	1.4	0.6 - 2.2	<100	51.7	46.8 - 56.5	2,300	5.1	2.4 - 7.7	200
Health Status before Pregnancy														
In good to excellent health	93.5 9	1.2 - 95.8	4,600	90.4 86.2 - 94.6	1,700				95.0	* 91.9 - 98.0	2,200			
Chronic conditions														
Diabetes ⁺	2.2	1.0 - 3.4	100	3.8 * 1.5 - 6.1	<100									
Hypertension ⁺	3.3	1.9 - 4.8	200	3.6 * 1.2 - 6.0	<100				2.6	* 0.7 - 4.5	<100			
Asthma ⁺	10.7	7.9 - 13.5	500	9.5 5.3 - 13.7	200				11.4	7.6 - 15.3	300			
Nutrition and Weight														
Daily folic acid use, month before pregnancy	38.7 3	4.0 - 43.4	1,900	32.0 24.9 - 39.1	600				43.8	37.1 - 50.4	1,000	64.6	38.9 - 90.3	100
Overweight before pregnancy	24.0 1	9.9 - 28.1	1,100	27.8 20.8 - 34.7	400				23.2	17.5 - 29.0	500			
Obese before pregnancy	21.7 1	7.5 - 25.9	1,000	29.2 21.7 - 36.8	500				18.1	13.0 - 23.2	400			
Inadequate weight gain during pregnancy	14.6 1	1.1 - 18.0	600	26.5 18.9 - 34.0	400				7.2	4.1 - 10.3	200			
Excessive weight gain during pregnancy	48.5 4	3.3 - 53.7	2,000	42.3 33.9 - 50.6	600				58.8	51.8 - 65.7	1,200			
Food insecurity during pregnancy	14.7 1	1.2 - 18.2	700	21.6 15.5 - 27.8	400				11.2	6.2 - 16.2	300			
Intimate Partner Violence (IPV) and Depressive Symptoms	;													
Physical or psychological IPV during pregnancy	6.0	3.5 - 8.6	300	8.1 3.8 - 12.4	100				5.3	* 1.4 - 9.3	100			
Prenatal depressive symptoms	14.1 1	0.8 - 17.5	700	15.2 10.0 - 20.4	300				13.8	8.7 - 19.0	300			
Postpartum depressive symptoms	13.7 1	0.5 - 16.9	700	13.1 8.3 - 17.9	200				13.1	8.3 - 17.9	300			
Hardships and Support during Pregnancy														
Experienced two or more hardships during childhood	27.2 2	3.0 - 31.4	1,300	27.8 21.1 - 34.5	500				27.2	21.0 - 33.3	600			
Homeless or did not have a regular place to sleep		1.6 - 5.7	200	3.6 * 0.9 - 6.4	<100									
Moved due to problems paying rent or mortgage	-	4.0 - 8.9	300	6.8 * 2.8 - 10.7	100				7.2	3.2 - 11.2	200			
Woman or partner lost job	13.0	9.6 - 16.4	600	15.8 10.3 - 21.3	300				12.3	7.0 - 17.5	300			
Woman or partner had pay or hours cut back	14.4 1	0.7 - 18.0	700	11.7 7.1 - 16.3	200				16.3	10.4 - 22.2	400			
Became separated or divorced		4.4 - 9.7	300	8.9 4.6 - 13.2	200					* 2.3 - 10.9	200			
Had no practical or emotional support	4.5	1.9 - 7.0	200	7.7 * 2.6 - 12.8	100									
Substance Use														
Any smoking, 3 months before pregnancy	13.7 1	0.4 - 17.1	700	6.5 2.9 - 10.1	100				20.1	14.4 - 25.9	500			
Any smoking, 3rd trimester [†]		1.2 - 5.0	200							* 0.6 - 7.2	<100			
Any smoking, postpartum	5.3	3.0 - 7.5	300	3.1 * 0.3 - 5.9	<100					3.3 - 10.8	200			
Any binge drinking, 3 months before pregnancy		4.6 - 22.2	900	11.9 6.9 - 16.9	200					17.3 - 29.3	500			
Any alcohol use, 3rd trimester ⁺	10.0	7.3 - 12.7	500	3.0 * 0.6 - 5.5	<100				16.5	11.7 - 21.4	400			
Pregnancy Intention and Family Planning														
Mistimed or unwanted pregnancy	26.1 2	1.9 - 30.4	1,300	32.0 25.0 - 39.1	600				23.1	16.8 - 29.3	500			
Unsure of pregnancy intentions	-	8.3 - 13.8	500	12.3 8.0 - 16.5	200					5.1 - 12.5	200			
Postpartum birth control use ⁺	-	4.0 - 90.2	4,300	87.0 81.8 - 92.2	1,600					85.4 - 93.3	2,100	77.1	55.2 - 98.9	200
Infant Sleep and Breastfeeding														
Placed infant on back to sleep	8378	0.1 - 87.3	4,000	81.5 75.6 - 87.3	1,500				83.8	78.6 - 89.0	1,900			
Infant always or often shared bed		5.0 - 44.4	1,900	51.8 44.3 - 59.3	900					25.1 - 37.9	700	27.0	5.4 - 48.7	<100
Intended to breastfeed, before birth		3.5 - 97.4	4,600	95.5 * 92.1 - 98.8	1,700					92.2 - 97.7	2,200	27.0	510.7	.100
Intended to breastfeed exclusively, before birth		7.3 - 84.7	3,900	74.0 67.5 - 80.4	1,400					81.9 - 91.0	2,000		45.9 - 93.4	200
Breastfeeding practices	01.0 /		5,500		1,.00				00.4		2,000	00.0		200
Any breastfeeding, 1 month after delivery	92.3 8	9.9 - 94.7	4,500	92.5 88.9 - 96.1	1,700				92.6	89.0 - 96.1	2,100			
Exclusive breastfeeding, 1 month after delivery		8.9 - 68.2	3,000	59.1 51.7 - 66.6	1,000					64.2 - 76.6	1,600		10.2 - 63.0	<100
Any breastfeeding, 3 months after delivery		8.6 - 87.1	3,000	83.9 78.0 - 89.7	1,200				-	76.0 - 88.8	1,400			
Exclusive breastfeeding, 3 months after delivery		3.4 - 54.6	1,800	43.9 35.7 - 52.2	600					47.2 - 63.3	900			



MIHA Data Snapshot, Sonoma County by Race/Ethnicity, 2013-2015

Maternal and Infant Health Assessment (MIHA) Survey

	Sonoma County			Hispanic			Black			White			Asian/Pacific Islander		
	% 95%	CI N	%	95% CI	N	%	95% CI	Ν	%	95% CI	N	%	95% CI	Ν	
Service Utilization and Health Insurance Coverage															
Had a usual source of pre-pregnancy care	80.9 77.1 -	34.6 4,000	77.8	71.8 - 83.8	1,400				79.8	74.1 - 85.6	1,900				
Initiated prenatal care in 1st trimester	92.3 89.9 -	94.7 4,500	87.8	82.8 - 92.9	1,600				95.8	93.6 - 98.1	2,200				
Mom or infant needed but couldn't afford care postpartum	8.5 5.6 -	11.5 400	6.7	3.4 - 10.0	100				10.2	5.2 - 15.2	200				
Participated in WIC during pregnancy ⁺	41.9 37.2 -	46.6 2,100	81.3	75.9 - 86.7	1,500				18.0	12.4 - 23.6	400				
Received CalFresh (food stamps) during pregnancy	19.3 15.6 -	23.0 1,000	30.0	23.4 - 36.6	500				11.5	6.7 - 16.3	300				
Pre-pregnancy insurance															
Medi-Cal	17.4 14.0 -	20.9 900	28.6	21.8 - 35.4	500				9.3	5.3 - 13.4	200				
Uninsured	24.4 20.4 -	28.5 1,200	44.8	37.4 - 52.2	800				14.4	9.4 - 19.3	300				
Private	55.3 50.6 -	50.0 2,700	23.9	17.6 - 30.3	400				74.4	68.1 - 80.6	1,700	71.3 *	47.7 - 94.8	200	
Prenatal insurance															
Medi-Cal	47.1 42.4 -	51.9 2,300	81.0	75.6 - 86.4	1,500				27.3	20.9 - 33.7	600				
Uninsured	1.1 * 0.3 -	1.8 <100													
Private	48.5 43.8 -	53.3 2,400	16.2	11.1 - 21.3	300				67.6	60.9 - 74.3	1,600	71.9*	48.4 - 95.5	200	
Maternal postpartum insurance															
Medi-Cal	30.6 26.1 -	35.0 1,500	46.3	38.8 - 53.7	800				19.9	13.8 - 26.0	500				
Uninsured	15.8 12.5 -	19.1 800	35.2	28.0 - 42.3	600				5.8	3.1 - 8.5	100				
Private	51.0 46.2 -	55.8 2,500	17.8	12.2 - 23.4	300				70.9	64.3 - 77.5	1,700	71.9*	48.4 - 95.5	200	
Infant health insurance															
Medi-Cal	48.2 43.5 -	53.0 2,300	84.0	78.8 - 89.3	1,500				25.4	19.2 - 31.6	600				
Uninsured	1.1 * 0.2 -	1.9 <100							1.6	* 0.2 - 3.0	<100				
Private	47.1 42.3 -	51.9 2,300	11.6	7.1 - 16.0	200				69.8	63.2 - 76.4	1,600	71.9*	48.4 - 95.5	200	
Maternal Demographics															
Age															
15-19	3.8 2.1 -	5.5 200	8.2	4.0 - 12.5	200				1.5	* 0.3 - 2.8	<100				
20-34	74.0 69.5 -	78.5 3,700	73.9	67.1 - 80.6	1,400				73.9	67.6 - 80.2	1,700	70.0*	42.9 - 97.2	200	
35+	22.2 17.8 -	26.5 1,100	17.9	11.9 - 23.9	300				24.6	18.3 - 30.9	600				
Did not complete high school (or GED)	14.9 11.6 -	18.3 700	29.8	22.9 - 36.8	500				3.7	* 0.8 - 6.5	<100				
Unmarried	33.2 28.9 -	37.5 1,600	44.5	37.1 - 51.8	800				26.8	21.0 - 32.7	600				
Born outside the U.S.	30.6 26.2 -		63.1	56.0 - 70.1	1,200				4.2	* 0.9 - 7.5	<100	66.3 *	42.2 - 90.4	200	
Speaks non-English language at home	24.8 20.8 -	28.8 1,200	54.4	47.0 - 61.8	1,000							39.3 *	12.5 - 66.0	<10	
Lives in a high poverty neighborhood	17.0 13.6 -	20.3 800	23.8	17.7 - 29.9	400				11.1	7.2 - 15.1	300				
Income as a percent of the Federal Poverty Guideline			_												
0-100%	27.2 22.8 -	31.5 1,300	48.6	40.7 - 56.4	800				13.5	8.2 - 18.8	300	29.4 *	5.0 - 53.9	<10	
101-200%	24.0 19.9 -		35.9	28.2 - 43.5	600					10.5 - 20.5	400				
> 200%	48.8 44.0 -			10.0 - 21.1	300					64.5 - 77.5	1,600	68.0*	43.3 - 92.7	200	

-- Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

* Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

⁺ Measure/definition may not be comparable to previous Snapshots (see Technical Notes).

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Sonoma County was 557. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.



MIHA Data Snapshot, Sonoma County by Total Live Births, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

		Sonoma County	,		First Live Birth		Second Live Birth or More			
	%	95% CI	Ν	%	95% CI	N	%	95% CI	Ν	
Total	100		4,900	41.0	36.3 - 45.6	2,000	59.0	54.4 - 63.7	2,900	
Health Status before Pregnancy										
In good to excellent health	93.5	91.2 - 95.8	4,600	93.8	90.6 - 96.9	1,900	93.3	90.1 - 96.4	2,700	
Chronic conditions										
Diabetes [†]	2.2	1.0 - 3.4	100	2.2 *	0.3 - 4.1	<100	2.2 *	0.8 - 3.7	<100	
Hypertension ⁺	3.3	1.9 - 4.8	200	3.6 *	1.3 - 5.8	<100	3.2 *	1.2 - 5.1	<100	
Asthma ⁺	10.7	7.9 - 13.5	500	14.3	9.7 - 18.9	300	8.2	4.7 - 11.7	200	
Nutrition and Weight										
Daily folic acid use, month before pregnancy	38.7	34.0 - 43.4	1,900	46.1	38.8 - 53.4	900	33.6	27.6 - 39.6	1,000	
Overweight before pregnancy	24.0	19.9 - 28.1	1,100	18.8	13.6 - 24.1	400	27.9	22.0 - 33.8	700	
Obese before pregnancy	21.7	17.5 - 25.9	1,000	17.9	11.6 - 24.1	400	24.6	18.9 - 30.3	600	
Inadequate weight gain during pregnancy	14.6	11.1 - 18.0	600	10.9	6.3 - 15.5	200	17.6	12.6 - 22.5	400	
Excessive weight gain during pregnancy	48.5	43.3 - 53.7	2,000	52.5	44.8 - 60.2	1,000	45.3	38.3 - 52.4	1,000	
Food insecurity during pregnancy	14.7	11.2 - 18.2	700	12.7	7.2 - 18.2	300	16.1	11.5 - 20.6	500	
Intimate Partner Violence (IPV) and Depressive Symptoms										
Physical or psychological IPV during pregnancy	6.0	3.5 - 8.6	300	7.4 *	2.5 - 12.4	100	5.0	2.5 - 7.5	100	
Prenatal depressive symptoms	14.1	10.8 - 17.5	700	15.6	10.0 - 21.3	300	12.9	8.9 - 17.0	400	
Postpartum depressive symptoms	13.7	10.5 - 16.9	700	12.2	7.5 - 17.0	200	14.5	10.2 - 18.8	400	
Hardships and Support during Pregnancy										
Experienced two or more hardships during childhood	27.2	23.0 - 31.4	1,300	27.5	21.0 - 34.1	600	27.0	21.5 - 32.5	800	
Homeless or did not have a regular place to sleep	3.7	1.6 - 5.7	200	4.7 *	0.8 - 8.6	<100	3.0 *	0.8 - 5.1	<100	
Moved due to problems paying rent or mortgage	6.5	4.0 - 8.9	300	6.5 *	2.4 - 10.5	100	6.5	3.4 - 9.6	200	
Woman or partner lost job	13.0	9.6 - 16.4	600	15.8	10.0 - 21.6	300	11.1	7.1 - 15.2	300	
Woman or partner had pay or hours cut back	14.4	10.7 - 18.0	700	14.0	8.4 - 19.7	300	14.6	9.9 - 19.4	400	
Became separated or divorced	7.1	4.4 - 9.7	300	8.2 *	3.3 - 13.1	200	6.2	3.2 - 9.1	200	
Had no practical or emotional support	4.5	1.9 - 7.0	200				6.0 *	2.3 - 9.7	200	
Substance Use										
Any smoking, 3 months before pregnancy	13.7	10.4 - 17.1	700	16.6	10.7 - 22.4	300	11.8	7.8 - 15.8	300	
Any smoking, 3rd trimester [†]	3.1 *	1.2 - 5.0	200	3.6 *	0.2 - 7.1	<100	2.7 *	0.6 - 4.8	<100	
Any smoking, postpartum	5.3	3.0 - 7.5	300	6.3 *	2.4 - 10.2	100	4.6	1.9 - 7.2	100	
Any binge drinking, 3 months before pregnancy	18.4	14.6 - 22.2	900	25.2	18.6 - 31.9	500	13.8	9.4 - 18.2	400	
Any alcohol use, 3rd trimester [†]	10.0	7.3 - 12.7	500	10.6	6.7 - 14.5	200	9.5	5.9 - 13.1	300	
Pregnancy Intention and Family Planning										
Mistimed or unwanted pregnancy	26.1	21.9 - 30.4	1,300	28.7	21.8 - 35.7	600	24.4	19.0 - 29.7	700	
Unsure of pregnancy intentions	11.0	8.3 - 13.8	500	11.8	7.5 - 16.0	200	10.5	6.9 - 14.1	300	
Postpartum birth control use ⁺	87.1	84.0 - 90.2	4,300	86.1	81.6 - 90.7	1,700	87.8	83.5 - 92.0	2,500	
Infant Sleep and Breastfeeding										
Placed infant on back to sleep	83.7	80.1 - 87.3	4,000	81.0	75.2 - 86.9	1,600	85.7	81.3 - 90.2	2,400	
Infant always or often shared bed	39.7	35.0 - 44.4	1,900	36.7	29.9 - 43.6	700	41.7	35.4 - 48.0	1,200	
Intended to breastfeed, before birth	95.4	93.5 - 97.4	4,600	97.2 *	95.1 - 99.3	1,900	94.1	91.1 - 97.1	2,700	
Intended to breastfeed exclusively, before birth	81.0	77.3 - 84.7	3,900	89.2	85.2 - 93.2	1,800	75.2	69.8 - 80.6	2,200	
Breastfeeding practices										
Any breastfeeding, 1 month after delivery	92.3	89.9 - 94.7	4,500	92.9	89.8 - 96.1	1,900	91.9	88.4 - 95.4	2,600	
Exclusive breastfeeding, 1 month after delivery	63.5	58.9 - 68.2	3,000	60.4	53.1 - 67.6	1,200	65.7	59.7 - 71.8	1,800	
Any breastfeeding, 3 months after delivery	82.8	78.6 - 87.1	3,000	81.2	74.6 - 87.8	1,200	83.9	78.4 - 89.4	1,800	
Exclusive breastfeeding, 3 months after delivery	49.0	43.4 - 54.6	1,800	49.2	40.4 - 58.1	700	48.7	41.4 - 56.0	1,000	



MIHA Data Snapshot, Sonoma County by Total Live Births, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

	Sonoma County					First Live Birth		Second Live Birth or More			
	%	95% CI	N	-	%	95% CI	N	%	95% CI	N	
Service Utilization and Health Insurance Coverage											
Had a usual source of pre-pregnancy care	80.9	77.1 - 84.6	4,000		71.1	64.3 - 78.0	1,400	87.8	84.0 - 91.5	2,600	
Initiated prenatal care in 1st trimester	92.3	89.9 - 94.7	4,500		92.1	88.6 - 95.6	1,800	92.5	89.2 - 95.7	2,600	
Mom or infant needed but couldn't afford care postpartum	8.5	5.6 - 11.5	400		10.1	4.7 - 15.5	200	7.4	4.1 - 10.7	200	
Participated in WIC during pregnancy [†]	41.9	37.2 - 46.6	2,100		33.9	26.9 - 40.9	700	47.6	41.4 - 53.9	1,400	
Received CalFresh (food stamps) during pregnancy	19.3	15.6 - 23.0	1,000		14.0	8.6 - 19.4	300	23.1	18.0 - 28.1	700	
Pre-pregnancy insurance											
Medi-Cal	17.4	14.0 - 20.9	900		10.8	6.7 - 14.9	200	22.1	17.0 - 27.2	600	
Uninsured	24.4	20.4 - 28.5	1,200		26.7	20.2 - 33.2	500	22.7	17.6 - 27.8	700	
Private	55.3	50.6 - 60.0	2,700		60.5	53.4 - 67.5	1,200	51.8	45.5 - 58.1	1,500	
Prenatal insurance											
Medi-Cal	47.1	42.4 - 51.9	2,300		40.6	33.4 - 47.8	800	51.7	45.4 - 57.9	1,500	
Uninsured	1.1 *	0.3 - 1.8	<100								
Private	48.5	43.8 - 53.3	2,400		55.0	47.7 - 62.2	1,100	44.2	37.9 - 50.4	1,300	
Maternal postpartum insurance											
Medi-Cal	30.6	26.1 - 35.0	1,500		32.0	25.0 - 39.0	600	29.5	23.7 - 35.2	900	
Uninsured	15.8	12.5 - 19.1	800		9.6	5.5 - 13.6	200	20.2	15.3 - 25.0	600	
Private	51.0	46.2 - 55.8	2,500		56.3	49.0 - 63.5	1,100	47.4	41.1 - 53.7	1,400	
Infant health insurance											
Medi-Cal	48.2	43.5 - 53.0	2,300		43.6	36.3 - 50.8	900	51.5	45.1 - 57.8	1,500	
Uninsured	1.1 *	0.2 - 1.9	<100								
Private	47.1	42.3 - 51.9	2,300		53.5	46.2 - 60.7	1,100	42.6	36.4 - 48.9	1,200	
Maternal Demographics											
Age											
15-19	3.8	2.1 - 5.5	200		7.4	3.8 - 11.1	200				
20-34	74.0	69.5 - 78.5	3,700		78.6	71.8 - 85.4	1,600	71.0	65.0 - 76.9	2,100	
35+	22.2	17.8 - 26.5	1,100		14.0	7.7 - 20.3	300	27.7	21.8 - 33.6	800	
Did not complete high school (or GED)	14.9	11.6 - 18.3	700		7.0	3.5 - 10.5	100	20.4	15.4 - 25.4	600	
Unmarried	33.2	28.9 - 37.5	1,600		42.7	35.7 - 49.8	800	26.8	21.5 - 32.0	800	
Race/Ethnicity											
White	51.7	46.8 - 56.5	2,300		61.3	54.1 - 68.5	1,100	45.3	38.8 - 51.7	1,200	
Hispanic	40.8	36.0 - 45.5	1,800		31.3	24.6 - 37.9	600	47.1	40.7 - 53.5	1,300	
Asian/Pacific Islander	5.1	2.4 - 7.7	200		5.0 *	0.8 - 9.2	<100	5.1	* 1.7 - 8.5	100	
Black	1.4	0.6 - 2.2	<100								
Born outside the U.S.	30.6	26.2 - 35.0	1,500		20.8	14.7 - 26.8	400	37.4	31.4 - 43.5	1,100	
Speaks non-English language at home	24.8	20.8 - 28.8	1,200		14.3	9.3 - 19.4	300	32.1	26.4 - 37.8	900	
Lives in a high poverty neighborhood	17.0	13.6 - 20.3	800		13.2	8.8 - 17.7	300	19.6	14.9 - 24.4	600	
Income as a percent of the Federal Poverty Guideline											
0-100%	27.2	22.8 - 31.5	1,300		22.1	15.7 - 28.5	400	30.7	24.8 - 36.6	900	
101-200%	24.0	19.9 - 28.2	1,100		21.2	15.1 - 27.2	400	26.0	20.4 - 31.6	700	
> 200%	48.8	44.0 - 53.7	2,300		56.7	49.3 - 64.2	1,100	43.3	37.0 - 49.6	1,200	

-- Estimate not shown because the relative standard error (RSE) is greater than 50%, fewer than 5 women reported, or the weighted population denominator for the column is less than 100 women.

* Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%)

⁺ Measure/definition may not be comparable to previous Snapshots (see Technical Notes).

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. The sample size of Sonoma County was 557. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.