

MIHA Data Snapshot, California by Maternal Age, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

| | California | | 15-19 Years | | 20-34 Years | 35 Years or Older | |
|---|-------------------------|---------|-------------------------|--------|-------------------------------|-----------------------------------|--|
| | % 95% CI | Ν | % 95% CI | Ν | % 95% CI N | % 95% CI N | |
| Total | 100 | 487,900 | 5.4 4.8 - 6.0 | 26,200 | 74.4 73.2 - 75.7 363,2 | 00 20.2 19.1 - 21.3 98,500 | |
| Health Status before Pregnancy | | | | | | | |
| In good to excellent health | 92.0 91.2 - 92.7 | 448,000 | 87.7 83.6 - 91.8 | 23,000 | 91.9 91.0 - 92.8 333,3 | 00 93.3 92.0 - 94.7 91,700 | |
| Chronic conditions | | | | | | | |
| Diabetes ⁺ | 2.5 2.1 - 3.0 | 12,200 | | | 2.1 1.7 - 2.6 7,7 | 00 4.1 2.9 - 5.3 4,000 | |
| Hypertension [†] | 2.6 2.3 - 3.0 | 12,600 | 1.3 * 0.2 - 2.3 | 300 | 2.1 1.8 - 2.5 7,7 | 00 4.8 3.6 - 6.1 4,600 | |
| Asthma† | 8.4 7.6 - 9.1 | 40,000 | 13.8 9.9 - 17.7 | 3,500 | 8.4 7.6 - 9.2 30,0 | 00 6.8 5.4 - 8.2 6,500 | |
| Nutrition and Weight | | | | | | | |
| Daily folic acid use, month before pregnancy | 33.8 32.5 - 35.0 | 164,400 | 15.8 12.4 - 19.1 | 4,100 | 32.2 30.7 - 33.6 116,6 | 00 44.5 41.3 - 47.6 43,700 | |
| Overweight before pregnancy | 25.1 23.8 - 26.3 | 116,100 | 22.2 17.1 - 27.2 | 5,600 | 25.9 24.4 - 27.3 89,2 | 00 22.8 20.2 - 25.4 21,300 | |
| Obese before pregnancy | 20.1 19.0 - 21.2 | 93,000 | 12.0 8.0 - 16.1 | 3,100 | 20.3 19.1 - 21.6 70,1 | 00 21.3 18.7 - 23.8 19,800 | |
| Inadequate weight gain during pregnancy | 18.2 17.0 - 19.4 | 74,300 | 17.3 12.5 - 22.0 | 3,900 | 18.6 17.2 - 20.0 56,8 | 00 17.0 14.3 - 19.6 13,700 | |
| Excessive weight gain during pregnancy | 41.2 39.7 - 42.6 | 168,100 | 49.2 42.7 - 55.7 | 11,100 | 40.4 38.7 - 42.1 123,3 | 00 41.8 38.3 - 45.3 33,700 | |
| Food insecurity during pregnancy | 15.6 14.6 - 16.5 | 75,100 | 22.2 17.2 - 27.2 | 5,700 | 16.3 15.2 - 17.5 58,6 | 00 11.1 9.3 - 12.9 10,800 | |
| Intimate Partner Violence (IPV) and Depressive Symptoms | | | | | | | |
| Physical or psychological IPV during pregnancy | 7.0 6.3 - 7.7 | 34,000 | 15.3 11.8 - 18.8 | 4,000 | 7.1 6.3 - 7.9 25,6 | 00 4.6 3.4 - 5.9 4,500 | |
| Prenatal depressive symptoms | 14.1 13.1 - 15.0 | 68,500 | 18.3 14.0 - 22.5 | 4,800 | 14.1 13.0 - 15.2 51,1 | 00 12.8 10.7 - 15.0 12,600 | |
| Postpartum depressive symptoms | 13.5 12.6 - 14.4 | 65,700 | 14.5 10.1 - 18.9 | 3,700 | 13.6 12.5 - 14.6 49,1 | 00 13.1 10.9 - 15.3 12,800 | |
| Hardships and Support during Pregnancy | | | | | | | |
| Experienced two or more hardships during childhood | 24.6 23.4 - 25.7 | 116,800 | 32.6 27.3 - 37.9 | 8,200 | 25.5 24.1 - 26.9 90,1 | 00 19.1 16.8 - 21.4 18,500 | |
| Homeless or did not have a regular place to sleep | 3.0 2.6 - 3.4 | 14,500 | 6.6 3.7 - 9.5 | 1,700 | 3.0 2.5 - 3.5 10,8 | 00 2.1 1.3 - 2.8 2,000 | |
| Moved due to problems paying rent or mortgage | 6.4 5.8 - 7.1 | 31,100 | 8.6 4.9 - 12.2 | 2,200 | 6.8 6.1 - 7.6 24,7 | 00 4.3 3.1 - 5.5 4,200 | |
| Woman or partner lost job | 14.3 13.3 - 15.2 | 68,700 | 20.8 15.5 - 26.0 | 5,200 | 14.4 13.3 - 15.5 51,6 | 00 12.2 10.3 - 14.2 11,900 | |
| Woman or partner had pay or hours cut back | 11.4 10.5 - 12.2 | 54,700 | 10.0 6.1 - 13.9 | 2,500 | 11.7 10.8 - 12.7 42,1 | 00 10.4 8.5 - 12.3 10,100 | |
| Became separated or divorced | 7.0 6.3 - 7.7 | 34,000 | 15.7 11.6 - 19.9 | 4,000 | 6.6 5.9 - 7.4 23,9 | 00 6.3 4.7 - 7.8 6,100 | |
| Had no practical or emotional support | 4.6 4.0 - 5.2 | 22,500 | 4.9 2.5 - 7.2 | 1,300 | 4.5 3.8 - 5.1 16,1 | 00 5.2 3.7 - 6.7 5,100 | |
| Substance Use | | | | | | | |
| Any smoking, 3 months before pregnancy | 10.8 9.9 - 11.6 | 52,100 | 10.5 7.4 - 13.7 | 2,800 | 11.8 10.8 - 12.8 42,4 | 00 7.0 5.6 - 8.5 6,900 | |
| Any smoking, 3rd trimester ⁺ | 2.7 2.3 - 3.1 | 13,200 | 2.5 1.3 - 3.7 | 700 | 3.1 2.6 - 3.7 11,3 | 00 1.2 0.9 - 1.6 1,200 | |
| Any smoking, postpartum | 5.4 4.8 - 6.0 | 26,100 | 6.6 4.0 - 9.1 | 1,700 | 5.9 5.2 - 6.6 21,3 | 00 3.1 2.2 - 4.1 3,100 | |
| Any binge drinking, 3 months before pregnancy | 15.1 14.1 - 16.0 | 72,600 | 10.0 6.4 - 13.6 | 2,600 | 16.3 15.1 - 17.4 58,3 | 00 12.0 10.0 - 14.0 11,700 | |
| Any alcohol use, 3rd trimester [†] | 7.3 6.6 - 8.1 | 35,600 | 0.4 * 0.1 - 0.7 | 100 | 6.9 6.0 - 7.7 24,7 | 00 11.0 9.1 - 12.9 10,800 | |
| Pregnancy Intention and Family Planning | | | | | | | |
| Mistimed or unwanted pregnancy | 30.5 29.3 - 31.8 | 147,900 | 63.4 58.0 - 68.9 | 16,600 | 30.9 29.4 - 32.4 111,5 | 00 20.3 17.7 - 22.9 19,800 | |
| Unsure of pregnancy intentions | 12.8 11.9 - 13.7 | 62,100 | 18.3 14.2 - 22.5 | 4,800 | 12.1 11.1 - 13.1 43,6 | 00 14.0 11.8 - 16.2 13,600 | |
| Postpartum birth control use [†] | 81.0 79.9 - 82.1 | 386,400 | 82.3 77.9 - 86.6 | 20,900 | 82.3 81.0 - 83.5 291,9 | 00 76.1 73.3 - 79.0 73,600 | |
| Infant Sleep and Breastfeeding | | | | | | | |
| Placed infant on back to sleep | 80.3 79.2 - 81.4 | 384,100 | 72.2 66.9 - 77.5 | 18,600 | 79.8 78.5 - 81.2 284,1 | 00 84.2 81.9 - 86.6 81,300 | |
| Infant always or often shared bed | 34.6 33.3 - 35.9 | 165,100 | 38.7 33.1 - 44.3 | 10,000 | 34.4 32.9 - 35.9 122,2 | 00 34.3 31.3 - 37.3 32,900 | |
| Intended to breastfeed, before birth | 92.4 91.7 - 93.2 | 441,900 | 88.7 85.1 - 92.4 | 22,700 | 92.5 91.6 - 93.3 329,0 | 00 93.3 91.7 - 95.0 90,100 | |
| Intended to breastfeed exclusively, before birth | 64.7 63.3 - 66.0 | 309,100 | 58.6 52.9 - 64.2 | 15,000 | 65.1 63.6 - 66.7 231,7 | 00 64.6 61.5 - 67.7 62,400 | |
| Breastfeeding practices | | | | | | | |
| Any breastfeeding, 1 month after delivery | 83.8 82.8 - 84.8 | 397,900 | 70.7 65.4 - 76.0 | 18,200 | 84.0 82.8 - 85.2 296,4 | 00 86.7 84.6 - 88.9 83,400 | |
| Exclusive breastfeeding, 1 month after delivery | 44.2 42.9 - 45.6 | 206,800 | 40.4 34.5 - 46.2 | 10,300 | 46.2 44.6 - 47.8 160,9 | 00 38.1 35.0 - 41.1 35,600 | |
| Any breastfeeding, 3 months after delivery | 67.3 65.8 - 68.8 | 252,100 | 41.8 35.2 - 48.4 | 8,600 | 66.5 64.8 - 68.3 185,5 | 00 77.3 74.2 - 80.3 58,000 | |
| Exclusive breastfeeding, 3 months after delivery | 29.1 27.7 - 30.5 | 107,600 | 16.6 11.8 - 21.5 | 3,400 | 29.9 28.3 - 31.5 82,6 | 00 29.3 26.1 - 32.6 21,600 | |

MIHA is a collaborative effort of the Maternal, Child and Adolescent Health Division and the Women, Infants and Children Division in the California Department of Public Health and the Center on Social Disparities in Health at the University of California San Francisco.



MIHA Data Snapshot, California by Maternal Age, 2013-2015 Maternal and Infant Health Assessment (MIHA) Survey

| | California | | 15-19 Years | | 20-34 Years | | 35 Years or | 35 Years or Older | |
|--|-------------------------|---------|-------------------------|--------|-------------------------|---------|-----------------------|-------------------|--|
| | % 95% CI | Ν | % 95% CI | N | % 95% CI | N | % 95% CI | Ν | |
| Service Utilization and Health Insurance Coverage | | | | | | | | | |
| Had a usual source of pre-pregnancy care | 70.6 69.4 - 71.9 | 343,800 | 64.4 58.8 - 70.0 | 16,900 | 68.7 67.2 - 70.1 | 248,700 | 79.6 77.0 - 82 | .2 78,200 | |
| Initiated prenatal care in 1st trimester | 89.4 88.6 - 90.2 | 428,900 | 77.3 73.0 - 81.6 | 19,800 | 89.8 88.8 - 90.7 | 320,900 | 91.1 89.2 - 92 | .9 88,200 | |
| Mom or infant needed but couldn't afford care postpartum | 11.5 10.6 - 12.4 | 54,200 | 11.8 8.3 - 15.2 | 3,000 | 11.9 10.8 - 13.0 | 41,900 | 9.8 8.0 - 11 | .5 9,200 | |
| Participated in WIC during pregnancy ⁺ | 53.4 52.4 - 54.4 | 254,400 | 92.0 89.7 - 94.3 | 23,800 | 55.7 54.4 - 57.0 | 198,000 | 34.4 31.6 - 37 | .1 32,500 | |
| Received CalFresh (food stamps) during pregnancy | 25.2 24.1 - 26.2 | 121,600 | 38.5 32.8 - 44.2 | 10,000 | 27.1 25.8 - 28.4 | 97,500 | 14.4 12.6 - 16 | .3 14,100 | |
| Pre-pregnancy insurance | | | | | | | | | |
| Medi-Cal | 26.7 25.6 - 27.8 | 129,800 | 62.5 56.9 - 68.0 | 16,400 | 27.2 25.9 - 28.5 | 98,400 | 15.3 13.2 - 17 | .3 15,000 | |
| Uninsured | 21.6 20.5 - 22.7 | 105,000 | 17.5 13.2 - 21.8 | 4,600 | 23.2 21.9 - 24.6 | 84,100 | 16.6 14.2 - 18 | .9 16,300 | |
| Private | 47.8 46.7 - 48.9 | 232,400 | 16.7 12.3 - 21.0 | 4,400 | 45.4 44.0 - 46.7 | 164,200 | 64.9 62.1 - 67 | .8 63,800 | |
| Prenatal insurance | | | | | | | | | |
| Medi-Cal | 50.1 49.0 - 51.1 | 243,600 | 85.7 81.5 - 89.9 | 22,400 | 52.5 51.2 - 53.9 | 190,000 | 31.7 29.0 - 34 | .4 31,200 | |
| Uninsured | 1.6 1.3 - 2.0 | 7,800 | 2.2 * 0.7 - 3.6 | 600 | 1.5 1.1 - 1.9 | 5,300 | 1.9 1.0 - 2 | .8 1,900 | |
| Private | 44.1 43.1 - 45.2 | 214,700 | 10.5 6.6 - 14.4 | 2,800 | 41.5 40.1 - 42.8 | 150,100 | 63.0 60.1 - 65 | .8 61,900 | |
| Maternal postpartum insurance | | | | | | | | | |
| Medi-Cal | 38.4 37.3 - 39.5 | 185,700 | 81.6 77.5 - 85.7 | 21,300 | 40.0 38.6 - 41.4 | 143,800 | 21.0 18.6 - 23 | .4 20,600 | |
| Uninsured | 13.0 12.0 - 13.9 | 62,600 | 7.8 5.4 - 10.2 | 2,000 | 13.5 12.4 - 14.6 | 48,500 | 12.3 10.3 - 14 | .3 12,100 | |
| Private | 45.3 44.2 - 46.3 | 218,800 | 9.1 5.7 - 12.6 | 2,400 | 42.7 41.4 - 44.1 | 153,600 | 64.1 61.3 - 66 | .9 62,900 | |
| Infant health insurance | | | | | | | | | |
| Medi-Cal | 52.8 51.7 - 53.8 | 251,800 | 90.6 87.2 - 94.0 | 23,400 | 55.3 53.9 - 56.6 | 196,200 | 33.4 30.7 - 36 | .2 32,200 | |
| Uninsured | 2.8 2.3 - 3.2 | 13,100 | 3.7 1.9 - 5.6 | 1,000 | 2.8 2.2 - 3.4 | 9,900 | 2.3 1.3 - 3 | .4 2,300 | |
| Private | 40.9 39.8 - 41.9 | 195,000 | 4.5 * 1.6 - 7.3 | 1,100 | 38.0 36.7 - 39.3 | 135,000 | 61.1 58.2 - 64 | .0 58,900 | |
| Maternal Demographics | | | | | | | | | |
| Did not complete high school (or GED) | 16.3 15.3 - 17.3 | 78,500 | 44.1 38.2 - 50.1 | 11,400 | 15.2 14.0 - 16.4 | 54,600 | 12.9 10.9 - 14 | .8 12,600 | |
| Unmarried | 40.5 39.2 - 41.7 | 196,000 | 87.0 82.7 - 91.3 | 22,600 | 41.3 39.8 - 42.7 | 148,700 | 25.2 22.6 - 27 | .8 24,700 | |
| Race/Ethnicity | | | | | | | | | |
| White | 29.3 28.1 - 30.4 | 140,000 | 10.3 7.7 - 12.9 | 2,700 | 29.4 28.1 - 30.7 | 104,700 | 34.0 31.0 - 36 | .9 32,600 | |
| Hispanic | 49.2 48.0 - 50.4 | 235,200 | 79.2 75.6 - 82.9 | 20,600 | 50.0 48.5 - 51.4 | 178,000 | 38.1 35.1 - 41 | .0 36,600 | |
| Asian/Pacific Islander | 15.4 14.3 - 16.5 | 73,800 | 1.4 0.6 - 2.2 | 400 | 14.2 13.0 - 15.5 | 50,700 | 23.6 20.5 - 26 | .7 22,700 | |
| Black | 5.6 5.4 - 5.9 | 26,900 | 8.4 6.0 - 10.8 | 2,200 | 5.8 5.5 - 6.2 | 20,800 | 4.2 3.3 - 5 | .0 4,000 | |
| Born outside the U.S. | 37.7 36.3 - 39.0 | 183,800 | 15.4 11.4 - 19.5 | 4,000 | 35.6 34.1 - 37.2 | 129,400 | 51.1 48.0 - 54 | .2 50,300 | |
| Speaks non-English language at home | 30.7 29.4 - 32.0 | 147,700 | 19.1 14.5 - 23.7 | 5,000 | 29.9 28.4 - 31.4 | 107,200 | 36.7 33.5 - 39 | | |
| Lives in a high poverty neighborhood | 38.8 37.5 - 40.1 | 182,400 | 61.0 55.4 - 66.6 | 15,200 | 40.0 38.5 - 41.6 | 140,000 | 28.5 25.6 - 31 | .3 27,100 | |

-- Estimate not shown because the relative standard error (RSE) is greater than 50%, or fewer than 5 women reported.

* Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%)

⁺ Measure/definition may not be comparable to previous Snapshots (see Technical Notes).

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2015 were combined, resulting in a statewide sample size of 20,762. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California in 2013-2015. Population estimate (N) is a three-year average. See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

Table Citation: Maternal and Infant Health Assessment (MIHA) Survey Data Snapshots, 2013-2015. California Department of Public Health; 2018.