

MIHA Data Snapshot, San Diego County, 2013-2014

Maternal and Infant Health Assessment (MIHA) Survey

✓ better than rest of California

✓ worse than rest of California

◆ no statistical difference

	San Diego County					California		
		%	95% CI	Population Estimate	%	95% CI	Population Estimate	
Health Status Before Pregnancy		,-			,,			
In good to excellent health	•	94.8	92.4 - 97.3	41,000	92.4	91.5 - 93.3	452,800	
Chronic conditions								
Diabetes†	*	4.6 *	1.4 - 7.7	1,900	2.3	1.8 - 2.8	10,900	
Hypertension†	•	3.1 *	1.0 - 5.2	1,300	2.5	2.0 - 2.9	11,900	
Asthma†	•	8.8	4.6 - 13.0	3,700	7.8	7.0 - 8.6	37,500	
Nutrition and Weight								
Daily folic acid use, month before pregnancy	\checkmark	41.9	35.2 - 48.7	18,200	32.7	31.1 - 34.3	159,900	
Overweight before pregnancy	•	23.8	17.8 - 29.8	9,800	24.7	23.2 - 26.2	114,900	
Obese before pregnancy	•	17.6	12.4 - 22.8	7,200	20.3	18.9 - 21.7	94,400	
Inadequate weight gain during pregnancy	•	17.0	10.9 - 23.1	6,100	17.7	16.2 - 19.2	72,900	
Excessive weight gain during pregnancy	•	35.7	28.5 - 42.9	12,900	41.4	39.5 - 43.2	170,100	
Food insecurity during pregnancy	•	15.7	10.6 - 20.7	6,700	16.6	15.4 - 17.9	80,500	
Intimate Partner Violence (IPV) and Depressive Symptoms								
Physical or psychological IPV during pregnancy	•	7.4	3.6 - 11.2	3,200	7.1	6.2 - 7.9	34,500	
Prenatal depressive symptoms	•	12.6	7.9 - 17.3	5,500	14.1	12.9 - 15.3	68,900	
Postpartum depressive symptoms	•	11.1	6.8 - 15.4	4,800	13.0	11.8 - 14.1	63,200	
Hardships and Support during Pregnancy								
Experienced two or more hardships during childhood	•	28.3	22.2 - 34.3	12,000	25.3	23.8 - 26.7	121,600	
Homeless or did not have a regular place to sleep	•	2.8 *	0.6 - 5.1	1,200	2.9	2.4 - 3.4	14,000	
Moved due to problems paying rent or mortgage	•	5.5	2.5 - 8.5	2,400	6.9	6.0 - 7.7	33,600	
Woman or partner lost job	•	11.3	7.4 - 15.3	4,900	14.6	13.4 - 15.8	71,000	
Woman or partner had pay or hours cut back	•	9.7	6.2 - 13.2	4,200	11.8	10.7 - 12.8	57,100	
Became separated or divorced	•	7.4	3.3 - 11.5	3,200	7.6	6.7 - 8.5	37,000	
Had no practical or emotional support	•	2.8 *	0.7 - 4.8	1,200	4.4	3.7 - 5.2	21,700	
Substance Use								
Any smoking, 3 months before pregnancy	•	13.0	8.5 - 17.5	5,600	11.6	10.5 - 12.7	56,500	
Any smoking, 3rd trimester†	•	1.5 *	0.2 - 2.7	600	2.9	2.3 - 3.4	13,900	
Any smoking, postpartum	•	5.8	2.7 - 8.9	2,500	5.6	4.8 - 6.3	27,300	
Any binge drinking, 3 months before pregnancy	•	18.8	13.3 - 24.3	7,900	15.1	13.8 - 16.3	72,900	
Any alcohol use, 3rd trimester†	×	14.9	9.3 - 20.5	6,400	7.6	6.6 - 8.6	37,000	
Pregnancy Intention and Family Planning								
Mistimed or unwanted pregnancy	•	29.8	23.5 - 36.1	12,600	31.2	29.6 - 32.8	151,800	
Unsure of pregnancy intentions	•	13.4	8.5 - 18.4	5,700	12.9	11.8 - 14.0	62,700	
Postpartum birth control use†	•	85.1	80.0 - 90.1	36,200	81.6	80.3 - 83.0	390,200	
Infant Sleep and Breastfeeding								
Placed infant on back to sleep	•	79.9	74.2 - 85.6	33,800	79.7	78.3 - 81.1	383,200	
Infant always or often shared bed	•	28.2	22.2 - 34.1	11,900	34.1	32.5 - 35.7	163,700	
Intended to breastfeed, before birth	•	93.9	90.9 - 96.9	39,800	92.4	91.5 - 93.4	444,100	
Intended to breastfeed exclusively, before birth	•	67.4	60.9 - 73.9	28,500	64.2	62.6 - 65.9	308,500	
Breastfeeding practices								
Any breastfeeding, 1 month after delivery	√	90.7	86.9 - 94.6	38,000	83.5	82.3 - 84.8	398,500	
Exclusive breastfeeding, 1 month after delivery	\checkmark	55.0	48.0 - 62.0	22,800	42.7	41.0 - 44.4	200,700	
Any breastfeeding, 3 months after delivery	•	72.3	65.0 - 79.6	22,900	66.2	64.3 - 68.2	244,200	
Exclusive breastfeeding, 3 months after delivery	√	34.4	26.6 - 42.1	10,800	27.4	25.7 - 29.1	99,800	

MIHA is a collaborative effort of the Maternal, Child and Adolescent Health Division and the Women, Infants and Children Division in the California Department of Public Health and the Center on Social Disparities in Health at the University of California San Francisco.



MIHA Data Snapshot, San Diego County, 2013-2014

Maternal and Infant Health Assessment (MIHA) Survey

↑ higher than rest of California ♦ lower than rest of California ♦ no statistical difference

	San Diego County			California			
				Population			Population
		%	95% CI	Estimate	%	95% CI	Estimate
Health Care Utilization and Public Program Participation							
Had a usual source of pre-pregnancy care	•	75.3	69.7 - 80.9	32,500	70.4	68.8 - 71.9	
Initiated prenatal care in 1st trimester	•	90.3	86.6 - 93.9	38,900	89.7	88.7 - 90.6	
Had a postpartum medical visit	•	90.2	85.7 - 94.6	•	87.5	86.4 - 88.7	
Mom or infant needed but couldn't afford care postpartum	+	12.4	7.6 - 17.2	5,200	12.6	11.4 - 13.7	59,700
Participated in WIC during pregnancy†	Ψ	44.4	39.0 - 49.8	18,900	54.1	52.9 - 55.4	259,300
Received CalFresh (food stamps) during pregnancy	Ψ	18.4	13.4 - 23.4	7,900	24.9	23.6 - 26.3	121,000
Health Insurance Coverage							
Pre-pregnancy insurance							
Medi-Cal	$\mathbf{\Psi}$	17.1	12.5 - 21.7	7,400	24.5	23.1 - 25.8	119,600
Uninsured	•	22.7	17.2 - 28.1	9,800	24.0	22.6 - 25.4	117,500
Private	*	50.4	44.6 - 56.1	21,800	47.5	46.2 - 48.9	232,500
Prenatal insurance	_						
Medi-Cal	$\mathbf{\Psi}$	39.0	34.0 - 44.1	16,900	49.7	48.4 - 51.0	242,800
Uninsured	*	2.4 *	0.3 - 4.6	1,100	1.6	1.2 - 2.0	7,900
Private	•	46.6	40.9 - 52.4	20,100	44.5	43.1 - 45.8	217,100
Maternal postpartum insurance	_						
Medi-Cal	$\mathbf{\Psi}$	25.7	20.5 - 30.9	11,000	36.2	34.7 - 37.6	175,700
Uninsured	•	16.1	11.3 - 20.9	6,900	15.0	13.8 - 16.2	72,800
Private	•	48.6	42.8 - 54.4	20,800	45.7	44.3 - 47.0	221,900
Infant health insurance							
Medi-Cal	$\mathbf{\Psi}$	41.4	35.7 - 47.0	17,500	52.2	50.9 - 53.6	250,400
Uninsured	*	4.8 *	1.9 - 7.8	2,000	2.9	2.2 - 3.5	13,700
Private	*	43.5	37.7 - 49.4	18,400	41.3	40.0 - 42.6	198,000
Maternal Demographics							
Age							
15-19	•	4.0	1.7 - 6.3	1,700	5.6	4.9 - 6.4	27,500
20-34	•	76.1	70.4 - 81.9	33,000	74.5	73.0 - 76.0	365,300
35+	*	19.9	14.4 - 25.3	8,600	19.9	18.5 - 21.3	97,400
Did not complete high school (or GED)	•	11.9	7.4 - 16.4	5,100	16.5	15.2 - 17.7	79,900
Unmarried	*	37.1	31.1 - 43.2	16,000	40.9	39.4 - 42.4	199,000
Race/Ethnicity							
Hispanic	•	43.3	36.7 - 49.9	17,300	49.2	47.7 - 50.7	236,500
Black	•	5.1	3.8 - 6.3	2,000	5.7	5.3 - 6.1	27,400
White	1	38.7	31.8 - 45.6	15,500	29.3	27.9 - 30.7	140,900
Asian/Pacific Islander	•	12.4	7.0 - 17.8	5,000	15.3	13.9 - 16.7	73,700
Nativity and Language							
Born outside the U.S.	•	32.7	26.2 - 39.2	14,200	37.9	36.2 - 39.6	185,700
Speaks non-English language at home	•	26.1	20.0 - 32.2	11,300	30.1	28.5 - 31.7	144,800
Lives in a high poverty neighborhood	$\mathbf{\Psi}$	30.9	24.8 - 37.0		38.9	37.3 - 40.5	
Income as a percent of the Federal Poverty Guideline							•
0-100%	$\mathbf{\Psi}$	34.0	28.2 - 39.9	14,000	41.0	39.5 - 42.4	188,800
101-200%	•	20.2	14.9 - 25.5	8,300	20.8	19.4 - 22.2	96,000
> 200%	^	45.8	40.0 - 51.6		38.2	36.9 - 39.6	•
Estimate not shown because the relative standard error (DSE) is greater than E00				10,500	50.2	30.5 - 33.0	110,200

 $[\]hbox{-- Estimate not shown because the relative standard error (RSE) is greater than 50\% or fewer than 5 women reported.}$

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2014 were combined, resulting in a statewide sample size of 13,963. The sample size of San Diego County was 404. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the region in 2013-2014. Population estimate is a two-year average. Symbols indicate whether the health indicator in the region was statistically different from the rest of the state (p-value < 0.05, chi-square test). See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA. Table Citation: Maternal and Infant Health Assessment (MIHA) Survey Data Snapshots, 2013-2014. California Department of Public Health. 2016.

 $^{^{*}}$ Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

[†] Measure/definition changed for this indicator and is not comparable to prior years (see Technical Notes).