

MIHA Data Snapshot, Santa Cruz County, 2013-2014

Maternal and Infant Health Assessment (MIHA) Survey

✓ better than rest of California

✓ worse than rest of California

◆ no statistical difference

	Santa Cruz County					California		
		%	95% CI	Population Estimate	%	95% CI	Population Estimate	
Health Status Before Pregnancy								
In good to excellent health	•	89.3	85.9 - 92.8	2,600	92.4	91.5 - 93.3	452,800	
Chronic conditions								
Diabetes†	•	1.7 *	0.5 - 3.0	< 100	2.3	1.8 - 2.8	10,900	
Hypertension†	•	2.2 *	0.8 - 3.5	100	2.5	2.0 - 2.9	11,900	
Asthma†	•	8.3	5.5 - 11.1	200	7.8	7.0 - 8.6	37,500	
Nutrition and Weight								
Daily folic acid use, month before pregnancy	•	36.7	31.5 - 41.9	1,100	32.7	31.1 - 34.3	159,900	
Overweight before pregnancy	•	24.6	19.7 - 29.5	600	24.7	23.2 - 26.2	114,900	
Obese before pregnancy	•	20.3	15.9 - 24.7	500	20.3	18.9 - 21.7	94,400	
Inadequate weight gain during pregnancy	•	16.5	12.1 - 20.9	400	17.7	16.2 - 19.2	72,900	
Excessive weight gain during pregnancy	x	48.2	42.2 - 54.1	1,100	41.4	39.5 - 43.2	170,100	
Food insecurity during pregnancy	•	19.2	15.0 - 23.5	600	16.6	15.4 - 17.9	80,500	
Intimate Partner Violence (IPV) and Depressive Symptoms								
Physical or psychological IPV during pregnancy	•	8.0	5.0 - 11.1	200	7.1	6.2 - 7.9	34,500	
Prenatal depressive symptoms	•	15.6	11.7 - 19.5	500	14.1	12.9 - 15.3	68,900	
Postpartum depressive symptoms	•	14.8	11.0 - 18.5	400	13.0	11.8 - 14.1	63,200	
Hardships and Support during Pregnancy								
Experienced two or more hardships during childhood	•	20.7	16.5 - 24.9	600	25.3	23.8 - 26.7	121,600	
Homeless or did not have a regular place to sleep	•	3.8	1.8 - 5.8	100	2.9	2.4 - 3.4	14,000	
Moved due to problems paying rent or mortgage	•	5.0	2.9 - 7.2	100	6.9	6.0 - 7.7	33,600	
Woman or partner lost job	•	12.4	9.1 - 15.8	400	14.6	13.4 - 15.8	71,000	
Woman or partner had pay or hours cut back	•	13.4	9.8 - 17.0	400	11.8	10.7 - 12.8	57,100	
Became separated or divorced	\checkmark	4.5	2.4 - 6.7	100	7.6	6.7 - 8.5	37,000	
Had no practical or emotional support	•	5.2	3.0 - 7.3	200	4.4	3.7 - 5.2	21,700	
Substance Use								
Any smoking, 3 months before pregnancy	\checkmark	6.5	4.1 - 9.0	200	11.6	10.5 - 12.7	56,500	
Any smoking, 3rd trimester†	\checkmark	1.1 *	0.1 - 2.1	< 100	2.9	2.3 - 3.4	13,900	
Any smoking, postpartum	\checkmark	2.3	1.0 - 3.6	100	5.6	4.8 - 6.3	27,300	
Any binge drinking, 3 months before pregnancy	•	18.3	14.3 - 22.4	500	15.1	13.8 - 16.3	72,900	
Any alcohol use, 3rd trimester†	×	10.8	7.7 - 14.0	300	7.6	6.6 - 8.6	37,000	
Pregnancy Intention and Family Planning								
Mistimed or unwanted pregnancy	•	26.3	21.7 - 30.9	800	31.2	29.6 - 32.8	151,800	
Unsure of pregnancy intentions	•	10.9	7.5 - 14.3	300	12.9	11.8 - 14.0	62,700	
Postpartum birth control use†	•	84.2	80.4 - 88.0	2,400	81.6	80.3 - 83.0	390,200	
Infant Sleep and Breastfeeding								
Placed infant on back to sleep	•	83.2	79.2 - 87.2	2,400	79.7	78.3 - 81.1	383,200	
Infant always or often shared bed	×	48.1	42.8 - 53.4	1,400	34.1	32.5 - 35.7	163,700	
Intended to breastfeed, before birth	\checkmark	97.3 *	95.6 - 99.0	2,800	92.4	91.5 - 93.4	444,100	
Intended to breastfeed exclusively, before birth	\checkmark	73.6	68.9 - 78.2	2,100	64.2	62.6 - 65.9	308,500	
Breastfeeding practices								
Any breastfeeding, 1 month after delivery	\checkmark	93.9	91.1 - 96.6	2,700	83.5	82.3 - 84.8	398,500	
Exclusive breastfeeding, 1 month after delivery	\checkmark	57.7	52.4 - 63.1	1,600	42.7	41.0 - 44.4	200,700	
Any breastfeeding, 3 months after delivery	\checkmark	82.5	77.7 - 87.3	1,700	66.2	64.3 - 68.2	244,200	
Exclusive breastfeeding, 3 months after delivery	\checkmark	40.6	34.5 - 46.7	900	27.4	25.7 - 29.1	99,800	

MIHA is a collaborative effort of the Maternal, Child and Adolescent Health Division and the Women, Infants and Children Division in the California Department of Public Health and the Center on Social Disparities in Health at the University of California San Francisco.



MIHA Data Snapshot, Santa Cruz County, 2013-2014

Maternal and Infant Health Assessment (MIHA) Survey

↑ higher than rest of California ♦ lower than rest of California ♦ no statistical difference

A lighter than rest of camorina + lower than rest of camorina + no		Santa Cruz County				California		
			Population		Por		Population	
		%	95% CI	Estimate	%	95% CI	Estimate	
Health Care Utilization and Public Program Participation								
Had a usual source of pre-pregnancy care	1	78.5	74.2 - 82.7	2,300	70.4	68.8 - 71.9	344,100	
Initiated prenatal care in 1st trimester	•	91.4	88.5 - 94.2	2,600	89.7	88.7 - 90.6	433,500	
Had a postpartum medical visit	•	88.1	84.8 - 91.4	2,500	87.5	86.4 - 88.7	427,500	
Mom or infant needed but couldn't afford care postpartum	•	16.1	12.2 - 20.0	500	12.6	11.4 - 13.7	59,700	
Participated in WIC during pregnancy†	•	53.2	47.9 - 58.5	1,500	54.1	52.9 - 55.4	259,300	
Received CalFresh (food stamps) during pregnancy	•	26.8	22.2 - 31.4	800	24.9	23.6 - 26.3	121,000	
Health Insurance Coverage								
Pre-pregnancy insurance								
Medi-Cal	•	22.6	18.3 - 26.9	700	24.5	23.1 - 25.8	119,600	
Uninsured	•	24.5	20.2 - 28.7	700	24.0	22.6 - 25.4	117,500	
Private	•	49.9	44.6 - 55.2	1,400	47.5	46.2 - 48.9	232,500	
Prenatal insurance								
Medi-Cal	1	55.4	50.1 - 60.7	1,600	49.7	48.4 - 51.0	242,800	
Uninsured	•	1.2 *	0.1 - 2.3	< 100	1.6	1.2 - 2.0	7,900	
Private	•	42.3	37.0 - 47.6	1,200	44.5	43.1 - 45.8	217,100	
Maternal postpartum insurance								
Medi-Cal	•	35.9	30.9 - 41.0	1,000	36.2	34.7 - 37.6	175,700	
Uninsured	1	20.3	16.3 - 24.3	600	15.0	13.8 - 16.2	72,800	
Private	*	43.4	38.1 - 48.6	1,300	45.7	44.3 - 47.0	221,900	
Infant health insurance								
Medi-Cal	1	58.0	52.8 - 63.3	1,700	52.2	50.9 - 53.6	250,400	
Uninsured					2.9	2.2 - 3.5	13,700	
Private	•	37.9	32.7 - 43.1	1,100	41.3	40.0 - 42.6	198,000	
Maternal Demographics								
Age								
15-19	•	5.8	3.2 - 8.3	200	5.6	4.9 - 6.4	27,500	
20-34	•	70.6	65.8 - 75.4	2,100	74.5	73.0 - 76.0	365,300	
35+	•	23.6	19.2 - 28.1	700	19.9	18.5 - 21.3	97,400	
Did not complete high school (or GED)	1	22.2	17.9 - 26.5	600	16.5	15.2 - 17.7	79,900	
Unmarried	*	40.5	35.4 - 45.7	1,200	40.9	39.4 - 42.4	199,000	
Race/Ethnicity								
Hispanic	1	56.8	51.6 - 62.1	1,600	49.2	47.7 - 50.7	236,500	
Black	-				5.7	5.3 - 6.1	27,400	
White	1	39.0	33.9 - 44.2	1,100	29.3	27.9 - 30.7		
Asian/Pacific Islander	Ψ.	3.1 *	0.8 - 5.4	100	15.3	13.9 - 16.7	73,700	
Nativity and Language								
Born outside the U.S.	•	37.0	31.9 - 42.1	1,100	37.9	36.2 - 39.6	185,700	
Speaks non-English language at home	•	34.6	29.6 - 39.6	1,000	30.1	28.5 - 31.7	144,800	
Lives in a high poverty neighborhood	•	39.8	34.5 - 45.1	1,100	38.9	37.3 - 40.5	182,900	
Income as a percent of the Federal Poverty Guideline				•			•	
0-100%	•	41.3	35.9 - 46.7	1,100	41.0	39.5 - 42.4	188,800	
101-200%	•	20.9	16.7 - 25.1	600	20.8	19.4 - 22.2	96,000	
> 200%	•	37.8	32.5 - 43.0	1,000	38.2	36.9 - 39.6		
Estimate not shown because the relative standard error (RSE) is greater than 509	/ or four			_,		3 2 2 2 2 3 3 1 0	,	

⁻⁻ Estimate not shown because the relative standard error (RSE) is greater than 50% or fewer than 5 women reported.

Notes: MIHA is an annual population-based survey of California resident women with a live birth. Data from MIHA 2013-2014 were combined, resulting in a statewide sample size of 13,963. The sample size of Santa Cruz County was 400. Percent (%), 95% confidence interval (95% CI) and estimated number of women in the population with the health indicator/characteristic (i.e., numerator of the percent rounded to the nearest hundred) are weighted to represent all women with a live birth who resided in California and the county in 2013-2014. Population estimate is a two-year average. Symbols indicate whether the health indicator in the county was statistically different from the rest of the state (p-value < 0.05, chi-square test). See the Technical Notes for information on weighting, comparability to prior years and technical definitions. Visit the MIHA website at www.cdph.ca.gov/MIHA.

Table Citation: Maternal and Infant Health Assessment (MIHA) Survey Data Snapshots, 2013-2014. California Department of Public Health. 2016.

 $^{^{*}}$ Estimate should be interpreted with caution due to low statistical reliability (RSE is between 30% and 50%).

[†] Measure/definition changed for this indicator and is not comparable to prior years (see Technical Notes).