











California – Chlamydia, Gonorrhea, and P&S Syphilis Rates by County, 2010



Note: Rates are per 100,000 population.
Source: California Department of Public Health, STD Control Branch

California – Chlamydia, Gonorrhea, and P&S Syphilis Rates by Age Group (2010), Race/Ethnicity (2010), and Year



■ Female ■ Male

— Female - - - Male



* Race data may be missing for a substantial number of cases. See the 'Data Limitations' page for further information.
 Note: Rates are per 100,000 population.
 Source: California Department of Public Health, STD Control Branch

California State – Chlamydia, Gonorrhea, and Early Syphilis Cases and Rates Tables for 2010

| Gender & Age Group | Chlamydia | | | Gonorrhea | | | P&S Syphilis | | | Early Latent Syphilis | | | Population | |
|---------------------|-----------|----------------------|---------|-----------|----------------------|-------|--------------|----------------------|------|-----------------------|----------------------|------|------------|----------------------|
| | Cases | Percent ¹ | Rate | Cases | Percent ¹ | Rate | Cases | Percent ¹ | Rate | Cases | Percent ¹ | Rate | Number | Percent ¹ |
| STATE TOTAL | 155,300 | 100.0% | 400.0 | 26,840 | 100.0% | 69.1 | 2,059 | 100.0% | 5.3 | 1,805 | 100.0% | 4.6 | 39,135,676 | 100.0% |
| Female Total | 106,339 | 68.5% | 542.3 | 10,757 | 40.1% | 54.9 | 75 | 3.6% | 0.4 | 126 | 7.0% | 0.6 | 19,608,793 | 50.1% |
| Ages | | | | | | | | | | | | | | |
| 0 – 9 | 52 | 0.0% | 1.9 | 12 | 0.1% | 0.4 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 2,701,285 | 13.8% |
| 10 – 14 | 908 | 0.9% | 69.1 | 148 | 1.4% | 11.3 | 1 | 1.3% | 0.1 | 0 | 0.0% | 0.0 | 1,313,498 | 6.7% |
| 15 – 19 | 33,440 | 31.5% | 2,247.0 | 3,349 | 31.2% | 225.0 | 10 | 13.3% | 0.7 | 21 | 16.7% | 1.4 | 1,488,238 | 7.6% |
| 20 – 24 | 41,406 | 39.1% | 2,950.5 | 3,637 | 33.9% | 259.2 | 12 | 16.0% | 0.9 | 28 | 22.2% | 2.0 | 1,403,352 | 7.2% |
| 25 – 29 | 16,646 | 15.7% | 1,296.8 | 1,732 | 16.1% | 134.9 | 14 | 18.7% | 1.1 | 21 | 16.7% | 1.6 | 1,283,661 | 6.5% |
| 30 – 34 | 6,961 | 6.6% | 574.2 | 887 | 8.3% | 73.2 | 4 | 5.3% | 0.3 | 14 | 11.1% | 1.2 | 1,212,291 | 6.2% |
| 35 – 44 | 4,998 | 4.7% | 184.8 | 691 | 6.4% | 25.6 | 18 | 24.0% | 0.7 | 22 | 17.5% | 0.8 | 2,703,970 | 13.8% |
| 45 + | 1,618 | 1.5% | 21.6 | 274 | 2.6% | 3.7 | 16 | 21.3% | 0.2 | 20 | 15.9% | 0.3 | 7,502,498 | 38.3% |
| Not Specified | 310 | 0.3% | - | 27 | 0.3% | - | 0 | 0.0% | - | 0 | 0.0% | - | | |
| Male Total | 48,358 | 31.1% | 247.6 | 15,961 | 59.5% | 81.7 | 1,983 | 96.3% | 10.2 | 1,676 | 92.9% | 8.6 | 19,526,883 | 49.9% |
| Ages | | | | | | | | | | | | | | |
| 0 – 9 | 26 | 0.1% | 0.9 | 4 | 0.0% | 0.1 | 0 | 0.0% | 0.0 | 1 | 0.1% | 0.0 | 2,812,744 | 14.4% |
| 10 – 14 | 174 | 0.4% | 12.7 | 32 | 0.2% | 2.3 | 2 | 0.1% | 0.1 | 0 | 0.0% | 0.0 | 1,371,611 | 7.0% |
| 15 – 19 | 8,762 | 18.2% | 559.4 | 1,894 | 11.9% | 120.9 | 65 | 3.3% | 4.2 | 39 | 2.3% | 2.5 | 1,566,183 | 8.0% |
| 20 – 24 | 16,432 | 34.1% | 1,086.6 | 4,011 | 25.2% | 265.2 | 270 | 13.6% | 17.9 | 171 | 10.2% | 11.3 | 1,512,182 | 7.7% |
| 25 – 29 | 10,072 | 20.9% | 726.5 | 3,237 | 20.3% | 233.5 | 311 | 15.7% | 22.4 | 222 | 13.3% | 16.0 | 1,386,408 | 7.1% |
| 30 – 34 | 5,045 | 10.5% | 395.9 | 2,080 | 13.1% | 163.2 | 297 | 15.0% | 23.3 | 212 | 12.7% | 16.6 | 1,274,379 | 6.5% |
| 35 – 44 | 5,007 | 10.4% | 179.7 | 2,813 | 17.7% | 101.0 | 559 | 28.2% | 20.1 | 570 | 34.0% | 20.5 | 2,786,242 | 14.3% |
| 45 + | 2,683 | 5.6% | 39.4 | 1,841 | 11.6% | 27.0 | 477 | 24.1% | 7.0 | 460 | 27.5% | 6.7 | 6,817,134 | 34.9% |
| Not Specified | 157 | 0.3% | - | 49 | 0.3% | - | 2 | 0.1% | - | 1 | 0.1% | - | | |

| Gender & Race/Ethnicity | Chlamydia | | | Gonorrhea | | | P&S Syphilis | | | Early Latent Syphilis | | | Population | |
|-------------------------------|-----------|----------------------|---------|-----------|----------------------|-------|--------------|----------------------|------|-----------------------|----------------------|------|------------|----------------------|
| | Cases | Percent ¹ | Rate | Cases | Percent ¹ | Rate | Cases | Percent ¹ | Rate | Cases | Percent ¹ | Rate | Number | Percent ¹ |
| STATE TOTAL | 155,300 | 100.0% | 400.0 | 26,840 | 100.0% | 69.1 | 2,059 | 100.0% | 5.3 | 1,805 | 100.0% | 4.6 | 39,135,676 | 100.0% |
| Female Total | 106,339 | 68.5% | 542.3 | 10,757 | 40.1% | 54.9 | 75 | 3.6% | 0.4 | 126 | 7.0% | 0.6 | 19,608,793 | 50.1% |
| American Indian/Alaska Native | 305 | 0.5% | 249.2 | 27 | 0.4% | 22.1 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 122,373 | 0.6% |
| Asian/Pacific Islander | 4,108 | 6.1% | 163.5 | 274 | 3.8% | 10.9 | 1 | 1.4% | 0.0 | 5 | 4.0% | 0.2 | 2,512,224 | 12.8% |
| Black/African-American | 15,902 | 23.7% | 1,359.3 | 3,511 | 48.8% | 300.1 | 32 | 45.7% | 2.7 | 44 | 35.5% | 3.8 | 1,169,844 | 6.0% |
| Latina/Hispanic | 33,189 | 49.5% | 466.8 | 1,968 | 27.3% | 27.7 | 14 | 20.0% | 0.2 | 56 | 45.2% | 0.8 | 7,110,156 | 36.3% |
| White | 13,591 | 20.3% | 164.2 | 1,421 | 19.7% | 17.2 | 23 | 32.9% | 0.3 | 19 | 15.3% | 0.2 | 8,278,622 | 42.2% |
| Other/Multi/Not Specified | 39,244 | 36.9% | - | 3,556 | 33.1% | - | 5 | 6.7% | - | 2 | 1.6% | - | 415,574 | 2.1% |
| Male Total | 48,358 | 31.1% | 247.6 | 15,961 | 59.5% | 81.7 | 1,983 | 96.3% | 10.2 | 1,676 | 92.9% | 8.6 | 19,526,883 | 49.9% |
| American Indian/Alaska Native | 112 | 0.4% | 94.6 | 34 | 0.3% | 28.7 | 7 | 0.4% | 5.9 | 3 | 0.2% | 2.5 | 118,348 | 0.6% |
| Asian/Pacific Islander | 1,570 | 5.0% | 67.6 | 506 | 4.4% | 21.8 | 92 | 4.8% | 4.0 | 67 | 4.2% | 2.9 | 2,321,659 | 11.9% |
| Black/African-American | 8,651 | 27.7% | 774.2 | 3,877 | 33.7% | 347.0 | 276 | 14.4% | 24.7 | 230 | 14.3% | 20.6 | 1,117,346 | 5.7% |
| Latina/Hispanic | 13,372 | 42.9% | 180.6 | 3,422 | 29.7% | 46.2 | 663 | 34.6% | 9.0 | 695 | 43.4% | 9.4 | 7,402,661 | 37.9% |
| White | 7,501 | 24.0% | 91.9 | 3,680 | 31.9% | 45.1 | 878 | 45.8% | 10.8 | 608 | 37.9% | 7.5 | 8,160,162 | 41.8% |
| Other/Multi/Not Specified | 17,152 | 35.5% | - | 4,442 | 27.8% | - | 67 | 3.4% | - | 73 | 4.4% | - | 406,707 | 2.1% |

¹ Gender specific age groups and race/ethnicity percent calculations exclude "Not Specified" from the denominator.

² Overall state rates were calculated using the July 1 county population estimates from the Department of Finance.
(For California, the July 1, 2010 estimate is 38,826,898).

Note: Rates are per 100,000 population.




Source: California Department of Public Health, STD Control Branch



California – Chlamydia, Gonorrhea, and P&S Syphilis

Ranking of County Rates for 2010

(with 95% Confidence Intervals*)



* Confidence intervals were calculated using Poisson exact method; not shown for counties with zero cases.

Note: Rates are per 100,000 population.

Source: California Department of Public Health, STD Control Branch

