

2011 MIHA County Charts


Geographic Comparisons of Select Indicators from the Maternal and Infant Health Assessment (MIHA) Survey

Maternal, Child and Adolescent Health Program
Center for Family Health
California Department of Public Health

September 2013



MIHA counties sampled in 2011



Top 20 Birthing Counties


% of resident women with a live birth in 2011

- Los Angeles 26.0%
- San Diego 8.7%
- Orange 7.6%
- Riverside 6.1%
- San Bernardino..... 6.1%
- Santa Clara 4.7%
- Sacramento 4.0%
- Alameda 3.8%
- Fresno 3.2%
- Kern 2.9%
- Contra Costa 2.4%
- San Joaquin 2.1%
- Ventura 2.1%
- San Mateo 1.8%
- San Francisco 1.7%
- Tulare 1.6%
- Stanislaus 1.5%
- Monterey 1.4%
- Santa Barbara 1.2%
- Sonoma 1.0%

Data Source: 2011 Birth Statistical Master File


Prepared by: Maternal, Child and Adolescent Health Program,
Center for Family Health, California Department of Public Health

Daily folic acid use in the month before pregnancy, MIHA 2011




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Obesity before pregnancy (BMI 30+), MIHA 2011




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Excessive weight gain during pregnancy, MIHA 2011




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Food insecurity during pregnancy, MIHA 2011




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Received CalFresh (food stamps) during pregnancy, MIHA 2011




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Physical or psychological intimate partner violence (IPV) during pregnancy, MIHA 2011




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Prenatal depressive symptoms, MIHA 2011



The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.



Postpartum depressive symptoms, MIHA 2011



The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.




Woman or partner lost job during pregnancy, MIHA 2011




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Had to move due to problems paying rent or mortgage during pregnancy, MIHA 2011




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars. *Data are not presented because fewer than 10 women reported.

Any smoking during the 1st or 3rd trimester, MIHA 2011




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars. *Data are not presented because fewer than 10 women reported.

Any binge drinking during the 3 months before pregnancy, MIHA 2011




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Mistimed or unwanted pregnancy, MIHA 2011




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Postpartum birth control use, MIHA 2011




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Placed infant on back to sleep, MIHA 2011




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Always or often bedshared, MIHA 2011



The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.



Exclusive breastfeeding at 3 months after delivery, MIHA 2011



The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.




Had a usual source of pre-pregnancy health care, MIHA 2011




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Initiated prenatal care in the 1st trimester, MIHA 2011




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Mom or infant needed but could not afford care postpartum, MIHA 2011




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Uninsured before pregnancy, MIHA 2011




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Covered by Medi-Cal during pregnancy, MIHA 2011




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Uninsured postpartum, MIHA 2011




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Infant covered by Medi-Cal, MIHA 2011




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Speaks non-English language at home, MIHA 2011




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Income within 0-100% of the Federal Poverty Guideline, MIHA 2011



The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Income within 0-200% of the Federal Poverty Guideline, MIHA 2011



The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,853 in 2011. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2011 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.