


2010 MIHA Charts by Prenatal Health Insurance

Indicators from the Maternal and Infant Health Assessment (MIHA) Survey

Maternal, Child and Adolescent Health Program
Center for Family Health
California Department of Public Health
September 2013




Daily folic acid use in the month before pregnancy, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Obese before pregnancy (BMI 30+), MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Excessive weight gain during pregnancy, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Food insecurity during pregnancy, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Participated in WIC during pregnancy, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Received CalFresh (food stamps) during pregnancy, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Physical or psychological intimate partner violence (IPV) during pregnancy, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Prenatal depressive symptoms, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Postpartum depressive symptoms, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Had a lot of unpaid bills during pregnancy, MIHA 2010



The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.


Woman or partner lost job during pregnancy, MIHA 2010



The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.




Any smoking during the 1st or 3rd trimester, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Any binge drinking during the 3 months before pregnancy, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Placed infant on back to sleep, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Exclusive breastfeeding at 3 months after delivery, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Had a usual source of pre-pregnancy health care, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Initiated prenatal care in the 1st trimester, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Mom or infant needed but could not afford care postpartum, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Uninsured before pregnancy, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Uninsured postpartum, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Infant covered by Medi-Cal, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Speaks non-English language at home, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Income within 0-100% of the Federal Poverty Guideline, MIHA 2010



The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Income within 0-200% of the Federal Poverty Guideline, MIHA 2010



The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.