

# 2010 MIHA Charts by Race/Ethnicity

Indicators from the Maternal and Infant Health Assessment (MIHA) Survey

Maternal, Child and Adolescent Health Program


Center for Family Health

California Department of Public Health

September 2013




# Daily folic acid use in the month before pregnancy, MIHA 2010



The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.



# Obesity before pregnancy (BMI 30+), MIHA 2010



The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.




# Excessive weight gain during pregnancy, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.

# Food insecurity during pregnancy, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.

# Participated in WIC during pregnancy, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.

# Received CalFresh (food stamps) during pregnancy, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.

# Physical or psychological intimate partner violence (IPV) during pregnancy, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.

# Prenatal depressive symptoms, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.

# Postpartum depressive symptoms, MIHA 2010



The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.



# Had a lot of unpaid bills during pregnancy, MIHA 2010



The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.





# Woman or partner lost job during pregnancy, MIHA 2010



The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.




# Any smoking during the 1<sup>st</sup> or 3<sup>rd</sup> trimester, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.

# Any binge drinking during the 3 months before pregnancy, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.

## Placed infant on back to sleep, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.

# Exclusive breastfeeding at 3 months after delivery, MIHA 2010



The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.



# Had a usual source of pre-pregnancy health care, MIHA 2010



The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.





# Initiated prenatal care in the 1<sup>st</sup> trimester, MIHA 2010



The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.




# Mom or infant needed but could not afford care postpartum, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.

# Uninsured before pregnancy, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.

# Covered by Medi-Cal during pregnancy, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.

# Uninsured postpartum, MIHA 2010



The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.



# Infant covered by Medi-Cal, MIHA 2010



The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.




# Speaks non-English language at home, MIHA 2010




The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.

# Income within 0-100% of the Federal Poverty Guideline, MIHA 2010



The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.

# Income within 0-200% of the Federal Poverty Guideline, MIHA 2010



The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of n=6,817 in 2010. Percentages and 95% confidence intervals are weighted to represent all women with a live birth in 2010 in California and in the counties shown. Confidence intervals are shown as thin black lines extending above and below the top of the bars.