



2012 MIHA Regional Charts

Geographic Comparisons of Selected Indicators from the
Maternal and Infant Health Assessment (MIHA) Survey



Maternal, Child and Adolescent Health Program
Center for Family Health
California Department of Public Health

December 2014




MIHA is a collaborative effort of the Maternal, Child and Adolescent Health and WIC Programs of the Center for Family Health, California Department of Public Health and the Center on Social Disparities in Health at the University of California, San Francisco. Visit the MIHA website at www.cdph.ca.gov/MIHA.

MIHA Regions of California






Daily folic acid use in the month before pregnancy, MIHA 2012



Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.




Obesity before pregnancy (BMI 30+), MIHA 2012




Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Excessive weight gain during pregnancy, MIHA 2012






Food insecurity during pregnancy, MIHA 2012



Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.




Participated in WIC during pregnancy, MIHA 2012



Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.




Received CalFresh (food stamps) during pregnancy, MIHA 2012



Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.




Physical or psychological intimate partner violence (IPV) during pregnancy, MIHA 2012



Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.




Prenatal depressive symptoms, MIHA 2012



Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.




Postpartum depressive symptoms, MIHA 2012



Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.



Woman or partner lost job during pregnancy, MIHA 2012




California
17.3
↓

Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.




Had to move due to problems paying rent or mortgage during pregnancy, MIHA 2012



Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.




Any smoking during the 1st or 3rd trimester, MIHA 2012



Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.




Any binge drinking during the 3 months before pregnancy, MIHA 2012



Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.




Mistimed or unwanted pregnancy, MIHA 2012



Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.




Postpartum birth control use, MIHA 2012



Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.




Placed infant on back to sleep, MIHA 2012



Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.




Exclusive breastfeeding at 3 months after delivery, MIHA 2012



Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.



Had a usual source of pre-pregnancy health care, MIHA 2012




**California
72.0**



Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.



Initiated prenatal care in the 1st trimester,[†] MIHA 2012




Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

[†] In 2012, the measure changed for this indicator and is not comparable to prior years (see Technical Document).



Had a dental visit during pregnancy, MIHA 2012




California
42.1
↓

Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.




Mom or infant needed but could not afford care postpartum, MIHA 2012



Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.




Uninsured before pregnancy, MIHA 2012



Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.




Covered by Medi-Cal during pregnancy, MIHA 2012




Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.

Uninsured postpartum, MIHA 2012




Infant covered by Medi-Cal, MIHA 2012






Speaks non-English language at home, MIHA 2012



Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.




Income within 0-100% of the Federal Poverty Guideline, MIHA 2012



Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.



Income within 0-200% of the Federal Poverty Guideline, MIHA 2012



Prepared by the Epidemiology, Assessment, and Program Development Branch, MCAH Division. The Maternal and Infant Health Assessment (MIHA) Survey is an annual population-based survey of women with a live birth, with a sample size of 6,810 in 2012. Percentages and 95% confidence intervals (95% CI) are weighted to represent all women with a live birth in 2012 in California and in the regions shown. Confidence intervals are shown as thin black lines extending above and below the top of the blue bars.