




MIHA County Comparison 2012

Maternal and Infant Health Assessment (MIHA) Survey



Physical or psychological intimate partner violence (IPV) during pregnancy

County	Percent	95% CI	Population Estimate of Indicator
California	8.8	7.5 - 10.2	43,400
Alameda	♦ 7.9	3.9 - 11.8	1,500
Contra Costa	♦ 7.3	2.5 - 12.1	900
Fresno	♦ 12.0	7.5 - 16.6	1,900
Kern	♦ 9.5	5.4 - 13.7	1,400
Los Angeles	♦ 9.4	5.0 - 13.9	12,200
Monterey	♦ 12.9	7.0 - 18.9	800
Orange	♦ 9.4	5.5 - 13.3	3,500
Riverside	♦ 7.9	4.7 - 11.1	2,300
Sacramento	♦ 9.0	5.5 - 12.5	1,700
San Bernardino	♦ 8.6	4.9 - 12.3	2,600
San Diego	♦ 7.4	4.0 - 10.9	3,200
San Francisco	♦ 6.1	2.7 - 9.5	500
San Joaquin	♦ 8.9	5.0 - 12.8	900
San Mateo	♦ 6.5	2.4 - 10.6	600
Santa Barbara	♦ 12.1	7.3 - 16.8	700
Santa Clara	♦ 9.6	5.3 - 13.9	2,300
Sonoma	✓ 4.1	1.5 - 6.7	200
Stanislaus	✗ 16.2	10.6 - 21.8	1,200
Tulare	♦ 11.5	6.3 - 16.6	900
Ventura	✓ 2.4	0.5 - 4.2	200

- ✓ Statistically better than the rest of California (p<0.05, chi-square test)
- ✗ Statistically worse than the rest of California (p<0.05, chi-square test)
- ♦ No statistical difference between county and the rest of California

Notes: MIHA is an annual population-based survey of California resident women with a live birth in 2012, with a sample size of 6,810. Percent (%), 95% confidence interval (95% CI), and estimated number of women in the population with the health indicator/characteristic (N), i.e., numerator of the percent rounded to the nearest hundred, are weighted to represent all women with a live birth in California and the county in 2012. See the Technical Document for information on weighting, comparability to prior years and technical definitions.