

CALIFORNIA DIABETES AND PREGNANCY PROGRAM - CALENDAR YEAR 2008

DATA REPORT for deliveries between January 1, 2008 and by December 31, 2008

Characteristics by Maternal Type of Diabetes and Intervention at Last Sweet Success Visit (number and percent) ¹

| Generated 03/17/2011 14:28:02 | Gestational Diabetes Mellitus (GDM) | | | | | | | | | | | | | | | | | |
|--|--|--------|---------|-------|----------------|-------|-----------|-------|-------|-------|---------------------|-------|----------------------------|-------|---------|-------|-------|---|
| State of California | Total | | Insulin | | Insulin & Oral | | Oral Meds | | Diet | | History of GDM Diet | | Impaired Glucose Tol. Diet | | Unknown | | None | |
| | num # | % | num # | % | num # | % | num # | % | num # | % | num # | % | num # | % | num # | % | num # | % |
| Sweet Success Clients | 12710 | 100.0% | 2690 | 21.2% | 70 | 0.6% | 1869 | 14.7% | 8081 | 63.6% | 78 | 0.0% | 669 | 0.0% | 586 | 0.0% | | |
| Live Births | 10583 | 83.3% | 2272 | 84.5% | 61 | 87.1% | 1514 | 81.0% | 6736 | 83.4% | 58 | 74.4% | 620 | 92.7% | 503 | 85.8% | | |
| Fetal Deaths | 14 | 0.1% | | | | | | | 11 | 0.1% | | | | 11 | 1.9% | | | |
| Spont. Abortions | 110 | 0.9% | 27 | 1.0% | | | | | 81 | 1.0% | 6 | 7.7% | 10 | 1.5% | | | | |
| Other ² | 44 | 0.3% | 8 | 0.3% | | | | | 33 | 0.4% | | | | | | | | |
| Unknown Outcomes | 1959 | 15.4% | 382 | 14.2% | 9 | 12.9% | 348 | 18.6% | 1220 | 15.1% | 13 | 16.7% | 34 | 5.1% | 68 | 11.6% | | |
| Maternal Characteristics ³ | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | |
| White | 2098 | 16.7% | 520 | 19.5% | 14 | 20.0% | 300 | 16.1% | 1264 | 15.8% | 20 | 27.8% | 145 | 22.0% | 68 | 11.7% | | |
| Hispanic | 6976 | 55.4% | 1423 | 53.4% | 40 | 57.1% | 1143 | 61.2% | 4370 | 54.6% | 33 | 45.8% | 345 | 52.4% | 207 | 35.6% | | |
| Black | 281 | 2.2% | 71 | 2.7% | | | 40 | 2.1% | 167 | 2.1% | | | 10 | 1.5% | 17 | 2.9% | | |
| Asian/Pacific Islander | 1914 | 15.2% | 413 | 15.5% | 6 | 8.6% | 207 | 11.1% | 1288 | 16.1% | 9 | 12.5% | 99 | 15.0% | 35 | 6.0% | | |
| Native American | 31 | 0.2% | 7 | 0.3% | | | 7 | 0.4% | 17 | 0.2% | | | | | | | | |
| Multiple Race | | | | | | | | | | | | | | | | | | |
| Other/Unknown | 1300 | 10.3% | 229 | 8.6% | 7 | 10.0% | 170 | 9.1% | 894 | 11.2% | 8 | 11.1% | 58 | 8.8% | 254 | 43.6% | | |
| Education | | | | | | | | | | | | | | | | | | |
| 0 - 3rd grade | 59 | 0.5% | 9 | 0.3% | | | 14 | 0.7% | 36 | 0.5% | | | | | | | | |
| 4th - 8th | 1239 | 9.8% | 230 | 8.6% | 7 | 10.0% | 250 | 13.4% | 752 | 9.4% | | | 68 | 10.3% | 39 | 6.7% | | |
| 9th - 12th | 4347 | 34.5% | 845 | 31.7% | 28 | 40.0% | 754 | 40.4% | 2720 | 34.0% | 23 | 31.9% | 238 | 36.1% | 127 | 21.8% | | |
| any post H.S. | 4670 | 37.1% | 965 | 36.2% | 28 | 40.0% | 605 | 32.4% | 3072 | 38.4% | 36 | 50.0% | 250 | 37.9% | 110 | 18.9% | | |
| Unknown | 2285 | 18.1% | 614 | 23.1% | 7 | 10.0% | 244 | 13.1% | 1420 | 17.8% | 9 | 12.5% | 101 | 15.3% | 304 | 52.2% | | |
| Payment Source | | | | | | | | | | | | | | | | | | |
| Private Ins. | | | | | | | | | | | | | | | | | | |
| HMO | 2613 | 20.7% | 584 | 21.9% | 16 | 22.9% | 343 | 18.4% | 1670 | 20.9% | 13 | 18.1% | 121 | 18.4% | 66 | 11.3% | | |
| PPO | 2563 | 20.3% | 543 | 20.4% | 9 | 12.9% | 293 | 15.7% | 1718 | 21.5% | 18 | 25.0% | 154 | 23.4% | 64 | 11.0% | | |
| MediCal | 6497 | 51.6% | 1319 | 49.5% | 43 | 61.4% | 1107 | 59.3% | 4028 | 50.4% | 35 | 48.6% | 349 | 53.0% | 185 | 31.8% | | |
| Self Pay | 144 | 1.1% | 30 | 1.1% | | | 30 | 1.6% | 84 | 1.1% | | | | | | | | |
| Multiple | 81 | 0.6% | 37 | 1.4% | | | | | 40 | 0.5% | | | | | | | | |
| None/Other | 702 | 5.6% | 150 | 5.6% | | | 90 | 4.8% | 460 | 5.8% | 5 | 6.9% | 32 | 4.9% | 263 | 45.2% | | |

1. Cell values less than 5 are suppressed.

2. "Other" includes induced terminations.

3. Tables by specific maternal characteristics (e.g. race/ethnicity, education, payment source) exclude mothers with spontaneous abortions.

CALIFORNIA DIABETES AND PREGNANCY PROGRAM - CALENDAR YEAR 2008

DATA REPORT for deliveries between January 1, 2008 and by December 31, 2008

Characteristics by Maternal Type of Diabetes and Intervention at Last Sweet Success Visit (number and percent)¹

| Generated 03/17/2011 14:28:02 | Gestational Diabetes Mellitus (GDM) | | | | | | | | | | | | | | | | | |
|--|--|--------|---------|--------|----------------|--------|-----------|--------|-------|--------|---------------------|--------|----------------------------|--------|---------|--------|-------|---|
| State of California | Total | | Insulin | | Insulin & Oral | | Oral Meds | | Diet | | History of GDM Diet | | Impaired Glucose Tol. Diet | | Unknown | | None | |
| | num # | % | num # | % | num # | % | num # | % | num # | % | num # | % | num # | % | num # | % | num # | % |
| Maternal Characteristics ³ | | | | | | | | | | | | | | | | | | |
| Maternal Age | | | | | | | | | | | | | | | | | | |
| < 18 years | 47 | 0.4% | | | | | | | 42 | 0.5% | | | 5 | 0.8% | 5 | 0.9% | | |
| 18 - 24 years | 1489 | 11.8% | 259 | 9.7% | 5 | 7.1% | 224 | 12.0% | 1001 | 12.5% | 8 | 11.1% | 107 | 16.2% | 42 | 7.2% | | |
| 25 - 34 years | 6440 | 51.1% | 1393 | 52.3% | 33 | 47.1% | 964 | 51.6% | 4050 | 50.6% | 30 | 41.7% | 353 | 53.6% | 192 | 33.0% | | |
| 35 - 39 years | 2828 | 22.4% | 638 | 24.0% | 23 | 32.9% | 426 | 22.8% | 1741 | 21.8% | 20 | 27.8% | 126 | 19.1% | 65 | 11.2% | | |
| 40 - 45 years | 932 | 7.4% | 224 | 8.4% | | | 141 | 7.6% | 565 | 7.1% | 7 | 9.7% | 41 | 6.2% | 27 | 4.6% | | |
| > 45 years | 33 | 0.3% | 8 | 0.3% | | | | | 21 | 0.3% | | | | | | | | |
| Unknown | 831 | 6.6% | 139 | 5.2% | 7 | 10.0% | 105 | 5.6% | 580 | 7.3% | 7 | 9.7% | 23 | 3.5% | 251 | 43.1% | | |
| Gravida | | | | | | | | | | | | | | | | | | |
| 1 | 2747 | 21.8% | 486 | 18.3% | 13 | 18.6% | 367 | 19.7% | 1881 | 23.5% | | | 174 | 26.4% | 73 | 12.5% | | |
| 2 to 4 | 7620 | 60.5% | 1602 | 60.2% | 46 | 65.7% | 1154 | 61.8% | 4818 | 60.2% | 53 | 73.6% | 389 | 59.0% | 190 | 32.6% | | |
| > 4 | 1970 | 15.6% | 501 | 18.8% | 10 | 14.3% | 321 | 17.2% | 1138 | 14.2% | 15 | 20.8% | 93 | 14.1% | 40 | 6.9% | | |
| Unknown | 263 | 2.1% | 74 | 2.8% | | | 25 | 1.3% | 163 | 2.0% | | | | | 279 | 47.9% | | |
| Para | | | | | | | | | | | | | | | | | | |
| 1 | 3556 | 28.2% | 687 | 25.8% | 17 | 24.3% | 467 | 25.0% | 2385 | 29.8% | 6 | 8.3% | 232 | 35.2% | 91 | 15.6% | | |
| 2 to 4 | 6891 | 54.7% | 1583 | 59.4% | 45 | 64.3% | 1039 | 55.7% | 4224 | 52.8% | 59 | 81.9% | 359 | 54.5% | 164 | 28.2% | | |
| > 4 | 792 | 6.3% | 203 | 7.6% | | | 123 | 6.6% | 462 | 5.8% | | | 31 | 4.7% | 16 | 2.7% | | |
| Unknown | 1361 | 10.8% | 190 | 7.1% | | | 238 | 12.7% | 929 | 11.6% | | | 37 | 5.6% | 311 | 53.4% | | |
| Method of Delivery | | | | | | | | | | | | | | | | | | |
| Vaginal (including VBAC) | 5973 | 47.4% | 1239 | 46.5% | 31 | 44.3% | 825 | 44.2% | 3878 | 48.5% | 41 | 56.9% | 363 | 55.1% | 110 | 18.9% | | |
| Cesarean - Primary | 2502 | 19.9% | 593 | 22.3% | 14 | 20.0% | 413 | 22.1% | 1482 | 18.5% | 8 | 11.1% | 126 | 19.1% | 40 | 6.9% | | |
| Cesarean - Secondary | 2048 | 16.3% | 535 | 20.1% | 19 | 27.1% | 335 | 17.9% | 1159 | 14.5% | 13 | 18.1% | 101 | 15.3% | 31 | 5.3% | | |
| Unknown | 2077 | 16.5% | 296 | 11.1% | 6 | 8.6% | 294 | 15.7% | 1481 | 18.5% | 10 | 13.9% | 69 | 10.5% | 401 | 68.9% | | |
| Preconception Care ⁴ | | | | | | | | | | | | | | | | | | |
| Yes | | | | | | | | | | | | | | | | | | |
| No | | | | | | | | | | | | | | | | | | |
| Not Applicable | 12600 | 100.0% | 2663 | 100.0% | 70 | 100.0% | 1867 | 100.0% | 8000 | 100.0% | 72 | 100.0% | 659 | 100.0% | 582 | 100.0% | | |
| Unknown | | | | | | | | | | | | | | | | | | |
| Hemoglobin A1c ⁵ | | | | | | | | | | | | | | | | | | |
| Pre-pregnancy known | 13 | | 6 | | | | | | 5 | | | | 10 | | | | | |
| Pre-pregnancy > 7.0 | | | | | | | | | | | | | | | | | | |
| Prenatal known | 3473 | | 1192 | | 28 | | 463 | | 1790 | | 25 | | 197 | | 62 | | | |
| Prenatal > 7.0 | 144 | 4.1% | 112 | 9.4% | | | 10 | 2.2% | 18 | 1.0% | | | 8 | 4.1% | 10 | 16.1% | | |

1. Cell values less than 5 are suppressed.
3. Tables by specific maternal characteristics (e.g. race/ethnicity, education, payment source) exclude mothers with spontaneous abortions.
4. Preconception care was reported only among women with pre-existing diabetes (Type 1 or 2). All other categories are reported as Not Applicable. Totals exclude spontaneous abortions.
5. Excludes clients with a spontaneous abortion and those whose HbA1c test results were not reported.

CALIFORNIA DIABETES AND PREGNANCY PROGRAM - CALENDAR YEAR 2008

DATA REPORT for deliveries between January 1, 2008 and by December 31, 2008

Characteristics by Maternal Type of Diabetes and Intervention at Last Sweet Success Visit (number and percent) ¹

| | | | | | | | | | | | | | | | | | | |
|-------------------------------|--|---|---------|---|----------------|---|-----------|---|-------|---|---------------------|---|----------------------------|---|---------|---|-------|---|
| Generated 03/17/2011 14:28:02 | Gestational Diabetes Mellitus (GDM) | | | | | | | | | | | | | | | | | |
| State of California | Total | | Insulin | | Insulin & Oral | | Oral Meds | | Diet | | History of GDM Diet | | Impaired Glucose Tol. Diet | | Unknown | | None | |
| | num # | % | num # | % | num # | % | num # | % | num # | % | num # | % | num # | % | num # | % | num # | % |

Maternal Characteristics ³

| Weeks Gestation at Entry Into Sweet Success Care | | | | | | | | | | | | | | | | | | |
|---|------|-------|------|--------|------|-------|------|--------|------|-------|-----|-------|-----|-------|-----|-------|-----|--|
| < 14 weeks | 946 | 7.5% | 388 | 14.6% | 8 | 11.4% | 161 | 8.6% | 389 | 4.9% | 27 | 37.5% | 61 | 9.3% | 23 | 4.0% | | |
| 14 to 27 weeks | 2900 | 23.0% | 868 | 32.6% | 27 | 38.6% | 515 | 27.6% | 1490 | 18.6% | 23 | 31.9% | 149 | 22.6% | 36 | 6.2% | | |
| 28 to 32 weeks | 5763 | 45.7% | 1053 | 39.5% | 26 | 37.1% | 883 | 47.3% | 3801 | 47.5% | 17 | 23.6% | 309 | 46.9% | 72 | 12.4% | | |
| 33 to 37 weeks | 2653 | 21.1% | 315 | 11.8% | 8 | 11.4% | 283 | 15.2% | 2047 | 25.6% | | | 121 | 18.4% | 39 | 6.7% | | |
| > 38 weeks | 210 | 1.7% | 18 | 0.7% | | | 12 | 0.6% | 180 | 2.3% | | | 15 | 2.3% | 6 | 1.0% | | |
| Unknown | 128 | 1.0% | 21 | 0.8% | | | 13 | 0.7% | 93 | 1.2% | | | | | 406 | 69.8% | | |
| Post-partum Testing Done | 5159 | 83.8% | 1161 | 22.5% | 37 | 0.7% | 888 | 17.2% | 3073 | 59.6% | 26 | 0.4% | 355 | 5.8% | 62 | 1.0% | | |
| SS Interventions (total) | | | | | | | | | | | | | | | | | | |
| 1 to 4 | 4106 | 32.6% | 407 | 15.3% | 5 | 7.1% | 369 | 19.8% | 3325 | 41.6% | 24 | 33.3% | 211 | 32.0% | 62 | 10.7% | | |
| 5 to 10 | 5221 | 41.4% | 1060 | 39.8% | 25 | 35.7% | 854 | 45.7% | 3282 | 41.0% | 26 | 36.1% | 268 | 40.7% | 61 | 10.5% | | |
| 11 to 15 | 1443 | 11.5% | 547 | 20.5% | 17 | 24.3% | 306 | 16.4% | 573 | 7.2% | 12 | 16.7% | 83 | 12.6% | 18 | 3.1% | | |
| > 15 | 1209 | 9.6% | 570 | 21.4% | 18 | 25.7% | 261 | 14.0% | 360 | 4.5% | 7 | 9.7% | 80 | 12.1% | 12 | 2.1% | | |
| Unknown | 621 | 4.9% | 79 | 3.0% | 5 | 7.1% | 77 | 4.1% | 460 | 5.8% | | | 17 | 2.6% | 429 | 73.7% | | |
| Average Number of Visits | 7.5 | | 11.2 | | 11.9 | | 9.3 | | 5.8 | | 8.5 | | 8.4 | | 1.8 | | 0.0 | |
| Treatment Change (from first to last SS visit) ⁶ | | | | | | | | | | | | | | | | | | |
| Insulin Added | 2192 | 59.1% | 2134 | 100.0% | 58 | 54.2% | | | | | | | | | | | | |
| Oral Medication Added | 1518 | 40.9% | | | 49 | 45.8% | 1469 | 100.0% | | | | | | | | | | |
| BMI less than 19.8 | 569 | 4.5% | 48 | 1.8% | | | 47 | 2.5% | 473 | 5.9% | | | 34 | 5.2% | 16 | 2.7% | | |
| Total Gain < 28 lbs ⁷ | 316 | 55.5% | 20 | 41.7% | | | 30 | 63.8% | 266 | 56.2% | | | 17 | 50.0% | 11 | 68.8% | | |
| Total Gain 28 - 40 lbs | 198 | 34.8% | 20 | 41.7% | | | 14 | 29.8% | 163 | 34.5% | | | 14 | 41.2% | 5 | 31.3% | | |
| Total Gain > 40 lbs | 55 | 9.7% | 8 | 16.7% | | | | | 44 | 9.3% | | | | | | | | |
| BMI 19.8 to 24.9 | 3149 | 25.0% | 441 | 16.6% | 13 | 18.6% | 336 | 18.0% | 2359 | 29.5% | 23 | 31.9% | 210 | 31.9% | 63 | 10.8% | | |
| Total Gain < 25 lbs | 1534 | 48.7% | 203 | 46.0% | 8 | 61.5% | 139 | 41.4% | 1184 | 50.2% | 14 | 60.9% | 98 | 46.7% | 45 | 71.4% | | |
| Total Gain 25 - 35 lbs | 1051 | 33.4% | 149 | 33.8% | | | 120 | 35.7% | 778 | 33.0% | 8 | 34.8% | 71 | 33.8% | 12 | 19.0% | | |
| Total Gain > 35 lbs | 564 | 17.9% | 89 | 20.2% | | | 77 | 22.9% | 397 | 16.8% | | | 41 | 19.5% | 6 | 9.5% | | |
| BMI 25.0 to 29.0 | 3082 | 24.5% | 555 | 20.8% | 15 | 21.4% | 440 | 23.6% | 2072 | 25.9% | 18 | 25.0% | 154 | 23.4% | 80 | 13.7% | | |
| Total Gain < 15 lbs | 961 | 31.2% | 161 | 29.0% | | | 140 | 31.8% | 657 | 31.7% | 7 | 38.9% | 37 | 24.0% | 55 | 68.8% | | |
| Total Gain 15 - 25 lbs | 1045 | 33.9% | 185 | 33.3% | | | 152 | 34.5% | 704 | 34.0% | 10 | 55.6% | 55 | 35.7% | 15 | 18.8% | | |
| Total Gain > 25 lbs | 1076 | 34.9% | 209 | 37.7% | 8 | 53.3% | 148 | 33.6% | 711 | 34.3% | | | 62 | 40.3% | 10 | 12.5% | | |
| BMI > 29.0 | 5776 | 45.8% | 1617 | 60.7% | 41 | 58.6% | 1040 | 55.7% | 3078 | 38.5% | 27 | 37.5% | 260 | 39.5% | 162 | 27.8% | | |
| Total Gain < 15 lbs | 2855 | 49.4% | 705 | 43.6% | 19 | 46.3% | 534 | 51.3% | 1597 | 51.9% | 19 | 70.4% | 117 | 45.0% | 129 | 79.6% | | |
| Total Gain 15 - 25 lbs | 1630 | 28.2% | 481 | 29.7% | 10 | 24.4% | 271 | 26.1% | 868 | 28.2% | 5 | 18.5% | 81 | 31.2% | 19 | 11.7% | | |
| Total Gain > 25 lbs | 1291 | 22.4% | 431 | 26.7% | 12 | 29.3% | 235 | 22.6% | 613 | 19.9% | | | 62 | 23.8% | 14 | 8.6% | | |
| BMI or Wt. Gain Unknown | 24 | 0.2% | | | | | | | 18 | 0.2% | | | | | 261 | 44.8% | | |

1. Cell values less than 5 are suppressed.

3. Tables by specific maternal characteristics (e.g. race/ethnicity, education, payment source) exclude mothers with spontaneous abortions.

6. Treatment change is reported for those clients remaining in the same general diagnostic grouping whose intervention at the last visit changed to include oral medications or insulin.

7. Total gain is based on the difference in weight measured at last SS visit and the pre-pregnancy weight reported.

CALIFORNIA DIABETES AND PREGNANCY PROGRAM - CALENDAR YEAR 2008

DATA REPORT for deliveries between January 1, 2008 and by December 31, 2008

Characteristics by Maternal Type of Diabetes and Intervention at Last Sweet Success Visit (number and percent)¹

| Generated 03/17/2011 14:28:02 | | | | | | | | | | | | | | | | | | |
|---|-------|-------|---------|-------|----------------|-------|-----------|-------|-------|-------|---------------------|-------|----------------------------|-------|---------|-------|-------|---|
| Gestational Diabetes Mellitus (GDM) | | | | | | | | | | | | | | | | | | |
| State of California | Total | | Insulin | | Insulin & Oral | | Oral Meds | | Diet | | History of GDM Diet | | Impaired Glucose Tol. Diet | | Unknown | | None | |
| | num # | % | num # | % | num # | % | num # | % | num # | % | num # | % | num # | % | num # | % | num # | % |
| Type of Pregnancy⁸ | | | | | | | | | | | | | | | | | | |
| Singletons | 11908 | 94.4% | 2554 | 95.9% | 66 | 94.3% | 1784 | 95.5% | 7504 | 93.7% | 70 | 97.2% | 625 | 94.8% | 570 | 97.9% | | |
| Multiples | 446 | 3.5% | 92 | 3.5% | | | 55 | 2.9% | 299 | 3.7% | | | 31 | 4.7% | 6 | 1.0% | | |
| - Twins | 427 | 3.4% | 86 | 3.2% | | | 52 | 2.8% | 289 | 3.6% | | | 28 | 4.2% | | | | |
| - Triplets or more | 19 | 0.2% | 6 | 0.2% | | | | | 10 | 0.1% | | | | | | | | |
| - Unknown multiples | | | | | | | | | | | | | | | | | | |
| Unknown | 260 | 2.1% | 18 | 0.7% | | | 30 | 1.6% | 208 | 2.6% | | | | | 6 | 1.0% | | |
| Gender⁹ | | | | | | | | | | | | | | | | | | |
| Male | 5165 | 43.4% | 1241 | 48.6% | 28 | 42.4% | 750 | 42.0% | 3146 | 41.9% | 29 | 41.4% | 287 | 45.9% | 91 | 16.0% | | |
| Female | 4571 | 38.4% | 992 | 38.8% | 30 | 45.5% | 728 | 40.8% | 2821 | 37.6% | 31 | 44.3% | 287 | 45.9% | 82 | 14.4% | | |
| Unknown | 2172 | 18.2% | 321 | 12.6% | 8 | 12.1% | 306 | 17.2% | 1537 | 20.5% | 10 | 14.3% | 51 | 8.2% | 397 | 69.6% | | |
| Congenital Anomalies | | | | | | | | | | | | | | | | | | |
| Known at Discharge | 98 | 71.0% | 31 | 31.6% | | | 17 | 17.3% | 49 | 50.0% | | | 7 | 5.1% | | | | |
| Method of Feeding at Discharge from Hospital | | | | | | | | | | | | | | | | | | |
| Breast Milk Only | 3944 | 33.1% | 820 | 32.1% | 21 | 31.8% | 625 | 35.0% | 2478 | 33.0% | 32 | 45.7% | 324 | 51.8% | 96 | 16.8% | | |
| Formula Only | 1124 | 9.4% | 250 | 9.8% | 10 | 15.2% | 169 | 9.5% | 695 | 9.3% | | | 47 | 7.5% | 14 | 2.5% | | |
| Breast milk & formula | 2518 | 21.1% | 592 | 23.2% | 17 | 25.8% | 368 | 20.6% | 1541 | 20.5% | 13 | 18.6% | 120 | 19.2% | 42 | 7.4% | | |
| Unknown | 4322 | 36.3% | 892 | 34.9% | 18 | 27.3% | 622 | 34.9% | 2790 | 37.2% | 23 | 32.9% | 134 | 21.4% | 418 | 73.3% | | |
| Singletons | | | | | | | | | | | | | | | | | | |
| Birthweight | | | | | | | | | | | | | | | | | | |
| < 1,500 grams | 45 | 0.4% | 8 | 0.3% | | | | | 33 | 0.4% | | | 10 | 1.6% | | | | |
| 1,500 to 2,499 grams | 565 | 4.7% | 135 | 5.3% | | | 71 | 4.0% | 359 | 4.8% | | | 26 | 4.2% | 9 | 1.6% | | |
| 2,500 to 3,999 grams | 8090 | 67.9% | 1795 | 70.3% | 43 | 65.2% | 1199 | 67.2% | 5053 | 67.3% | 51 | 72.9% | 474 | 75.8% | 146 | 25.6% | | |
| 4,000 to 4,499 grams | 741 | 6.2% | 206 | 8.1% | 11 | 16.7% | 132 | 7.4% | 392 | 5.2% | | | 48 | 7.7% | 14 | 2.5% | | |
| > 4,500 grams | 196 | 1.6% | 60 | 2.3% | | | 39 | 2.2% | 93 | 1.2% | | | 9 | 1.4% | 5 | 0.9% | | |
| Unknown | 2271 | 19.1% | 350 | 13.7% | 8 | 12.1% | 339 | 19.0% | 1574 | 21.0% | 10 | 14.3% | 58 | 9.3% | 396 | 69.5% | | |
| Completed Weeks Gestation at Delivery¹⁰ | | | | | | | | | | | | | | | | | | |
| 20 to 24 weeks | 6 | 0.1% | | | | | | | | | | | | | | | | |
| 24 to 28 weeks | 35 | 0.3% | 13 | 0.5% | | | | | 18 | 0.2% | | | 8 | 1.3% | | | | |
| 29 to 32 weeks | 57 | 0.5% | 15 | 0.6% | | | 8 | 0.4% | 34 | 0.5% | | | | | | | | |
| 33 to 36 weeks | 774 | 6.5% | 191 | 7.5% | 5 | 7.6% | 110 | 6.2% | 468 | 6.2% | 6 | 8.6% | 36 | 5.8% | 10 | 1.8% | | |
| 37 to 41 weeks | 9033 | 75.9% | 2080 | 81.4% | 52 | 78.8% | 1374 | 77.0% | 5527 | 73.7% | 50 | 71.4% | 517 | 82.7% | 121 | 21.2% | | |
| > 41 weeks | 173 | 1.5% | 24 | 0.9% | | | 27 | 1.5% | 121 | 1.6% | | | 15 | 2.4% | | | | |
| Unknown | 1830 | 15.4% | 229 | 9.0% | 8 | 12.1% | 259 | 14.5% | 1334 | 17.8% | 12 | 17.1% | 45 | 7.2% | 434 | 76.1% | | |

1. Cell values less than 5 are suppressed.

8. Tables by type of pregnancy and infant characteristics exclude spontaneous abortions

9. Tables by specific infant characteristics (e.g. gender, congenital anomalies, method of feeding, birthweight, completed weeks gestation, birthweight for gestational age, and not discharged with mother) are for singleton infants not spontaneously aborted.

10. Completed weeks gestation at delivery calculated using estimated date of delivery and actual date of delivery (Birth Date).

If either value is missing, case is reported as unknown.

CALIFORNIA DIABETES AND PREGNANCY PROGRAM - CALENDAR YEAR 2008

DATA REPORT for deliveries between January 1, 2008 and by December 31, 2008

Characteristics by Maternal Type of Diabetes and Intervention at Last Sweet Success Visit (number and percent)¹

| | | | | | | | | | | | | | | | | | | |
|-------------------------------|--|---|---------|---|----------------|---|-----------|---|-------|---|---------------------|---|----------------------------|---|---------|---|-------|---|
| Generated 03/17/2011 14:28:02 | Gestational Diabetes Mellitus (GDM) | | | | | | | | | | | | | | | | | |
| State of California | Total | | Insulin | | Insulin & Oral | | Oral Meds | | Diet | | History of GDM Diet | | Impaired Glucose Tol. Diet | | Unknown | | None | |
| | num # | % | num # | % | num # | % | num # | % | num # | % | num # | % | num # | % | num # | % | num # | % |

| Birthweight (Size) for Gestational Age^{9,11} | | | | | | | | | | | | | | | | | | |
|--|------|-------|------|-------|----|-------|------|-------|------|-------|----|-------|-----|-------|-----|-------|--|--|
| Small (<10th %tile) | 1160 | 9.7% | 240 | 9.4% | | | 156 | 8.7% | 760 | 10.1% | 7 | 10.0% | 68 | 10.9% | 14 | 2.5% | | |
| Average (11-89th %tile) | 7288 | 61.2% | 1636 | 64.1% | 38 | 57.6% | 1071 | 60.0% | 4543 | 60.5% | 41 | 58.6% | 430 | 68.8% | 94 | 16.5% | | |
| Large (>90 %tile) | 939 | 7.9% | 279 | 10.9% | 12 | 18.2% | 183 | 10.3% | 465 | 6.2% | 6 | 8.6% | 57 | 9.1% | 14 | 2.5% | | |
| Unknown | 2521 | 21.2% | 399 | 15.6% | 12 | 18.2% | 374 | 21.0% | 1736 | 23.1% | 16 | 22.9% | 70 | 11.2% | 448 | 78.6% | | |
| Not Discharged with Mother | | | | | | | | | | | | | | | | | | |
| Total | 768 | 74.3% | 217 | 28.3% | 7 | 0.9% | 147 | 19.1% | 397 | 51.7% | | | 37 | 3.6% | 16 | 1.5% | | |
| Length of Stay < 7 days | 259 | 33.7% | 84 | 38.7% | | | 49 | 33.3% | 125 | 31.5% | | | 14 | 37.8% | 6 | 37.5% | | |
| Length of Stay > 7 days | 207 | 27.0% | 60 | 27.6% | | | 43 | 29.3% | 103 | 25.9% | | | 13 | 35.1% | 6 | 37.5% | | |
| Unknown | 302 | 39.3% | 73 | 33.6% | 5 | 71.4% | 55 | 37.4% | 169 | 42.6% | | | 10 | 27.0% | | | | |

1. Cell values less than 5 are suppressed.

9. Tables by specific infant characteristics (e.g. gender, congenital anomalies, method of feeding, birthweight, completed weeks gestation, birthweight for gestational age, and not discharged with mother) are for singleton infants not spontaneously aborted.

11. Birthweight (size) for gestational age is calculated using infant birthweight and gestational age as calculated by difference of estimated date of delivery (EDD) and actual Delivery Date (DD) when compared to standard U.S. growth charts. If birthweight, EDD, or DD is missing, case is reported as unknown.