

Table 8. Rates of central line-associated bloodstream infection and central line insertion practices (CLIP) adherence percent in patient care locations providing burn critical care, reported by California hospitals, January - December 2012 (inclusive).

Reporting hospital*	Cases	Central line-days	Rate	95% Confidence Interval	Statistical interpretation**	CLIP Adherence %
STATE OF CALIFORNIA POOLED DATA	37	12772	2.90			97.1
ARROWHEAD RGN MED CTR, COLTON	0	480	0.00	(0.00,6.24)	⊙	100
COMMUNITY REGIONAL MED CTR, FRESNO	9	1742	5.17	(2.36,9.81)	⊙	85.7
LAC+USC MED CTR, LOS ANGELES	0	1033	0.00	(0.00,2.90)	⊙	100
SAINTE FRANCIS MEMORIAL HOSP, SAN FRANCISCO	4	1207	3.31	(0.90,8.49)	⊙	97.6
SAN JOAQUIN COMMUNITY HOSP, BAKERSFIELD	5	219	22.83	(7.41,53.28)	●	60.0
SANTA CLARA VALLEY MED CTR, SAN JOSE	1	491	2.04	(0.05,11.35)	⊙	97.0
TORRANCE MEMORIAL MED CTR, TORRANCE	0	1263	0.00	(0.00,2.37)	○	ND
UNIVERSITY OF CALIFORNIA DAVIS MED CTR, SACRAMENTO	7	3026	2.31	(0.93,4.77)	⊙	97.2
UNIVERSITY OF CALIFORNIA IRVINE MED CTR, ORANGE	3	1335	2.25	(0.46,6.57)	⊙	100
† UNIVERSITY OF CALIFORNIA, SAN DIEGO	5	1425	3.51	(1.14,8.19)	⊙	98.6
UNIVERSITY OF CALIFORNIA, SAN DIEGO MED CTR						
UCSD-LA JOLLA, JOHN M.& SALLY B. THORNTON HOSP						
WEST HILLS HOSP AND MED CTR	3	453	6.62	(1.37,19.35)	⊙	100
WESTERN MED CTR SANTA ANA	0	98	0.00	(0.00,30.57)	⊙	100

* Hospitals with only one acute care campus associated with their license were listed by the campus name; hospitals with more than one campus associated with their license were listed by licensee names with acute care campus name(s) indented below

**Statistical interpretation indicates hospital-specific rates that were statistically higher (●), lower (○), or no different (⊙) than California average rates, or rates that were not tested due to too few central line-days or too few hospitals reported (†)

† Hospital reported infection data aggregated over all acute care campuses

‡ Hospital reported infection data separately by acute care campus

ND: No CLIP data reported.

Notes: Rate per 1000 central line days; 95% confidence interval calculated for rates based on exact Poisson distribution; burn critical care locations specialize in care of critically ill patients with significant/major burns

Source: Central Line-Associated Bloodstream Infections and Central Line Insertion Practices in California Hospitals, January through December 2012. California Department of Public Health