

CLABSI Table 9. Rates of central line-associated bloodstream infection and central line insertion practices (CLIP) adherence percent in patient care locations providing trauma critical care, reported by California hospitals, January - December 2013 (inclusive).

Reporting hospital*	Cases	Central line-days	Rate	95% Confidence Interval	Statistical interpretation**	CLIP Adherence %
STATE OF CALIFORNIA POOLED DATA	25	26372	0.95			98.4
CALIFORNIA HOSP MED CTR, LOS ANGELES	0	847	0.00	(0.00,3.54)	⊙	96.8
DESERT REGIONAL MED CTR, PALM SPRINGS	0	1274	0.00	(0.00,2.35)	⊙	66.7
LAC/HARBOR-UCLA MED CTR, TORRANCE	7	2388	2.93	(1.18,6.04)	●	98.2
MERCY SAN JUAN MED CTR, CARMICHAEL	3	1137	2.64	(0.54,7.71)	⊙	100
† PALOMAR HEALTH	0	2039	0.00	(0.00,1.47)	⊙	99.5
PALOMAR HEALTH DOWNTOWN CAMPUS, ESCONDIDO
PALOMAR MED CTR, ESCONDIDO
REGIONAL MED CTR OF SAN JOSE	0	1235	0.00	(0.00,2.43)	⊙	ND
SAN FRANCISCO GENERAL HOSP	3	3119	0.96	(0.20,2.81)	⊙	100
SANTA CLARA VALLEY MED CTR, SAN JOSE	0	1105	0.00	(0.00,2.71)	⊙	100
SCRIPPS MEMORIAL HOSP, LA JOLLA	1	1957	0.51	(0.01,2.85)	⊙	96.2
SUTTER ROSEVILLE MED CTR	0	1979	0.00	(0.00,1.51)	⊙	99.7
UNIVERSITY OF CALIFORNIA DAVIS MED CTR, SACRAMENTO	8	4748	1.68	(0.73,3.32)	⊙	97.7
UNIVERSITY OF CALIFORNIA IRVINE MED CTR, ORANGE	3	4544	0.66	(0.14,1.93)	⊙	97.9

* Hospitals with only one acute care campus associated with their license were listed by the campus name; hospitals with more than one campus associated with their license were listed by licensee names with acute care campus name(s) indented below

**Statistical interpretation indicates hospital-specific rates that were statistically higher (●), lower (○), or no different (⊙) than California average rates, or rates that were not tested due to too few central line-days or too few hospitals reported in the location category (†)

† Hospital reported infection data aggregated over all acute care campuses

‡ Hospital reported infection data separately by acute care campus

ND: No CLIP data reported.

Notes: Rate per 1000 central line days; 95% confidence interval calculated for rates based on exact Poisson distribution; trauma critical care locations specialize in care of critically ill patients who require a high level of monitoring and/or intervention following trauma or during critical illness related to trauma

Source: Central Line-Associated Bloodstream Infections and Central Line Insertion Practices in California Hospitals, January through December 2013. California Department of Public Health