

**CLABSI Table 18. Rates of central line-associated bloodstream infection in patient care locations providing neonatal step-down care, reported by California hospitals, January - December 2013 (inclusive)**

Reporting hospital*	Cases	Central line-days	Rate	95% Confidence Interval	Statistical interpretation**
<b>STATE OF CALIFORNIA POOLED DATA</b>	<b>0</b>	<b>1094</b>	<b>0.00</b>		
† ALAMEDA COUNTY MED CTR	0	1	-----	-----	+
ALAMEDA COUNTY MED CTR, OAKLAND	.	.	.		
ALAMEDA COUNTY MED CTR-FAIRMONT CAMPUS, SAN LEANDRO	.	.	.		
COMMUNITY HOSP OF THE MONTEREY PENINSULA, MONTEREY	0	2	-----	-----	+
KAISER FOUNDATION HOSP, FRESNO	0	8	-----	-----	+
KAISER FOUNDATION HOSP, SANTA ROSA	0	6	-----	-----	+
KAISER FOUNDATION HOSP-SOUTH SACRAMENTO	0	9	-----	-----	+
LUCILE SALTER PACKARD CHILDREN'S HOSP AT STANFORD, PALO ALTO	0	88	0.00	(0.00,34.04)	⊙
MEMORIAL MED CTR, MODESTO	0	6	-----	-----	+
METHODIST HOSP OF SACRAMENTO	0	98	0.00	(0.00,30.57)	⊙
POMERADO HOSP, POWAY	0	7	-----	-----	+
SHARP CHULA VISTA MED CTR	0	84	0.00	(0.00,35.66)	⊙
SUTTER DELTA MED CTR, ANTIOCH	0	12	-----	-----	+
UNIVERSITY OF CALIFORNIA DAVIS MED CTR, SACRAMENTO	0	747	0.00	(0.00,4.01)	⊙
WATSONVILLE COMMUNITY HOSP	0	26	-----	-----	+

\* Hospitals with only one acute care campus associated with their license were listed by the campus name; hospitals with more than one campus associated with their license were listed by licensee names with acute care campus name(s) indented below

\*\*Statistical interpretation indicates hospital-specific rates that were statistically higher (●), lower (○), or no different (⊙) than California average rates, or rates that were not tested due to too few central line-days or too few hospitals reported in the location category (†)

† Hospital reported infection data aggregated over all acute care campuses

‡ Hospital reported infection data separately by acute care campus

Notes: Rate per 1000 central line days; 95% confidence interval calculated for rates based on exact Poisson distribution; ----- indicates data not shown due to too few central line-days reported; neonatal step down general care locations specialize in preterm and/or ill infants with birth weight ≥1500 grams requiring resuscitation and stabilization before transfer to a facility at which newborn intensive care is provided

Source: Central Line-Associated Bloodstream Infections and Central Line Insertion Practices in California Hospitals, January through December 2013. California Department of Public Health