

CLABSI Table 6. Central Line-Associated Bloodstream Infection (CLABSI) Rates and Central Line Insertion Practices (CLIP) Adherence Percentages in Medical/Surgical Critical Care Units, Reported by California Major Teaching Hospitals, January - December 2014

Hospitals highlighted in red did not participate in 2014 data validation; completeness of reported infections has not been evaluated

County	Hospital Name	CLABSI count	Central line-days	Rate per 1000*	95% Confidence Interval**	Statistical Interpretation***	CLIP Adherence %^
	STATE OF CALIFORNIA POOLED DATA	67	57765	1.16			99.2
Alameda	Highland Hospital	3	3339	0.9	(0.19,2.63)	N	99.5
Fresno	Community Regional Medical Center	19	7224	2.63	(1.58,4.11)	H	99.7
Kern	Kern Medical Center	0	2058	0	(0.00,1.46)	N	100
Los Angeles	Cedars-Sinai Medical Center	2	955	2.09	(0.25,7.57)	N	98.6
Los Angeles	Kaiser Foundation Hospital, Los Angeles	1	3231	0.31	(0.01,1.72)	N	96.7
Los Angeles	Keck Hospital of USC	6	2989	2.01	(0.74,4.37)	N	98.2
<i>Los Angeles</i>	<i>Santa Monica - UCLA Medical Center and Orthopaedic Hospital</i>	<i>3</i>	<i>2884</i>	<i>1.04</i>	<i>(0.21,3.04)</i>	<i>N</i>	<i>99.7</i>
Riverside	Eisenhower Medical Center	2	3579	0.56	(0.07,2.02)	N	99.4
Riverside	Riverside County Regional Medical Center	1	5436	0.18	(0.00,1.02)	L	100
Sacramento	University of California Davis Medical Center	2	1416	1.41	(0.17,5.10)	N	99
San Bernardino	Loma Linda University Medical Center	1	261	3.83	(0.10,21.35)	N	100
San Diego	Scripps Mercy Hospital	0	28	.	-----	---	100
San Diego	University of California, San Diego Medical Center	8	6610	1.21	(0.52,2.38)	N	97.1
San Francisco	Kaiser Foundation Hospital, San Francisco	0	1152	0	(0.00,2.60)	N	98.9
San Francisco	UCSF Medical Center	8	6001	1.33	(0.58,2.63)	N	99.8
Santa Clara	O'Connor Hospital	2	2973	0.67	(0.08,2.43)	N	96.6
Santa Clara	Stanford Hospital	9	7629	1.18	(0.54,2.24)	N	99.8

*Rate per 1000 central line days

**95% confidence interval calculated for rates based on exact Poisson distribution.

*** Statistical interpretation indicates hospital-specific rates that were statistically higher (H), lower (L), or no different (N) than California average rates, or rates that were not tested due to too few central line-days or too few hospitals reported (---).

^The CLIP bundle includes eight components: hand hygiene, recommend skin preparation agent, allowing the skin preparation agent to dry, and five maximal sterile barriers (cap, mask, gown, gloves, and drape).

Source: Central Line-Associated Bloodstream Infections and Central Line Insertion Practices in California Hospitals, January 2014 through December 2014 (Inclusive).