

CLABSI Table 45. Central Line-Associated Bloodstream Infection (CLABSI) Rates Related to Temporary-Type Central Lines in Solid Organ Transplant Care Units, Reported by California Hospitals, January - December 2014

Hospitals highlighted in red did not participate in 2014 data validation; completeness of reported infections has not been evaluated.

County	Hospital Name	CLABSI count	Central line-days	Rate per 1000*	95% Confidence Interval**	Statistical Interpretation***
	STATE OF CALIFORNIA POOLED DATA	13	14286	0.91		
Los Angeles	Keck Hospital of USC	4	2035	1.97	(0.54,5.03)	---
<i>Los Angeles</i>	<i>LAC/Harbor-UCLA Medical Center</i>	<i>0</i>	<i>332</i>	<i>0</i>	<i>(0.00,9.02)</i>	<i>---</i>
Los Angeles	Ronald Reagan UCLA Medical Center	7	5284	1.32	(0.53,2.73)	---
Los Angeles	Saint Vincent Medical Center	0	415	0	(0.00,7.22)	---
Riverside	Riverside Community Hospital	0	1963	0	(0.00,1.53)	---
San Diego	Sharp Memorial Hospital	2	2155	0.93	(0.11,3.35)	---
San Francisco	UCSF Medical Center	0	2102	0	(0.00,1.43)	---

*Rate per 1000 central line days.

**95% confidence interval calculated for rates based on exact Poisson distribution.

*** Statistical interpretation indicates hospital-specific rates that were statistically higher (H), lower (L), or no different (N) than California average rates, or rates that were not tested due to too few central line-days or too few hospitals reported (---).

Source: Central Line-Associated Bloodstream Infections and Central Line Insertion Practices in California Hospitals, January 2014 through December 2014 (Inclusive).