

CLABSI Table 37. Central Line-Associated Bloodstream Infection (CLABSI) Rates Related to Permanent -Type Central Lines in Bone Marrow Transplant Pediatric Care Units, Reported by California Hospitals, January-December 2015

County	Hospital Name	CLABSI count	Central line days	Rate per 1000*	95% Confidence Interval**	Statistical Interpretation***
	STATE OF CALIFORNIA POOLED DATA	20	8121	2.46		
Alameda	Children's Hospital & Research Center Oakland	1	774	1.29	(0.03,7.20)	
Los Angeles	Children's Hospital Los Angeles	5	3514	1.42	(0.46,3.32)	
Los Angeles	Ronald Reagan UCLA Medical Center	3	1473	2.04	(0.42,5.95)	
San Francisco	UCSF Medical Center	11	2360	4.66	(2.33,8.34)	

*CLABSI rate per 1000 central line days.

**95% confidence interval calculated for rates based on exact Poisson distribution.

*** Statistical interpretation indicates hospital-specific rates that were statistically higher, lower, or no different than California average rates, or rates that were not tested due to too few central line days or too few hospitals reported .

Source: Central Line-Associated Bloodstream Infections and Central Line Insertion Practices in California Hospitals, January through December 2015. California Department of Public Health.