

CLABSI Table 43. Central Line-Associated Bloodstream Infection (CLABSI) Rates Related to Temporary -Type Central Lines in Bone Marrow Transplant Care Units, Reported by California Hospitals, January-December 2015

County	Hospital Name	CLABSI count	Central line days	Rate per 1000*	95% Confidence Interval**	Statistical Interpretation***
	STATE OF CALIFORNIA POOLED DATA	159	42549	3.74		
Los Angeles	City of Hope Helford Clinical Research Hospital	54	15834	3.41	(2.56,4.45)	
Los Angeles	Ronald Reagan UCLA Medical Center	42	6615	6.35	(4.58,8.58)	
Sacramento	Sutter Medical Center, Sacramento	13	4388	2.96	(1.58,5.07)	
Sacramento	University of California Davis Medical Center	17	4449	3.82	(2.23,6.12)	
San Diego	University of California, San Diego Medical Center	30	7142	4.20	(2.83,6.00)	
San Francisco	UCSF Medical Center	0	384	0.00	(0.00,7.80)	
Santa Clara	Stanford Health Care	3	3737	0.80	(0.17,2.35)	

*CLABSI rate per 1000 central line days.

**95% confidence interval calculated for rates based on exact Poisson distribution.

*** Statistical interpretation indicates hospital-specific rates that were statistically higher, lower, or no different than California average rates, or rates that were not tested due to too few central line days or too few hospitals reported .

Source: Central Line-Associated Bloodstream Infections and Central Line Insertion Practices in California Hospitals, January through December 2015. California Department of Public Health.