

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
ADVENTIST MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
ADVENTIST MEDICAL CENTER - REEDLEY	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	79
	Surgery patients who were given the right kind of antibiotic to help prevent infection	93
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
AHMC ANAHEIM REGIONAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	93
ALAMEDA COUNTY MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	**NSPP
	Surgery patients who were given the right kind of antibiotic to help prevent infection	**NSPP
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	**NSPP

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
ALAMEDA HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	93
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	88
ALHAMBRA HOSPITAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	95
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	93
ALTA BATES SUMMIT MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
ALTA BATES SUMMIT MEDICAL CENTER - ALTA BATES CAMP	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
ALVARADO HOSPITAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
ANAHEIM GENERAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
ANTELOPE VALLEY HOSPITAL MED CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
ARROWHEAD REGIONAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
ARROYO GRANDE COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
BAKERSFIELD HEART HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
BAKERSFIELD MEMORIAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
BANNER LASSEN MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	94
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
BARSTOW COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
BARTON MEMORIAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
BEAR VALLEY COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
BELLFLOWER MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	80
	Surgery patients who were given the right kind of antibiotic to help prevent infection	87
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	74

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
BEVERLY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
BIGGS GRIDLEY MEMORIAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
BROTMAN MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	95
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	89
CALIFORNIA HOSPITAL MEDICAL CENTER LA	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
CALIFORNIA PACIFIC MEDICAL CTR - ST. LUKE'S CAMPUS	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
CALIFORNIA PACIFIC MEDICAL CTR-DAVIES CAMPUS HOSP	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
CALIFORNIA PACIFIC MEDICAL CTR-PACIFIC CAMPUS HOSP	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
CATALINA ISLAND MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
CEDARS-SINAI MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
CENTINELA HOSPITAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
CENTRAL VALLEY GENERAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
CHAPMAN MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	90

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
CHILDREN'S HOSP OF LOS ANGELES	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
CHINESE HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	84
CHINO VALLEY MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
CITRUS VALLEY MEDICAL CENTER-IC CAMPUS	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
CLOVIS COMMUNITY MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
COALINGA REGIONAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
COAST PLAZA HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
COASTAL COMMUNITIES HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
COLLEGE HOSPITAL COSTA MESA	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
COLORADO RIVER MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
COLUSA REGIONAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
COMMUNITY AND MISSION HOSPITAL OF HUNTINGTON PARK	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	81
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
COMMUNITY HOSPITAL OF LONG BEACH	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	91
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
COMMUNITY HOSPITAL OF SAN BERNARDINO	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	94
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	88
COMMUNITY HOSPITAL OF THE MONTEREY PENINSULA	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
COMMUNITY MEMORIAL HOSPITAL SAN BUENAVENTURA	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	94
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
COMMUNITY REGIONAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
CONTRA COSTA REGIONAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	89
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	86
CORCORAN DISTRICT HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
CORONA REGIONAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	95
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
DAMERON HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
DELANO REGIONAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	85
	Surgery patients who were given the right kind of antibiotic to help prevent infection	80
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
DESERT REGIONAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
DESERT VALLEY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	92
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
DOCTORS HOSPITAL OF MANTECA	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
DOCTORS HOSPITAL OF WEST COVINA, INC	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	82
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	84
DOCTORS MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
DOCTORS MEDICAL CENTER-SAN PABLO	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
DOMINICAN HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
DOWNEY REGIONAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
EAST LOS ANGELES DOCTORS HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	94
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
EAST VALLEY HOSPITAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
EASTERN PLUMAS HOSPITAL - PORTOLA CAMPUS	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
EDEN MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
EISENHOWER MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
EL CAMINO HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
EL CENTRO REGIONAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	91
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
EMANUEL MEDICAL CENTER INC	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
ENCINO HOSPITAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	**NSPP
	Surgery patients who were given the right kind of antibiotic to help prevent infection	**NSPP
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	**NSPP
ENLOE MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
FAIRCHILD MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	92
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
FALLBROOK HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
FEATHER RIVER HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
FOOTHILL PRESBYTERIAN HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
FOUNTAIN VALLEY REGIONAL HOSPITAL & MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
FRANK R HOWARD MEMORIAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
FRENCH HOSPITAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
FRESNO HEART AND SURGICAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
FRESNO SURGICAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
FRESNO VA MEDICAL CENTER (VA CENTRAL CALIFORNIA HEALTHCARE SYSTEM)	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
GARDEN GROVE HOSPITAL & MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
GARFIELD MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
GEORGE L MEE MEMORIAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	71
	Surgery patients who were given the right kind of antibiotic to help prevent infection	95
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
GLENDALE ADVENTIST MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	93
GLENDALE MEM HOSPITAL & HLTH CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
GLENN MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
GOLETA VALLEY COTTAGE HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
GOOD SAMARITAN HOSPITAL, BAKERSFIELD	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	**NSPP
	Surgery patients who were given the right kind of antibiotic to help prevent infection	**NSPP
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	**NSPP
GOOD SAMARITAN HOSPITAL, LOS ANGELES	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	93
GOOD SAMARITAN HOSPITAL, SAN JOSE	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
GREATER EL MONTE COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	92
	Surgery patients who were given the right kind of antibiotic to help prevent infection	92
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
GROSSMONT HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
HAZEL HAWKINS MEMORIAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	95
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	91
HEALDSBURG DISTRICT HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	93
	Surgery patients who were given the right kind of antibiotic to help prevent infection	93
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	89

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
HEMET VALLEY MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	90
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94
HENRY MAYO NEWHALL MEMORIAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94
HI-DESERT MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	89
	Surgery patients who were given the right kind of antibiotic to help prevent infection	77
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	93
HOAG MEMORIAL HOSPITAL PRESBYTERIAN	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
HOAG ORTHOPEDIC INSTITUTE	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
HOLLYWOOD COMMUNITY HOSPITAL OF HOLLYWOOD	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	**NSPP
	Surgery patients who were given the right kind of antibiotic to help prevent infection	**NSPP
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	**NSPP
HOLLYWOOD PRESBYTERIAN MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
HUNTINGTON BEACH HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
HUNTINGTON MEMORIAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
JEROLD PHELPS COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
JOHN C FREMONT HEALTHCARE DISTRICT	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
JOHN F KENNEDY MEMORIAL HOSPITAL, INC	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
JOHN MUIR MEDICAL CENTER - CONCORD CAMPUS	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
JOHN MUIR MEDICAL CENTER - WALNUT CREEK CAMPUS	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
KAISER FOUNDATION HOSP SO SACRAMENTO	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
KAISER FOUNDATION HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
KAISER FOUNDATION HOSPITAL - ANTIOCH	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
KAISER FOUNDATION HOSPITAL - BALDWIN PARK	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
KAISER FOUNDATION HOSPITAL - DOWNEY	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
KAISER FOUNDATION HOSPITAL - FRESNO	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
KAISER FOUNDATION HOSPITAL - LOS ANGELES	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
KAISER FOUNDATION HOSPITAL - PANORAMA CITY	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
KAISER FOUNDATION HOSPITAL - REDWOOD CITY	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
KAISER FOUNDATION HOSPITAL - ROSEVILLE	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
KAISER FOUNDATION HOSPITAL - SACRAMENTO	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
KAISER FOUNDATION HOSPITAL - SAN DIEGO	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
KAISER FOUNDATION HOSPITAL - SAN FRANCISCO	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
KAISER FOUNDATION HOSPITAL - SOUTH BAY	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
KAISER FOUNDATION HOSPITAL - SOUTH SAN FRANCISCO	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
KAISER FOUNDATION HOSPITAL - VACAVILLE	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
KAISER FOUNDATION HOSPITAL - WALNUT CREEK	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
KAISER FOUNDATION HOSPITAL - WEST LA	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
KAISER FOUNDATION HOSPITAL - WOODLAND HILLS	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
KAISER FOUNDATION HOSPITAL ANAHEIM	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
KAISER FOUNDATION HOSPITAL AND REHAB CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
KAISER FOUNDATION HOSPITAL FONTANA	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
KAISER FOUNDATION HOSPITAL FREMONT/HAYWARD	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
KAISER FOUNDATION HOSPITAL MANTECA	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
KAISER FOUNDATION HOSPITAL OAKLAND/RICHMOND	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
KAISER FOUNDATION HOSPITAL, RIVERSIDE	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
KAISER FOUNDATION HOSPITAL-MORENO VALLEY	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
KAISER FOUNDATION HOSPITAL-SAN JOSE	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
KAISER FOUNDATION HOSPITAL-SANTA CLARA	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
KAISER FOUNDATION HOSPITAL-SANTA ROSA	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
KAWEAH DELTA MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	88
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	93
KECK HOSPITAL OF USC	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
KERN MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	93
	Surgery patients who were given the right kind of antibiotic to help prevent infection	86
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	79
KERN VALLEY HEALTHCARE DISTRICT	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
LA PALMA INTERCOMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	95
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
LAC/HARBOR-UCLA MED CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
LAC/OLIVE VIEW-UCLA MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
LAC/RANCHO LOS AMIGOS NATIONAL REHABILITATION CTR	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	94
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
LAC+USC MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
LAGUNA HONDA HOSPITAL & REHABILITATION CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
LAKEWOOD REGIONAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
LODI MEMORIAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
LOMA LINDA UNIVERSITY MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
LOMA LINDA UNIVERSITY MEDICAL CENTER-MURRIETA	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	85
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	83
LOMA LINDA VA MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
LOMPOC VALLEY MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	89
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	84

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
LONG BEACH MEMORIAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
LOS ALAMITOS MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
LOS ANGELES COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	67
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	57
LOS ANGELES METROPOLITAN MEDICAL CTR	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
LOS ROBLES HOSPITAL & MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
LUCILE SALTER PACKARD CHILDREN'S HSP AT STANFORD	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
MAD RIVER COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	92
	Surgery patients who were given the right kind of antibiotic to help prevent infection	95
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
MADERA COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	94
	Surgery patients who were given the right kind of antibiotic to help prevent infection	94
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	88

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
MAMMOTH HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	88
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
MARIAN MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
MARIN GENERAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
MARINA DEL REY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
MARK TWAIN ST JOSEPH'S HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
MARSHALL MEDICAL CENTER (1-RH)	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
MAYERS MEMORIAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
MEMORIAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
MEMORIAL HOSPITAL LOS BANOS	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
MEMORIAL HOSPITAL OF GARDENA	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
MENDOCINO COAST DISTRICT HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	88
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
MENIFEE VALLEY MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	90

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
MENLO PARK SURGICAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	**NSPP
	Surgery patients who were given the right kind of antibiotic to help prevent infection	**NSPP
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	**NSPP
MERCY GENERAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
MERCY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
MERCY HOSPITAL OF FOLSOM	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
MERCY MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
MERCY MEDICAL CENTER MT SHASTA	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
MERCY MEDICAL CENTER REDDING	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
MERCY SAN JUAN MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
METHODIST HOSPITAL OF SACRAMENTO	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
METHODIST HOSPITAL OF SOUTHERN CA	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	95
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
MILLER CHILDREN'S HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
MIRACLE MILE MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
MISSION COMMUNITY HOSPITAL - PANORAMA	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	92
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
MISSION HOSPITAL REGIONAL MED CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
MONTCLAIR HOSPITAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
MONTEREY PARK HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	94
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
MOTION PICTURE & TELEVISION HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
MOUNTAINS COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
NATIVIDAD MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	90
NORTHBAY MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
NORTHERN INYO HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	76
	Surgery patients who were given the right kind of antibiotic to help prevent infection	95
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
NORTHRIDGE HOSPITAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
NOVATO COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
OAK VALLEY DISTRICT HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	89
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
O'CONNOR HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
OJAI VALLEY COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	71
OLYMPIA MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
ORANGE COAST MEMORIAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
OROVILLE HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
PACIFIC ALLIANCE MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	93
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	90
PACIFIC HOSPITAL OF LONG BEACH	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	85
PACIFICA HOSPITAL OF THE VALLEY	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	74
	Surgery patients who were given the right kind of antibiotic to help prevent infection	89
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	70

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
PALM DRIVE HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
PALMDALE REGIONAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	93
PALO ALTO VA MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
PALO VERDE HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	82
	Surgery patients who were given the right kind of antibiotic to help prevent infection	71
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	75

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
PALOMAR MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
PARADISE VALLEY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
PARKVIEW COMMUNITY HOSPITAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	93
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	85
PATIENTS' HOSPITAL OF REDDING	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
PENINSULA MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
PETALUMA VALLEY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
PIONEERS MEMORIAL HEALTHCARE DISTRICT	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	94
	Surgery patients who were given the right kind of antibiotic to help prevent infection	86
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	79
PLACENTIA LINDA HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
POMERADO HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
POMONA VALLEY HOSPITAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
PRESBYTERIAN INTERCOMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
PROVIDENCE HOLY CROSS MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
PROVIDENCE LITTLE CO OF MARY MED CTR SAN PEDRO	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
PROVIDENCE LITTLE COMPANY OF MARY MED CTR TORRANCE	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
PROVIDENCE SAINT JOSEPH MEDICAL CTR	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
PROVIDENCE TARZANA MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
QUEEN OF THE VALLEY MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
REDLANDS COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
REDWOOD MEMORIAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
REGIONAL MEDICAL CENTER OF SAN JOSE	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
RIDEOUT MEMORIAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
RIDGECREST REGIONAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	89
	Surgery patients who were given the right kind of antibiotic to help prevent infection	92
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	91
RIVERSIDE COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
RIVERSIDE COUNTY REGIONAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
RONALD REAGAN UCLA MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
SADDLEBACK MEMORIAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
SAINT AGNES MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
SAINT FRANCIS MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
SAINT FRANCIS MEMORIAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
SAINT JOHN'S HEALTH CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
SAINT LOUISE REGIONAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	93
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
SAINT VINCENT MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
SALINAS VALLEY MEMORIAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
SAN ANTONIO COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
SAN DIMAS COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
SAN FRANCISCO GENERAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
SAN FRANCISCO VA MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
SAN GABRIEL VALLEY MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	94
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	89
SAN GORGONIO MEMORIAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	80
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
SAN JOAQUIN COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
SAN JOAQUIN GENERAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	93
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94
SAN MATEO MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	94
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	87
SAN RAMON REGIONAL MEDICAL CTR	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	95
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
SANTA BARBARA COTTAGE HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
SANTA CLARA VALLEY MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	90
SANTA MONICA - UCLA MED CTR & ORTHOPAEDIC HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
SANTA ROSA MEMORIAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
SANTA YNEZ VALLEY COTTAGE HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
SCRIPPS GREEN HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
SCRIPPS MEMORIAL HOSPITAL - ENCINITAS	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
SCRIPPS MEMORIAL HOSPITAL LA JOLLA	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
SCRIPPS MERCY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
SEQUOIA HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
SETON MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
SHARP CHULA VISTA MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
SHARP CORONADO HOSPITAL AND HLTHCR CTR	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
SHARP MEMORIAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
SHASTA REGIONAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
SHERMAN OAKS HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
SIERRA NEVADA MEMORIAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
SIERRA VIEW DISTRICT HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	94
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94
SIERRA VISTA REGIONAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
SILVER LAKE MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
SIMI VALLEY HOSPITAL & HEALTH CARE SERVICES	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	91
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
SONOMA DEVELOPMENTAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
SONOMA VALLEY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
SONORA REGIONAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
SOUTHERN INYO HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
SOUTHWEST HEALTHCARE SYSTEM	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	91
ST BERNARDINE MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	93
ST ELIZABETH COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
ST HELENA HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
ST HELENA HOSPITAL-CLEARLAKE	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
ST JOHNS PLEASANT VALLEY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
ST JOHNS REGIONAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
ST JOSEPH HOSPITAL, EUREKA	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
ST JOSEPH HOSPITAL, ORANGE COUNTY	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
ST JOSEPHS MED CENTER OF STOCKTON	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
ST JUDE MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
ST MARY MEDICAL CENTER, APPLE VALLEY	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
ST MARY MEDICAL CENTER, SAN FRANCISCO	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
ST MARY'S MEDICAL CENTER, SAN FRANCISCO	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
ST ROSE HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	95
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94
STANFORD HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
STANISLAUS SURGICAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
SURPRISE VALLEY COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
SUTTER AMADOR HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
SUTTER AUBURN FAITH HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
SUTTER COAST HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
SUTTER DAVIS HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
SUTTER DELTA MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
SUTTER GENERAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
SUTTER LAKESIDE HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	93
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	91
SUTTER MATERNITY & SURGERY CENTER OF SANTA CRUZ	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
SUTTER MEDICAL CENTER OF SANTA ROSA	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
SUTTER ROSEVILLE MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
SUTTER SOLANO MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
SUTTER SURGICAL HOSPITAL - NORTH VALLEY	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
SUTTER TRACY COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
TAHOE FOREST HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	93

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
TEHACHAPI HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
TEMPLE COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
THOUSAND OAKS SURGICAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
TORRANCE MEMORIAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
TRI-CITY MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
TRI-CITY REGIONAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	87
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	88
TRINITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
TULARE REGIONAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	92
	Surgery patients who were given the right kind of antibiotic to help prevent infection	94
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
TWIN CITIES COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
UCSF MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
UKIAH VALLEY MEDICAL CENTER/HOSPITAL D	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	93
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94
UNIVERSITY OF CALIFORNIA DAVIS MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
UNIVERSITY OF CALIFORNIA IRVINE MED CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	95
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
UNIVERSITY OF CALIFORNIA SAN DIEGO MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94
USC KENNETH NORRIS JR CANCER HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
VA GREATER LOS ANGELES HEALTHCARE SYSTEM	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
VA LONG BEACH HEALTHCARE SYSTEM	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
VA N CALIFORNIA HEALTHCARE SYSTEM	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
VA SAN DIEGO HEALTHCARE SYSTEM	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
VALLEY PRESBYTERIAN HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	86

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
VALLEYCARE MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	93
VENTURA COUNTY MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
VERDUGO HILLS HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	93
	Surgery patients who were given the right kind of antibiotic to help prevent infection	95
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
VICTOR VALLEY COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	90
	Surgery patients who were given the right kind of antibiotic to help prevent infection	88
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	85

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
WASHINGTON HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
WATSONVILLE COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94
WEST ANAHEIM MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
WEST HILLS HOSPITAL & MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
WESTERN MEDICAL CENTER HOSP ANAHEIM	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94
WESTERN MEDICAL CENTER SANTA ANA	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	94
	Surgery patients who were given the right kind of antibiotic to help prevent infection	93
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	85
WHITE MEMORIAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
WHITTIER HOSPITAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	89
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)

**Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals,
April 2011 - March 2012**

Acute Care Hospitals	Care Measures	Percent Adherence
WOODLAND MEMORIAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	95
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100

*N/A (Not Available OR Too Few Surgical Procedures).

**NSPP (No Surgical Procedures Performed)