

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>ADVENTIST MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	91
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	85
<b>AHMC ANAHEIM REGIONAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	90
<b>ALAMEDA COUNTY MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	67
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>ALAMEDA HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
<b>ALHAMBRA HOSPITAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	87

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
ALTA BATES SUMMIT MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
ALTA BATES SUMMIT MEDICAL CENTER - ALTA BATES CAMP	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
ALVARADO HOSPITAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
ANAHEIM GENERAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
ANTELOPE VALLEY HOSPITAL MED CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	89

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>ARROWHEAD REGIONAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94
<b>ARROYO GRANDE COMMUNITY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	94
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
<b>BAKERSFIELD HEART HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	91
<b>BAKERSFIELD MEMORIAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
<b>BANNER LASSEN MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>BARSTOW COMMUNITY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	91
<b>BARTON MEMORIAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	90
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
<b>BEAR VALLEY COMMUNITY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	80
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	25
<b>BELLFLOWER MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	81
	Surgery patients who were given the right kind of antibiotic to help prevent infection	93
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	70
<b>BEVERLY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	94
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	93

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
BIGGS GRIDLEY MEMORIAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
BROTMAN MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	86
	Surgery patients who were given the right kind of antibiotic to help prevent infection	90
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	76
CALIFORNIA HOSPITAL MEDICAL CENTER LA	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	93
CALIFORNIA PACIFIC MEDICAL CTR - ST. LUKE'S CAMPUS	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
CALIFORNIA PACIFIC MEDICAL CTR- DAVIES CAMPUS HOSP	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	89

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>CALIFORNIA PACIFIC MEDICAL CTR- PACIFIC CAMPUS HOSP</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
<b>CATALINA ISLAND MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
<b>CEDARS-SINAI MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
<b>CENTINELA HOSPITAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>CENTRAL VALLEY GENERAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	77
	Surgery patients who were given the right kind of antibiotic to help prevent infection	79
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>CHAPMAN MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	93
	Surgery patients who were given the right kind of antibiotic to help prevent infection	82
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	81
<b>CHINESE HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	82
<b>CHINO VALLEY MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	93
	Surgery patients who were given the right kind of antibiotic to help prevent infection	93
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>CITRUS VALLEY MEDICAL CENTER-IC CAMPUS</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	91
<b>CLOVIS COMMUNITY MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
COALINGA REGIONAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
COAST PLAZA HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	78
	Surgery patients who were given the right kind of antibiotic to help prevent infection	85
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
COASTAL COMMUNITIES HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	94
	Surgery patients who were given the right kind of antibiotic to help prevent infection	94
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	93
COLLEGE HOSPITAL COSTA MESA	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
COLORADO RIVER MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>COLUSA REGIONAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	0
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>COMMUNITY AND MISSION HOSPITAL OF HUNTINGTON PARK</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	93
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	91
<b>COMMUNITY HOSPITAL OF LONG BEACH</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	83
	Surgery patients who were given the right kind of antibiotic to help prevent infection	86
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>COMMUNITY HOSPITAL OF SAN BERNARDINO</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	89
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>COMMUNITY HOSPITAL OF THE MONTEREY PENINSULA</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	85

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>COMMUNITY MEMORIAL HOSPITAL SAN BUENAVENTURA</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
<b>COMMUNITY REGIONAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	95
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
<b>CONTRA COSTA REGIONAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	81
	Surgery patients who were given the right kind of antibiotic to help prevent infection	92
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	87
<b>CORCORAN DISTRICT HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
<b>CORONA REGIONAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	94
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	93

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>DAMERON HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
<b>DELANO REGIONAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	90
	Surgery patients who were given the right kind of antibiotic to help prevent infection	90
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	89
<b>DESERT REGIONAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
<b>DESERT VALLEY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	89
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>DOCTORS HOSPITAL OF MANTECA</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>DOCTORS HOSPITAL OF WEST COVINA, INC</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	75
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	85
<b>DOCTORS MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	93
<b>DOCTORS MEDICAL CENTER-SAN PABLO</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	94
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	80
<b>DOMINICAN HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
<b>DOWNEY REGIONAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	91

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>EAST LOS ANGELES DOCTORS HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	85
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	80
<b>EAST VALLEY HOSPITAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>EASTERN PLUMAS HOSPITAL - PORTOLA CAMPUS</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
<b>EDEN MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
<b>EISENHOWER MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>EL CAMINO HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
<b>EL CENTRO REGIONAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	90
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>EMANUEL MEDICAL CENTER INC</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
<b>ENCINO HOSPITAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>ENLOE MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	94
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	93

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>FAIRCHILD MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	50
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>FALLBROOK HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>FEATHER RIVER HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
<b>FOOTHILL PRESBYTERIAN HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	90
<b>FOUNTAIN VALLEY REGIONAL HOSPITAL &amp; MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>FRANK R HOWARD MEMORIAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
<b>FRENCH HOSPITAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
<b>FRESNO HEART AND SURGICAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	95
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
<b>FRESNO SURGICAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
<b>FRESNO VA MEDICAL CENTER (VA CENTRAL CALIFORNIA HEALTHCARE SYSTEM)</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>GARDEN GROVE HOSPITAL &amp; MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>GARFIELD MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
<b>GEORGE L MEE MEMORIAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	79
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>GLENDALE ADVENTIST MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	84
<b>GLENDALE MEM HOSPITAL &amp; HLTH CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>GLENN MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
<b>GOLETA VALLEY COTTAGE HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94
<b>GOOD SAMARITAN HOSPITAL, BAKERSFIELD</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	0
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>GOOD SAMARITAN HOSPITAL, LOS ANGELES</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	89
<b>GOOD SAMARITAN HOSPITAL, SAN JOSE</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>GREATER EL MONTE COMMUNITY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	92
	Surgery patients who were given the right kind of antibiotic to help prevent infection	86
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>GROSSMONT HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
<b>HAZEL HAWKINS MEMORIAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	67
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	87
<b>HEALDSBURG DISTRICT HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	95
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	59
<b>HEMET VALLEY MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>HENRY MAYO NEWHALL MEMORIAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	94
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
<b>HI-DESERT MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	77
	Surgery patients who were given the right kind of antibiotic to help prevent infection	86
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	76
<b>HOAG MEMORIAL HOSPITAL PRESBYTERIAN</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
<b>HOAG ORTHOPEDIC INSTITUTE</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
<b>HOLLYWOOD COMMUNITY HOSPITAL OF HOLLYWOOD</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	50
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>HOLLYWOOD PRESBYTERIAN MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	88
<b>HUNTINGTON BEACH HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>HUNTINGTON MEMORIAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	93
<b>JEROLD PHELPS COMMUNITY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
<b>JOHN C FREMONT HEALTHCARE DISTRICT</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>JOHN F KENNEDY MEMORIAL HOSPITAL, INC</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	91
<b>JOHN MUIR MEDICAL CENTER - CONCORD CAMPUS</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
<b>JOHN MUIR MEDICAL CENTER - WALNUT CREEK CAMPUS</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
<b>KAISER FOUNDATION HOSP SO SACRAMENTO</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>KAISER FOUNDATION HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>KAISER FOUNDATION HOSPITAL - ANTIOCH</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
<b>KAISER FOUNDATION HOSPITAL - BALDWIN PARK</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
<b>KAISER FOUNDATION HOSPITAL - DOWNEY</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
<b>KAISER FOUNDATION HOSPITAL - FRESNO</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
<b>KAISER FOUNDATION HOSPITAL - LOS ANGELES</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>KAISER FOUNDATION HOSPITAL - PANORAMA CITY</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
<b>KAISER FOUNDATION HOSPITAL - REDWOOD CITY</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>KAISER FOUNDATION HOSPITAL - SACRAMENTO</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
<b>KAISER FOUNDATION HOSPITAL - SAN DIEGO</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>KAISER FOUNDATION HOSPITAL - SAN FRANCISCO</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>KAISER FOUNDATION HOSPITAL - SOUTH BAY</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>KAISER FOUNDATION HOSPITAL - SOUTH SAN FRANCISCO</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>KAISER FOUNDATION HOSPITAL - VACAVILLE</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	93
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>KAISER FOUNDATION HOSPITAL - WALNUT CREEK</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
<b>KAISER FOUNDATION HOSPITAL - WEST LA</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>KAISER FOUNDATION HOSPITAL - WOODLAND HILLS</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>KAISER FOUNDATION HOSPITAL ANAHEIM</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
<b>KAISER FOUNDATION HOSPITAL AND REHAB CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
<b>KAISER FOUNDATION HOSPITAL FONTANA</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94
<b>KAISER FOUNDATION HOSPITAL FREMONT/HAYWARD</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>KAISER FOUNDATION HOSPITAL MANTECA</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>KAISER FOUNDATION HOSPITAL OAKLAND/RICHMOND</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
<b>KAISER FOUNDATION HOSPITAL, RIVERSIDE</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
<b>KAISER FOUNDATION HOSPITAL- MORENO VALLEY</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
<b>KAISER FOUNDATION HOSPITAL-SAN JOSE</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
KAISER FOUNDATION HOSPITAL-SANTA CLARA	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
KAISER FOUNDATION HOSPITAL-SANTA ROSA	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
KAWEAH DELTA MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	90
	Surgery patients who were given the right kind of antibiotic to help prevent infection	91
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	87
KECK HOSPITAL OF USC	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
KERN MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	94
	Surgery patients who were given the right kind of antibiotic to help prevent infection	84
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	86

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
KERN VALLEY HEALTHCARE DISTRICT	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
LA PALMA INTERCOMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
LAC/HARBOR-UCLA MED CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	94
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
LAC/OLIVE VIEW-UCLA MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
LAC/RANCHO LOS AMIGOS NATIONAL REHABILITATION CTR	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	92
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	86

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>LAC+USC MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	91
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
<b>LAGUNA HONDA HOSPITAL &amp; REHABILITATION CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
<b>LAKWOOD REGIONAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
<b>LODI MEMORIAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	90
<b>LOMA LINDA UNIVERSITY MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>LOMA LINDA UNIVERSITY MEDICAL CENTER-MURRIETA</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
<b>LOMA LINDA VA MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
<b>LOMPOC VALLEY MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	91
	Surgery patients who were given the right kind of antibiotic to help prevent infection	86
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	74
<b>LONG BEACH MEMORIAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	89
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94
<b>LOS ALAMITOS MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
LOS ANGELES COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	82
	Surgery patients who were given the right kind of antibiotic to help prevent infection	91
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	45
LOS ANGELES METROPOLITAN MEDICAL CTR	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	60
	Surgery patients who were given the right kind of antibiotic to help prevent infection	80
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	60
LOS ROBLES HOSPITAL & MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
MAD RIVER COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	95
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	85
MADERA COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	92
	Surgery patients who were given the right kind of antibiotic to help prevent infection	89
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	93

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>MARIAN MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
<b>MARIN GENERAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
<b>MARINA DEL REY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	91
<b>MARK TWAIN ST JOSEPH'S HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>MARSHALL MEDICAL CENTER (1-RH)</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>MAYERS MEMORIAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
<b>MEMORIAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
<b>MEMORIAL HOSPITAL LOS BANOS</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
<b>MEMORIAL HOSPITAL OF GARDENA</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	93
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	93
<b>MENDOCINO COAST DISTRICT HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	88
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>MENIFEE VALLEY MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
<b>MENLO PARK SURGICAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
<b>MERCY GENERAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
<b>MERCY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
<b>MERCY HOSPITAL OF FOLSOM</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>MERCY MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94
<b>MERCY MEDICAL CENTER MT SHASTA</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>MERCY MEDICAL CENTER REDDING</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
<b>MERCY SAN JUAN MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
<b>METHODIST HOSPITAL OF SACRAMENTO</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>METHODIST HOSPITAL OF SOUTHERN CA</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	93
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
<b>MIRACLE MILE MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
<b>MISSION COMMUNITY HOSPITAL - PANORAMA</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	91
	Surgery patients who were given the right kind of antibiotic to help prevent infection	94
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	91
<b>MISSION HOSPITAL REGIONAL MED CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
<b>MONTCLAIR HOSPITAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	88
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	93

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>MONTEREY PARK HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
<b>MOTION PICTURE &amp; TELEVISION HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>MOUNTAINS COMMUNITY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
<b>NATIVIDAD MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	91
<b>NORTHBAY MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>NORTHERN INYO HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	62
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>NORTHRIDGE HOSPITAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
<b>NOVATO COMMUNITY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
<b>OAK VALLEY DISTRICT HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	86
	Surgery patients who were given the right kind of antibiotic to help prevent infection	93
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	89
<b>O'CONNOR HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
OJAI VALLEY COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	80
OLYMPIA MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
ORANGE COAST MEMORIAL MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	93
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	93
OROVILLE HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	94
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
PACIFIC ALLIANCE MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	94
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>PACIFIC HOSPITAL OF LONG BEACH</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	87
<b>PACIFICA HOSPITAL OF THE VALLEY</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	73
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	71
<b>PALM DRIVE HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	92
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
<b>PALMDALE REGIONAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	95
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	88
<b>PALO ALTO VA MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>PALO VERDE HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	64
	Surgery patients who were given the right kind of antibiotic to help prevent infection	50
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	91
<b>PALOMAR MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
<b>PARADISE VALLEY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
<b>PARKVIEW COMMUNITY HOSPITAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	87
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
<b>PATIENTS' HOSPITAL OF REDDING</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	94
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>PENINSULA MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
<b>PETALUMA VALLEY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
<b>PIONEERS MEMORIAL HEALTHCARE DISTRICT</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	89
	Surgery patients who were given the right kind of antibiotic to help prevent infection	84
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	78
<b>PLACENTIA LINDA HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
<b>PLUMAS DISTRICT HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>POMERADO HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
<b>POMONA VALLEY HOSPITAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	93
<b>PRESBYTERIAN INTERCOMMUNITY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
<b>PROVIDENCE HOLY CROSS MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
<b>PROVIDENCE LITTLE CO OF MARY MED CTR SAN PEDRO</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>PROVIDENCE LITTLE COMPANY OF MARY MED CTR TORRANCE</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
<b>PROVIDENCE SAINT JOSEPH MEDICAL CTR</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
<b>PROVIDENCE TARZANA MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
<b>QUEEN OF THE VALLEY MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
<b>REDLANDS COMMUNITY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>REDWOOD MEMORIAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
<b>REGIONAL MEDICAL CENTER OF SAN JOSE</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
<b>RIDEOUT MEMORIAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
<b>RIDGECREST REGIONAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	82
	Surgery patients who were given the right kind of antibiotic to help prevent infection	59
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	75
<b>RIVERSIDE COMMUNITY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>RIVERSIDE COUNTY REGIONAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94
<b>RONALD REAGAN UCLA MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	91
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	91
<b>SADDLEBACK MEMORIAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
<b>SAINT AGNES MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
<b>SAINT FRANCIS MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	89

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
SAINT FRANCIS MEMORIAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
SAINT JOHN'S HEALTH CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
SAINT LOUISE REGIONAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
SAINT VINCENT MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
SALINAS VALLEY MEMORIAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>SAN ANTONIO COMMUNITY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
<b>SAN DIMAS COMMUNITY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
<b>SAN FRANCISCO GENERAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	90
<b>SAN FRANCISCO VA MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
<b>SAN GABRIEL VALLEY MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	85

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>SAN GORGONIO MEMORIAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	85
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>SAN JOAQUIN COMMUNITY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	93
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
<b>SAN JOAQUIN GENERAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	91
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
<b>SAN MATEO MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	95
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	82
<b>SAN RAMON REGIONAL MEDICAL CTR</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	95
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	91

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>SANTA BARBARA COTTAGE HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	94
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	91
<b>SANTA CLARA VALLEY MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	85
<b>SANTA MONICA - UCLA MED CTR &amp; ORTHOPAEDIC HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
<b>SANTA ROSA MEMORIAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	92
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
<b>SANTA YNEZ VALLEY COTTAGE HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>SCRIPPS GREEN HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
<b>SCRIPPS MEMORIAL HOSPITAL - ENCINITAS</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>SCRIPPS MEMORIAL HOSPITAL LA JOLLA</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
<b>SCRIPPS MERCY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
<b>SEQUOIA HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>SETON MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94
<b>SHARP CHULA VISTA MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
<b>SHARP CORONADO HOSPITAL AND HLTHCR CTR</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
<b>SHARP MEMORIAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
<b>SHASTA REGIONAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>SHERMAN OAKS HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>SIERRA KINGS DISTRICT HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	75
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>SIERRA NEVADA MEMORIAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
<b>SIERRA VIEW DISTRICT HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	94
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	91
<b>SIERRA VISTA REGIONAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>SILVER LAKE MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	40
	Surgery patients who were given the right kind of antibiotic to help prevent infection	80
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	80
<b>SIMI VALLEY HOSPITAL &amp; HEALTH CARE SERVICES</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	87
<b>SONOMA DEVELOPMENTAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
<b>SONOMA VALLEY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
<b>SONORA REGIONAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>SOUTHERN INYO HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
<b>SOUTHWEST HEALTHCARE SYSTEM</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	94
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	80
<b>ST BERNARDINE MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	93
<b>ST ELIZABETH COMMUNITY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
<b>ST HELENA HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>ST HELENA HOSPITAL-CLEARLAKE</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
<b>ST JOHNS PLEASANT VALLEY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
<b>ST JOHNS REGIONAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	91
<b>ST JOSEPH HOSPITAL, EUREKA</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
<b>ST JOSEPH HOSPITAL, Orange County</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>ST JOSEPHS MED CENTER OF STOCKTON</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
<b>ST JUDE MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
<b>ST MARY MEDICAL CENTER, APPLE VALLEY</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	94
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	88
<b>ST MARY MEDICAL CENTER, SAN FRANCISCO</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
<b>ST MARY'S MEDICAL CENTER, SAN FRANCISCO</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>ST ROSE HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	92
	Surgery patients who were given the right kind of antibiotic to help prevent infection	93
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
<b>STANFORD HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94
<b>STANISLAUS SURGICAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
<b>SURPRISE VALLEY COMMUNITY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
<b>SUTTER AMADOR HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
SUTTER AUBURN FAITH HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
SUTTER COAST HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	91
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
SUTTER DAVIS HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
SUTTER DELTA MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
SUTTER GENERAL HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>SUTTER LAKESIDE HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	94
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
<b>SUTTER MATERNITY &amp; SURGERY CENTER OF SANTA CRUZ</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
<b>SUTTER MEDICAL CENTER OF SANTA ROSA</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
<b>SUTTER ROSEVILLE MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
<b>SUTTER SOLANO MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>SUTTER SURGICAL HOSPITAL - NORTH VALLEY</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	94
	Surgery patients who were given the right kind of antibiotic to help prevent infection	95
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
<b>SUTTER TRACY COMMUNITY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
<b>TAHOE FOREST HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	91
	Surgery patients who were given the right kind of antibiotic to help prevent infection	89
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	82
<b>TEHACHAPI HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
<b>TEMPLE COMMUNITY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	33

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>TORRANCE MEMORIAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	94
<b>TRI-CITY MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
<b>TRI-CITY REGIONAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	90
	Surgery patients who were given the right kind of antibiotic to help prevent infection	90
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	84
<b>TRINITY HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
<b>TULARE REGIONAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	87

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
TWIN CITIES COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
UCSF MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	96
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
UKIAH VALLEY MEDICAL CENTER/HOSPITAL D	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	93
	Surgery patients who were given the right kind of antibiotic to help prevent infection	93
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
UNIVERSITY OF CALIFORNIA DAVIS MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
UNIVERSITY OF CALIFORNIA IRVINE MED CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	94
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>UNIVERSITY OF CALIFORNIA SAN DIEGO MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	96
<b>USC KENNETH NORRIS JR CANCER HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	*N/A
	Surgery patients who were given the right kind of antibiotic to help prevent infection	*N/A
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	*N/A
<b>VA GREATER LOS ANGELES HEALTHCARE SYSTEM</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	98
	Surgery patients who were given the right kind of antibiotic to help prevent infection	89
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	97
<b>VA LONG BEACH HEALTHCARE SYSTEM</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
<b>VA N CALIFORNIA HEALTHCARE SYSTEM</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	94
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
VA SAN DIEGO HEALTHCARE SYSTEM	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
VALLEY PRESBYTERIAN HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	93
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	84
VALLEYCARE MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
VENTURA COUNTY MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	98
VERDUGO HILLS HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	96
	Surgery patients who were given the right kind of antibiotic to help prevent infection	92
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	91

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
VICTOR VALLEY COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	83
	Surgery patients who were given the right kind of antibiotic to help prevent infection	69
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	67
WASHINGTON HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	99
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
WATSONVILLE COMMUNITY HOSPITAL	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99
WEST ANAHEIM MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	100
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100
WEST HILLS HOSPITAL & MEDICAL CENTER	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	100
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	99

\*No surgical procedures performed.

## Surgical Care Improvement Project (SCIP) Antimicrobial Prophylaxis Measures in California Acute Care Hospitals, April 2010 - March 2011

Acute Care Hospitals	Care Measures	Percent Adherence
<b>WESTERN MEDICAL CENTER HOSP ANAHEIM</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	90
<b>WESTERN MEDICAL CENTER SANTA ANA</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	95
	Surgery patients who were given the right kind of antibiotic to help prevent infection	93
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	91
<b>WHITE MEMORIAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	99
	Surgery patients who were given the right kind of antibiotic to help prevent infection	98
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	95
<b>WHITTIER HOSPITAL MEDICAL CENTER</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	94
	Surgery patients who were given the right kind of antibiotic to help prevent infection	90
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	92
<b>WOODLAND MEMORIAL HOSPITAL</b>	Surgery patients who were given an antibiotic at the right time (within one hour before surgery) to help prevent infection	97
	Surgery patients who were given the right kind of antibiotic to help prevent infection	97
	Surgery patients whose preventive antibiotics were stopped at the right time (within 24 hours after surgery)	100

\*No surgical procedures performed.