

Table 27. Rates of central line-associated bloodstream infection in patient care locations providing general pediatric rehabilitation care, reported by California hospitals, April 2010 - March 2011 (inclusive)

Reporting hospital*	Cases	Central line-days	Patient days	Device utilization ratio	Rate	95% Confidence Interval	Statistical interpretation**	
							CA	U.S.
STATE OF CALIFORNIA POOLED DATA	7	2674	15129	0.18	2.6			
CHILDREN'S HOSP CENTRAL CA, MADERA	6	1696	4534	0.37	3.5	(1.3, 7.7)	*	*
RADY CHILDREN'S HOSP, SAN DIEGO	1	978	10595	0.09	1.0	(0.0, 5.7)	*	*

* Hospitals with only one acute care campus associated with their license were listed by the campus name; hospitals with more than one campus associated with their license were listed by licensee names with acute care campus name(s) indented below

**Statistical interpretation indicates, compared with California (CA) and U.S. average rates, hospital-specific rates that were significantly (statistically) higher (●), lower (○), no different (⊙), not tested due to unstable rates (—) or not tested due to too few hospitals reporting (*)

† Hospital reported infection data aggregated over all acute care campuses

‡ Hospital reported infection data separately by acute care campus

Notes: Rate per 1000 central line days; device utilization ratio is central line days/patient days for months when central lines were used; 95% confidence interval calculated for rates based on exact Poisson distribution; rehabilitation general care locations provide care to patients who have lost function due to acute or chronic pain, musculoskeletal problems, stroke, or catastrophic events resulting in complete or partial paralysis

Source: Central Line-Associated Bloodstream Infections in California Hospitals, April 2010 through March 2011. California Department of Public Health