

Table 35. Rates of temporary central line-associated bloodstream infection in patient care locations providing pediatric bone marrow transplant care, reported by California hospitals, April 2010 - March 2011 (inclusive)

Reporting hospital*	Cases	Central line-days	Patient days	Device utilization ratio	Rate	95% Confidence Interval	Statistical interpretation**	
							CA	U.S.
<b>STATE OF CALIFORNIA POOLED DATA</b>	<b>2</b>	<b>3734</b>	<b>6859</b>	<b>0.54</b>	<b>0.5</b>			
CHILDRENS HOSP AND RESEARCH CTR AT OAKLAND	0	31	164	0.19	-----	-----	*	*
CHILDRENS HOSP OF LOS ANGELES	1	3208	3757	0.85	0.3	(0.0, 1.7)	*	*
CHILDREN'S HOSP OF ORANGE COUNTY, ORANGE	1	495	2938	0.17	2.0	(0.1, 11.3)	*	*

\* Hospitals with only one acute care campus associated with their license were listed by the campus name; hospitals with more than one campus associated with their license were listed by licensee names with acute care campus name(s) indented below

\*\*Statistical interpretation indicates, compared with California (CA) and U.S. average rates, hospital-specific rates that were significantly (statistically) higher (●), lower (○), no different (⊙), not tested due to unstable rates (—) or not tested due to too few hospitals reporting (\*)

† Hospital reported infection data aggregated over all acute care campuses

‡ Hospital reported infection data separately by acute care campus

Notes: Rate per 1000 central line days; device utilization ratio is central line days/patient days for months when central lines were used; 95% confidence interval calculated for rates based on exact Poisson distribution; ----- indicates data not shown due to unstable rates; bone marrow transplant special care locations specialize in patients who undergo bone marrow (stem cell) transplant for the treatment of various disorders

Source: Central Line-Associated Bloodstream Infections in California Hospitals, April 2010 through March 2011. California Department of Public Health