

2007 KINDERGARTEN RETROSPECTIVE SURVEY RESULTS  
CALIFORNIA DEPARTMENT OF PUBLIC HEALTH, IMMUNIZATION BRANCH

### Introduction

The annual Kindergarten Retrospective Survey (KRS) is a primary source of information about childhood immunization coverage in California. This survey provides estimates of immunization coverage among kindergarten students at various age checkpoints. This report presents results from this year's Kindergarten Retrospective Survey and the attached tables and figures present comparisons of immunization coverage by a variety of factors, including region and race/ethnicity.

### Methods and Participants

This year's Kindergarten Retrospective Survey was conducted in a sample of approximately 3% of California's kindergartens in concurrence with selective review. Local health departments visited 248 schools with kindergartens and collected copies of every sixth student immunization record (blue card). Demographic information and immunization history were extracted from each record. Data were analyzed using birth dates and immunization dates to retrospectively estimate immunization coverage at various age checkpoints. Age checkpoints are defined according to whether or not children are up-to-date for DTaP, Polio, MMR, Hep B, and Varicella vaccines at 3, 5, 7, 13, 19 and 24 months.

This year's sample consists of 2,438 kindergarten students. Children in this sample were born between 2000 and 2002. Since this is a retrospective survey, estimates of immunization coverage represent levels among toddlers approximately 3-4 years ago.

### Results and Discussion

Results from this year's Kindergarten Retrospective Survey indicate that the percent of California's kindergarten students at 24 months of age with the 4:3:1 series (4+ DTP, 3+ Polio, and 1+ MMR) was 75.7±2.1%. Immunization coverage for individual vaccines ranged from 78.8±2.2% for 4+ DTaP to 92.4±1.3% for 1+ MMR. Table 1 shows a comparison of this year's Kindergarten Retrospective Survey and the 2003 National Immunization Survey. Estimates of immunization coverage for individual vaccines and the 4:3:1 series were similar in both surveys.

Immunization coverage among California's children is similar across regions (Table 2 and Figure A). This year, the percent of kindergarten students at 24 months of age with the 4:3:1 vaccine series was 74.3±3.5% among children in Los Angeles, 78.9±3.7% among children in Southern California, 80.3±5.9% among children in the Bay Area, 71.7±11.7% among children in the Central Coast, 73.1±8.5% among children in the Central Valley, 65.8±10.5% among children in the North Central Valley, and 74.7±4.9% among children in rural Northern California. Although the point estimates may seem different, once confidence intervals are factored in, it becomes clear that any observed differences are not statistically significant.

Table 3 shows immunization coverage among California's children by race/ethnicity. This year, the percent of kindergarten students at 24 months of age with the 4:3:1 vaccine series was 76.6±4.2% among white children, 75.0±2.4% among Hispanic children, 72.0±8.4% among Black children, and 81.8±6.5% among Asian children. The Kindergarten Retrospective Survey is one of the only sources of information on subpopulations of Black and Asian children and data suggest a serious and persistent disparity in immunization coverage among Black children. Over the past 5 years, coverage with the 4:3:1 vaccine series among Black children has been significantly lower than White, Hispanic, and Asian children (Figure B) and this is consistent at all age checkpoints (Figure C).

#### Description of Attached Tables

Enclosed are tables and figures showing this year's results. Table 1 compares immunization coverage as measured by the Kindergarten Retrospective Survey and the National Immunization Survey. Table 2 shows immunization coverage among kindergarten students at 24 months of age by region. Figure A is a map of 4:3:1 vaccine series coverage by region. Table 3 shows immunization coverage among kindergarten students at 24 months of age by race/ethnicity. Figures B&C show disparities in 4:3:1 vaccine series coverage among children by race/ethnicity over time (Figure B) and at all age checkpoints (Figure C). Tables 4-8 show immunization coverage among kindergarten students at 3, 5, 7, 13 and 19 months of age by race/ethnicity.

*For further information, please contact Madina Sharifi at (510) 620-3757 or [Madina.Sharifi@cdph.ca.gov](mailto:Madina.Sharifi@cdph.ca.gov) at the California Department of Public Health, Immunization Branch.*

**KINDERGARTEN RETROSPECTIVE SURVEY**  
CALIFORNIA, 2007

TABLE 1: COMPARISON OF IMMUNIZATION COVERAGE AS MEASURED BY THE KINDERGATEN RETROSPECTIVE SURVEY (KRS) AND THE NATIONAL IMMUNIZATION SURVEY (NIS)<sup>1</sup>

	n	DTaP (4+) <sup>2</sup>	Polio (3+) <sup>3</sup>	MMR (1+) <sup>4</sup>	Hep B (3+) <sup>5</sup>	Var (1+) <sup>6</sup>	4:3:1 <sup>7</sup>
KRS (24 mo.)	2,438	78.8% ± 2.2%	90.5% ± 1.6%	92.4% ± 1.3%	89.8% ± 1.4%	84.8% ± 1.9%	75.7% ± 2.1%
NIS (24 mo.)	n/a	79.3% ± 4.3%	89.3% ± 3.7%	90.1% ± 3.4%	89.1% ± 3.6%	87.4% ± 3.6%	75.7% ± 4.6%

1. Data from the 2003 NIS are used for comparison because data from the KRS are retrospective; Children in both surveys were born between 2000 and 2002
2. Four or more doses of any diphtheria and tetanus toxoids and pertussis vaccines including diphtheria and tetanus toxoids, and any acellular pertussis vaccine (DTP/DTaP/DT)
3. Three or more doses of any poliovirus vaccine
4. One or more doses of measles-mumps-rubella vaccine
5. Three or more doses of hepatitis B vaccine
6. One or more doses of varicella vaccine, unadjusted for history of varicella disease
7. Four or more doses of DTaP, three or more doses of Polio, and one or more doses of MMR

**KINDERGARTEN RETROSPECTIVE SURVEY  
CALIFORNIA, 2007**

TABLE 2: IMMUNIZATION COVERAGE AMONG KINDERGARTEN STUDENTS AT 24 MONTHS OF AGE BY REGION<sup>1</sup>

	n	DTaP (4+) <sup>2</sup>	Polio (3+) <sup>3</sup>	MMR (1+) <sup>4</sup>	Hep B (3+) <sup>5</sup>	Var (1+) <sup>6</sup>	4:3:1 <sup>7</sup>	4:3:1:-:3 <sup>8</sup>	4:3:1:-:3:1 <sup>9</sup>
ALL	2,438	78.8% ± 2.2%	90.5% ± 1.6%	92.4% ± 1.3%	89.8% ± 1.4%	84.8% ± 1.9%	75.7% ± 2.1%	73.2% ± 2.2%	69.6% ± 2.3%
Los Angeles	767	78.8% ± 3.7%	91.7% ± 2.9%	93.5% ± 2.5%	88.2% ± 3.1%	84.4% ± 3.0%	74.3% ± 3.5%	71.0% ± 3.5%	67.0% ± 3.8%
Other So. California	612	81.0% ± 4.1%	91.2% ± 3.0%	93.5% ± 1.8%	90.2% ± 2.1%	85.4% ± 3.6%	78.9% ± 3.7%	75.7% ± 4.2%	72.0% ± 4.3%
SF Bay Area	466	82.3% ± 5.6%	91.3% ± 4.2%	92.3% ± 4.0%	91.6% ± 4.2%	87.0% ± 4.9%	80.3% ± 5.9%	79.1% ± 5.9%	75.2% ± 6.4%
Central Coast	93	76.0% ± 12.5%	84.4% ± 8.8%	91.0% ± 8.0%	86.1% ± 9.6%	87.9% ± 10.2%	71.7% ± 11.7%	70.2% ± 12.4%	69.7% ± 12.3%
Central Valley	190	75.6% ± 9.8%	88.6% ± 6.3%	94.3% ± 3.0%	92.6% ± 3.0%	86.9% ± 6.7%	73.1% ± 8.5%	71.8% ± 8.2%	68.3% ± 9.9%
North Central Valley	199	71.6% ± 10.3%	84.7% ± 5.2%	82.7% ± 5.5%	88.2% ± 6.7%	76.1% ± 8.5%	65.8% ± 10.5%	64.9% ± 10.5%	61.0% ± 10.3%
Rural No. California	111	76.2% ± 4.7%	96.7% ± 4.0%	93.0% ± 6.4%	92.2% ± 5.5%	83.4% ± 4.9%	74.7% ± 4.9%	72.5% ± 5.1%	69.0% ± 6.4%

1. Regions do not exactly match Field Rep assigned regions:

Los Angeles includes Los Angeles County

Other So. California includes Imperial, Orange, Riverside, San Bernardino, and San Diego Counties

SF Bay Area includes Alameda, Contra Costa, Marin, Napa, San Francisco, San Mateo, Santa Clara, Solano, and Sonoma Counties

Central Coast includes Monterey, San Luis Obispo, Santa Barbara, Santa Cruz, and Ventura Counties

Central Valley includes Fresno, Kern, Kings, Madera, Merced, and Tulare Counties

North Central Valley includes Sacramento, San Joaquin, and Stanislaus Counties

Rural No. California includes Alpine, Amador, Butte, Calaveras, Colusa, Del Norte, El Dorado, Glenn, Humboldt, Inyo, Lake, Lassen, Mariposa, Mendocino, Modoc, Mono, Nevada, Placer, Plumas, San Benito, Shasta, Sierra, Siskiyou, Sutter, Tehama, Trinity, Tuolumne, Yolo, and Yuba Counties

2. Four or more doses of any diphtheria and tetanus toxoids and pertussis vaccines including diphtheria and tetanus toxoids, and any acellular pertussis vaccine (DTP/DTaP/DT)

3. Three or more doses of any poliovirus vaccine

4. One or more doses of measles-mumps-rubella vaccine

5. Three or more doses of hepatitis B vaccine

6. One or more doses of varicella vaccine, unadjusted for history of varicella disease


7. Four or more doses of DTaP, three or more doses of Polio, and one or more doses of MMR

8. Four or more doses of DTaP, three or more doses of Polio, one or more doses of MMR, and three or more doses of Hep B

9. Four or more doses of DTaP, three or more doses of Polio, one or more doses of MMR, three or more doses of Hep B, and one or more doses of varicella

KINDERGARTEN RETROSPECTIVE SURVEY  
CALIFORNIA, 2007

FIGURE A: 4:3:1<sup>1</sup> COVERAGE AMONG KINDERGARTEN STUDENTS AT 24 MONTHS OF AGE BY REGION<sup>2</sup>



Central Valley includes Fresno, Kern, Kings, Madera, Merced, and Tulare Counties

North Central Valley includes Sacramento, San Joaquin, and Stanislaus Counties

Rural No. California includes Alpine, Amador, Butte, Calaveras, Colusa, Del Norte, El Dorado, Glenn, Humboldt, Inyo, Lake, Lassen, Mariposa, Mendocino, Modoc, Mono, Nevada, Placer, Plumas, San Benito, Shasta, Sierra, Siskiyou, Sutter, Tehama, Trinity, Tuolumne, Yolo, and Yuba Counties

**KINDERGARTEN RETROSPECTIVE SURVEY**  
CALIFORNIA, 2007


TABLE 3: IMMUNIZATION COVERAGE AMONG KINDERGARTEN STUDENTS AT 24 MONTHS OF AGE BY RACE/ETHNICITY<sup>1</sup>

	n	DTaP (4+) <sup>2</sup>	Polio (3+) <sup>3</sup>	MMR (1+) <sup>4</sup>	Hep B (3+) <sup>5</sup>	Var (1+) <sup>6</sup>	4:3:1 <sup>7</sup>	4:3:1:-:3 <sup>8</sup>	4:3:1:-:3:1 <sup>9</sup>
ALL	2,438	78.8% ± 2.2%	90.5% ± 1.6%	92.4% ± 1.3%	89.8% ± 1.4%	84.8% ± 1.9%	75.7% ± 2.1%	73.2% ± 2.2%	69.6% ± 2.3%
White	541	80.7% ± 4.5%	88.3% ± 3.0%	91.9% ± 2.6%	90.5% ± 2.4%	86.5% ± 3.7%	76.6% ± 4.2%	74.6% ± 4.2%	70.5% ± 4.8%
Hispanic	1,080	78.1% ± 2.5%	92.6% ± 1.9%	93.3% ± 1.6%	91.2% ± 1.5%	84.4% ± 2.2%	75.0% ± 2.4%	72.7% ± 2.5%	69.2% ± 2.7%
Black	346	73.7% ± 8.5%	86.1% ± 7.7%	88.9% ± 5.9%	85.8% ± 7.0%	82.0% ± 7.4%	72.0% ± 8.4%	70.3% ± 8.5%	66.3% ± 8.9%
Asian	223	85.9% ± 5.9%	92.0% ± 4.6%	93.5% ± 3.7%	89.2% ± 5.3%	86.3% ± 5.3%	81.8% ± 6.5%	78.4% ± 7.0%	74.5% ± 7.2%
Other	106	77.3% ± 11.0%	89.6% ± 6.7%	91.0% ± 7.4%	85.6% ± 9.0%	83.7% ± 9.6%	76.2% ± 11.4%	73.7% ± 11.7%	70.1% ± 12.5%
Unknown	142	69.8% ± 11.9%	85.2% ± 9.3%	90.3% ± 8.1%	81.5% ± 8.7%	82.3% ± 9.9%	69.8% ± 11.9%	64.3% ± 12.2%	63.2% ± 12.2%

1. As recorded on student immunization record (blue card). Individual racial groups do not include Hispanic children. Children of Hispanic ethnicity may be of any race
2. Four or more doses of any diphtheria and tetanus toxoids and pertussis vaccines including diphtheria and tetanus toxoids, and any acellular pertussis vaccine (DTP/DTaP/DT)
3. Three or more doses of any poliovirus vaccine
4. One or more doses of measles-mumps-rubella vaccine
5. Three or more doses of hepatitis B vaccine
6. One or more doses of varicella vaccine, unadjusted for history of varicella disease
7. Four or more doses of DTaP, three or more doses of Polio, and one or more doses of MMR
8. Four or more doses of DTaP, three or more doses of Polio, one or more doses of MMR, and three or more doses of Hep B
9. Four or more doses of DTaP, three or more doses of Polio, one or more doses of MMR, three or more doses of Hep B, and one or more doses of varicella

**KINDERGARTEN RETROSPECTIVE SURVEY  
CALIFORNIA, 2007**

FIGURE B: 4:3:1<sup>1</sup> COVERAGE AMONG KINDERGARTEN STUDENTS AT 24 MONTHS OF AGE BY YEAR OF KRS<sup>2</sup> AND RACE/ETHNICITY<sup>3</sup>




1. Four or more doses of DTaP, three or more doses of polio, and one or more doses of MMR

2. Year of KRS: year collected is the school year that students were in kindergarten. Since this is a retrospective survey, year estimated reflects the year when kindergarten students were 24 months

3. As recorded on student immunization record (blue card). Individual racial groups do not include Hispanic children. Children of Hispanic ethnicity may be of any race

**KINDERGARTEN RETROSPECTIVE SURVEY**  
CALIFORNIA, 2007

FIGURE C: 4:3:1<sup>1</sup> COVERAGE AMONG KINDERGARTEN STUDENTS BY AGE CHECKPOINT<sup>2</sup> AND RACE/ETHNICITY<sup>3</sup>



1. Age appropriate doses of DTaP, polio, and MMR

2. Age checkpoints are defined according to whether or not children are up-to-date for age appropriate doses of DTaP, polio, and MMR at 3, 5, 7, 13, 19 and 24 months

3. As recorded on student immunization record (blue card). Individual racial groups do not include Hispanic children. Children of Hispanic ethnicity may be of any race

**KINDERGARTEN RETROSPECTIVE SURVEY**  
CALIFORNIA, 2007

TABLE 4: IMMUNIZATION COVERAGE AMONG KINDERGARTEN STUDENTS AT 3 MONTHS OF AGE BY RACE/ETHNICITY<sup>1</sup>

	n	DTaP (1+) <sup>2</sup>	Polio (1+) <sup>3</sup>	Hep B (1+) <sup>4</sup>	1:1:0 <sup>5</sup>	1:1:0:-:1 <sup>6</sup>
ALL	2,438	85.0% ± 2.0%	84.3% ± 2.0%	85.9% ± 1.9%	82.5% ± 2.1%	78.1% ± 2.2%
White	541	88.0% ± 3.2%	84.9% ± 3.8%	84.5% ± 3.9%	84.2% ± 4.0%	77.1% ± 4.5%
Hispanic	1,080	84.1% ± 2.6%	84.0% ± 2.7%	87.4% ± 2.2%	81.9% ± 2.8%	79.3% ± 2.7%
Black	346	76.8% ± 9.9%	78.1% ± 8.4%	80.9% ± 7.9%	75.9% ± 9.7%	71.6% ± 9.9%
Asian	223	86.8% ± 5.3%	89.8% ± 5.1%	88.7% ± 4.6%	86.2% ± 5.6%	82.4% ± 5.8%
Other	106	88.2% ± 7.9%	86.4% ± 8.7%	87.6% ± 7.6%	86.4% ± 8.7%	80.1% ± 9.4%
Unknown	142	83.4% ± 8.1%	79.3% ± 10.4%	77.3% ± 10.0%	78.4% ± 10.2%	70.7% ± 10.6%

1. As recorded on student immunization record (blue card). Individual racial groups do not include Hispanic children. Children of Hispanic ethnicity may be of any race
2. One or more doses of any diphtheria and tetanus toxoids and pertussis vaccines including diphtheria and tetanus toxoids, and any pertussis vaccine (DTP/DTaP/DT)
3. One or more doses of any poliovirus vaccine
4. One or more doses of hepatitis B vaccine
5. One or more doses of DTaP and one or more doses of Polio
6. One or more doses of DTaP, one or more doses of polio, and one or more doses of Hep B

**KINDERGARTEN RETROSPECTIVE SURVEY**  
CALIFORNIA, 2007

TABLE 5: IMMUNIZATION COVERAGE AMONG KINDERGARTEN STUDENTS AT 5 MONTHS OF AGE BY RACE/ETHNICITY<sup>1</sup>

	n	DTaP (2+) <sup>2</sup>	Polio (2+) <sup>3</sup>	Hep B (2+) <sup>4</sup>	2:2:0 <sup>5</sup>	2:2:0:-:2 <sup>6</sup>
ALL	2,438	73.4% ± 2.6%	72.5% ± 2.5%	74.8% ± 2.5%	70.3% ± 2.6%	63.6% ± 2.7%
White	541	79.0% ± 4.9%	75.5% ± 5.0%	74.1% ± 4.8%	74.0% ± 5.2%	63.9% ± 5.4%
Hispanic	1,080	71.1% ± 3.2%	71.6% ± 3.2%	76.2% ± 3.0%	68.8% ± 3.4%	64.0% ± 3.4%
Black	346	53.3% ± 9.4%	53.4% ± 9.4%	62.6% ± 8.1%	51.4% ± 9.3%	45.2% ± 9.3%
Asian	223	84.9% ± 6.0%	85.2% ± 5.8%	84.1% ± 5.2%	83.3% ± 6.3%	78.0% ± 6.4%
Other	106	78.1% ± 10.2%	77.8% ± 10.3%	73.6% ± 9.2%	77.8% ± 10.3%	67.9% ± 9.0%
Unknown	142	67.9% ± 11.2%	62.2% ± 12.0%	63.9% ± 12.9%	58.8% ± 12.3%	50.4% ± 13.6%

1. As recorded on student immunization record (blue card). Individual racial groups do not include Hispanic children. Children of Hispanic ethnicity may be of any race
2. Two or more doses of any diphtheria and tetanus toxoids and pertussis vaccines including diphtheria and tetanus toxoids, and any pertussis vaccine (DTP/DTaP/DT)
3. Two or more doses of any poliovirus vaccine
4. Two or more doses of hepatitis B vaccine
5. Two or more doses of DTaP and two or more doses of Polio
6. Two or more doses of DTaP, two or more doses of polio, and two or more doses of Hep B

**KINDERGARTEN RETROSPECTIVE SURVEY**  
CALIFORNIA, 2007

TABLE 6: IMMUNIZATION COVERAGE AMONG KINDERGARTEN STUDENTS AT 7 MONTHS OF AGE BY RACE/ETHNICITY<sup>1</sup>

	n	DTaP (3+) <sup>2</sup>	Polio (2+) <sup>3</sup>	Hep B (2+) <sup>4</sup>	3:2:0 <sup>5</sup>	3:2:0:-:2 <sup>6</sup>
ALL	2,438	61.1% ± 3.1%	88.5% ± 1.6%	86.6% ± 1.6%	60.2% ± 3.0%	57.1% ± 3.0%
White	541	66.3% ± 6.2%	89.0% ± 2.8%	83.7% ± 3.4%	64.4% ± 6.0%	59.2% ± 5.7%
Hispanic	1,080	57.1% ± 3.4%	88.7% ± 2.4%	89.5% ± 1.9%	56.4% ± 3.4%	54.7% ± 3.4%
Black	346	39.7% ± 8.8%	80.8% ± 8.2%	80.2% ± 6.9%	39.7% ± 8.8%	37.7% ± 8.3%
Asian	223	77.1% ± 7.7%	93.0% ± 4.2%	88.7% ± 5.0%	76.5% ± 7.8%	73.2% ± 7.7%
Other	106	71.5% ± 11.4%	89.5% ± 8.4%	86.9% ± 9.3%	71.3% ± 11.5%	69.3% ± 11.9%
Unknown	142	62.0% ± 11.2%	84.2% ± 8.7%	75.7% ± 10.9%	60.9% ± 11.2%	54.8% ± 12.4%

1. As recorded on student immunization record (blue card). Individual racial groups do not include Hispanic children. Children of Hispanic ethnicity may be of any race
2. Three or more doses of any diphtheria and tetanus toxoids and pertussis vaccines including diphtheria and tetanus toxoids, and any pertussis vaccine (DTP/DTaP/DT)
3. Two or more doses of any poliovirus vaccine
4. Two or more doses of hepatitis B vaccine
5. Three or more doses of DTaP and two or more doses of Polio
6. Three or more doses of DTaP, two or more doses of polio, and two or more doses of Hep B

**KINDERGARTEN RETROSPECTIVE SURVEY**  
CALIFORNIA, 2007

TABLE 7: IMMUNIZATION COVERAGE AMONG KINDERGARTEN STUDENTS AT 13 MONTHS OF AGE BY RACE/ETHNICITY<sup>1</sup>

	n	DTaP (3+) <sup>2</sup>	Polio (2+) <sup>3</sup>	MMR (1+) <sup>4</sup>	Hep B (2+) <sup>5</sup>	Var (1+) <sup>6</sup>	3:2:1 <sup>7</sup>	3:2:1:-:2 <sup>8</sup>	3:2:1:-:2:1 <sup>9</sup>
ALL	2,438	88.4% ± 1.8%	95.0% ± 1.0%	53.6% ± 3.0%	93.8% ± 1.1%	46.0% ± 3.0%	51.7% ± 3.0%	51.0% ± 2.9%	38.5% ± 2.8%
White	541	90.2% ± 3.2%	95.0% ± 2.0%	50.4% ± 5.1%	93.9% ± 2.0%	47.2% ± 6.4%	48.7% ± 5.1%	47.7% ± 5.0%	34.5% ± 5.2%
Hispanic	1,080	89.0% ± 2.3%	95.8% ± 1.4%	57.1% ± 4.0%	95.2% ± 1.3%	46.4% ± 3.6%	55.5% ± 3.9%	54.7% ± 3.7%	41.8% ± 3.6%
Black	346	78.8% ± 7.1%	93.5% ± 5.5%	43.4% ± 9.0%	92.6% ± 5.1%	34.2% ± 8.7%	38.3% ± 9.0%	38.2% ± 9.0%	28.3% ± 8.5%
Asian	223	90.5% ± 5.0%	95.5% ± 3.4%	57.6% ± 8.5%	92.7% ± 4.1%	50.3% ± 7.6%	56.1% ± 8.7%	56.1% ± 8.7%	41.2% ± 8.1%
Other	106	92.6% ± 6.6%	95.1% ± 5.1%	50.1% ± 10.4%	92.3% ± 6.8%	44.5% ± 9.1%	49.7% ± 10.4%	49.6% ± 10.4%	38.1% ± 9.4%
Unknown	142	78.7% ± 9.3%	87.6% ± 8.1%	43.2% ± 10.3%	83.9% ± 9.3%	44.9% ± 10.9%	40.8% ± 10.8%	38.3% ± 10.6%	35.4% ± 10.0%

1. As recorded on student immunization record (blue card). Individual racial groups do not include Hispanic children. Children of Hispanic ethnicity may be of any race
2. Three or more doses of any diphtheria and tetanus toxoids and pertussis vaccines including diphtheria and tetanus toxoids, and any acellular pertussis vaccine (DTP/DTaP/DT)
3. Two or more doses of any poliovirus vaccine
4. One or more doses of measles-mumps-rubella vaccine
5. Two or more doses of hepatitis B vaccine
6. One or more doses of varicella vaccine, unadjusted for history of varicella disease
7. Three or more doses of DTaP, two or more doses of Polio, and one or more doses of MMR
8. Three or more doses of DTaP, two or more doses of Polio, one or more doses of MMR, and two or more doses of Hep B
9. Three or more doses of DTaP, two or more doses of Polio, one or more doses of MMR, two or more doses of Hep B, and one or more doses of varicella

**KINDERGARTEN RETROSPECTIVE SURVEY**  
CALIFORNIA, 2007

TABLE 8: IMMUNIZATION COVERAGE AMONG KINDERGARTEN STUDENTS AT 19 MONTHS OF AGE BY RACE/ETHNICITY<sup>1</sup>

	n	DTaP (4+) <sup>2</sup>	Polio (3+) <sup>3</sup>	MMR (1+) <sup>4</sup>	Hep B (3+) <sup>5</sup>	Var (1+) <sup>6</sup>	4:3:1 <sup>7</sup>	4:3:1:-:3 <sup>8</sup>	4:3:1:-:3:1 <sup>9</sup>
ALL	2,438	65.6% ± 2.7%	85.3% ± 1.7%	88.0% ± 1.6%	86.5% ± 1.7%	79.5% ± 2.1%	61.7% ± 2.6%	59.2% ± 2.7%	55.5% ± 2.8%
White	541	65.9% ± 5.1%	82.1% ± 3.3%	87.5% ± 2.9%	87.8% ± 2.9%	80.1% ± 4.2%	60.6% ± 4.9%	59.2% ± 4.8%	53.9% ± 5.7%
Hispanic	1,080	65.5% ± 3.3%	87.6% ± 2.1%	89.3% ± 2.0%	88.1% ± 1.8%	79.1% ± 2.7%	62.3% ± 3.1%	59.6% ± 3.1%	56.5% ± 3.1%
Black	346	56.3% ± 9.4%	81.0% ± 8.4%	81.6% ± 7.6%	79.1% ± 9.0%	77.6% ± 8.3%	52.7% ± 9.4%	51.0% ± 9.4%	48.7% ± 9.5%
Asian	223	77.5% ± 7.4%	90.3% ± 5.0%	88.4% ± 4.8%	86.7% ± 5.4%	83.0% ± 5.8%	71.3% ± 7.8%	67.8% ± 8.3%	64.6% ± 8.6%
Other	106	59.6% ± 14.4%	82.5% ± 9.5%	87.7% ± 8.6%	83.5% ± 10.2%	79.2% ± 9.6%	58.1% ± 15.2%	56.0% ± 15.2%	52.5% ± 15.3%
Unknown	142	57.9% ± 11.2%	77.4% ± 9.4%	84.3% ± 9.5%	76.2% ± 9.6%	76.4% ± 10.0%	57.1% ± 11.4%	51.1% ± 11.6%	48.3% ± 10.9%

1. As recorded on student immunization record (blue card). Individual racial groups do not include Hispanic children. Children of Hispanic ethnicity may be of any race
2. Four or more doses of any diphtheria and tetanus toxoids and pertussis vaccines including diphtheria and tetanus toxoids, and any acellular pertussis vaccine (DTP/DTaP/DT)
3. Three or more doses of any poliovirus vaccine
4. One or more doses of measles-mumps-rubella vaccine
5. Three or more doses of hepatitis B vaccine
6. One or more doses of varicella vaccine, unadjusted for history of varicella disease
7. Four or more doses of DTaP, three or more doses of Polio, and one or more doses of MMR
8. Four or more doses of DTaP, three or more doses of Polio, one or more doses of MMR, and three or more doses of Hep B
9. Four or more doses of DTaP, three or more doses of Polio, one or more doses of MMR, three or more doses of Hep B, and one or more doses of varicella