

California Influenza Surveillance Project

California Department of Public Health

2009-2010

Influenza Update


During week 17 (April 25 – May 1, 2010), overall influenza activity in California remained “sporadic” (defined by the CDC as “small numbers of laboratory-confirmed influenza cases or a single laboratory-confirmed influenza outbreak has been reported, but there is no increase in cases of influenza-like-illness (ILI)”). Reports of ILI from sentinel providers decreased during *MMWR* week 17 (April 25 – May 1, 2010). The percentage of Kaiser hospitalizations for pneumonia and influenza (P&I) remained steady in northern California and decreased in southern California. Laboratory detections of influenza remain low. Detections of respiratory syncytial virus (RSV) declined.

A. Syndromic Surveillance

1. Kaiser Permanente Hospitalization Data (“Flu Admits”)

The admission diagnoses of flu, pneumonia, and influenza (“Flu Admits”) serve as surrogate markers for the more accurate, but less timely, diagnoses contained in discharge data. Influenza activity is tracked by dividing the number of Flu Admits by the total number of hospital admissions for the same day to obtain a percentage of influenza and pneumonia admissions. As indicated in the circle, Figure 1 shows that during week 17 (April 25 – May 1, 2010), the percentage of Kaiser hospitalizations for pneumonia and influenza (P&I) remained steady in northern California. Figure 2 shows a decrease in southern California. Both data points remain within the range of percentages seen for seasonal influenza in previous years.


Figures 1-2. Inpatient “Flu” Admissions at Kaiser Facilities, 2005-2010.



2. CDC Influenza Sentinel Providers

Sentinel providers report the number of outpatient visits for influenza-like illness (ILI) and the total number of visits per week. These data are reported weekly as a percentage of total visits. Figure 3 shows a peak in Weeks 17-18 (April 26 – May 9, 2009) when 2009 H1N1 influenza was first identified. ILI decreased sharply in week 17 (April 25 – May 1, 2010). As a result of a delay in reporting, the actual percentage may be higher. A total of 67 sentinel providers reported in Week 17.

Figure 3. California Sentinel Providers – Influenza-Like Visits, 2005-2010.



B. Laboratory Update

1. Respiratory Laboratory Network (RLN) Influenza PCR Surveillance Results

As noted in Table 1, during Week 17 (April 25 – May 1, 2010), none of the specimens received by the Respiratory Laboratory Network were positive for influenza.

Table 1. Respiratory Laboratory Network (RLN) Influenza PCR Surveillance Results from Selected Laboratories*, Week 17 (April 25 – May 1, 2010)

	Total Flu A tested	Flu A (% of total)	H1 (% of Flu A)	H3 (% of Flu A)	Unsubtypeable (% of Flu A)	Total Flu B tested	Flu B (% of total)	Total RSV tested (R-mix)	RSV (% of total)
Total RLN*	65	0 (0%)	0 (0%)	0 (0%)	0 (0%)	65	0 (0%)	14	0 (0%)
Northern	23	0 (0%)	0 (0%)	0 (0%)	0 (0%)	23	0 (0%)	0 (0%)	0 (0%)
Central	13	0 (0%)	0 (0%)	0 (0%)	0 (0%)	13	0 (0%)	0 (0%)	0 (0%)
Southern	29	0 (0%)	0 (0%)	0 (0%)	0 (0%)	29	0 (0%)	0 (0%)	0 (0%)

* 18 RLN laboratories reporting, including:

Northern CA: Contra Costa, El Dorado, Marin, Sacramento, San Francisco, Santa Clara, Shasta, Sonoma

Central CA: Fresno, San Joaquin, Tulare

Southern CA: Long Beach, Los Angeles, Orange, Riverside, San Diego, San Luis Obispo, Santa Barbara

2. Laboratory Positive Results Data

Table 2 shows positive influenza and other virus results from sentinel laboratories, local public health laboratories and VRDL.

Table 2. Influenza and other respiratory virus detections, April 25 – May 1, 2010.

		Sentinel Laboratories/Respiratory Laboratory Network [†]	Sentinel Providers
Week 17	Number	23 sites reporting	581 specimens submitted (267 positive by PCR, 6 pending)
	Influenza A	1 ^a Total tested week 17: 455	0
	Influenza B	0 Total tested week 17: 455	0
	RSV	17 ^b Total tested week 17: 420	N/A
	Other Respiratory Viruses	14 ^d Total tested week 17: 209	N/A

[†]Sentinel laboratories are hospital, academic, private, and public health laboratories located throughout California that provide data on the number of laboratory-confirmed influenza and other respiratory virus detections and isolations. The Respiratory Laboratory Network (RLN) is a network of 23 local public health laboratories that offer enhanced diagnostic testing with the "R-mix" shell vial assay, which detects several respiratory pathogens, including influenza A and B viruses, respiratory syncytial virus, parainfluenza virus, and adenovirus. Some RLN labs also offer PCR testing for influenza A and B.

^a Alameda (1)

^b Contra Costa (1); Fresno (1); Placer (1); Sacramento (3); San Diego (2); San Francisco (1); San Mateo (1); Santa Clara (6); Solano (1)

^c parainfluenza type 3 (5); rhinovirus (4); adenovirus (3); human metapneumovirus (1); parainfluenza type 1 (1)

Figure 4 shows that laboratory detections for influenza peaked in week 27 (July 5 - 11, 2009). Influenza remained low during the past month. Figure 5 shows that RSV has continued to decrease during the past month.

Figure 4. Influenza detections at sentinel laboratories/Respiratory Laboratory Network (RLN), 2006-2010.




Figure 5. RSV detections at sentinel laboratories, 2005-2010.

