

**SECTION: CERTIFICATION**

**SUBJECT: Eligibility Requirement**

**ITEM: Determining Clinical Nutrition Need for All Categories**

---

## **Purpose**

To provide local agencies (LA) standardized procedures for determining clinical risk when certifying women, infants and children for WIC benefits; to define allowable clinical risk criteria, and to indicate the priority level for each dietary risk and participant category.

## **Policy**

Qualified Staff shall assess whether a clinical need exists at each certification using approved nutrition assessment tools, the appropriate screens in the Integrated Statewide Information System (ISIS) and self-reported information.

## **Required Procedures**

- I. Clinical nutritional need shall be determined by evaluating the applicant's age, medical history, social, or other issues asked during individual counseling as part of the certification process.
- II. The LA shall document sources of information, other than the referral form or the nutrition questionnaire (NQ), in ISIS on the "Individual Nutrition Education Plan (INEP)" or in "Individual Comments."
- III. Self-reporting of a diagnosis by a medical professional shall not be confused with self-diagnosis, where a person simply claims to have or to have had a medical condition without any reference to professional diagnosis. A self-reported medical diagnosis ("My doctor says that I have/my son or daughter has...") should prompt the LA to validate the presence of the condition by asking more pointed questions related to that diagnosis.
- IV. The LA shall document identified indicators of clinical nutrition need in ISIS beginning with the appropriate "Health Information" screen. At a minimum, local agencies shall ask applicants all questions contained on the "Health Questions" and "Additional Health Questions" screens. Other clinical indicators of nutrition needs can be selected using the option key in the "other conditions" field.

- V. The LA shall ask the supplemental alcohol and/or drug questions in ISIS if a woman admits that she is currently using drugs or alcohol or has a history of an alcohol or drug problem.

## **Authority**

7 CFR 246.7 (e)

USDA Policy Memorandum 98-9, Revisions 1, 7, 8, 9, and 10; Nutrition Risk Criteria Value Enhanced Nutrition Assessment (VENA)

WRO Policy Memo 11/25/13: Transmittal of Revised Nutrition Risk Criteria

## **Resources**

USDA Risk Descriptions:

<http://www.cdph.ca.gov/programs/wicworks/Pages/NutritionRiskDescriptions.aspx>;

Protocols for various nutrition risk criteria and template for developing protocols at:  
<http://www.cdph.ca.gov/programs/wicworks/Pages/WIC-PEPB-Protocols.aspx>.

## CLINICAL RISK CRITERIA

**WIC 210-12**

The table below includes the clinical criteria to assess nutrition need of all categories of WIC applicants. The priority level is indicated for each category and the ISIS risk code and corresponding USDA risk code is provided for reference. “N” indicates that ISIS will generate the question “Do you want to see the Nutritionist?” On the Summary screen when this risk code is identified.

ISIS CODE (USDA CODE)	INDICATOR OF NUTRITIONAL NEED DEFINITION	PRIORITIES				
		PREGNANT WOMEN	BREASTFEEDING WOMEN	NON-BREASTFEEDING WOMEN	INFANTS	CHILDREN
<b>C10*</b> (302)	<b>Current Gestational Diabetes Mellitus</b>	I N				
<b>C11*</b> (303)	<b>History of Gestational Diabetes Mellitus</b>	Any history I	Most recent pregnancy I	Most recent pregnancy III		
<b>C12*</b> (345)	<b>Current Pregnancy Hypertension</b>	I N				
<b>C14*</b> (336)	<b>Fetal Growth Restriction (FGR):</b> usually defined as a fetal weight < 10th percentile for gestational age	I N				
<b>C15*</b> (301)	<b>Hyperemesis Gravidarum:</b> severe nausea and vomiting to the extent that the pregnant woman becomes dehydrated and acidotic)	I N				
<b>C17*</b> (304)	<b>History of Preeclampsia</b>	I N	I N	III N		

## CLINICAL RISK CRITERIA

WIC 210-12

ISIS CODE (USDA CODE)	INDICATOR OF NUTRITIONAL NEED DEFINITION	PRIORITIES				
		PREGNANT WOMEN	BREASTFEEDING WOMEN	NON-BREASTFEEDING WOMEN	INFANTS	CHILDREN
<b>C20</b> (311)	<b>History of Preterm Delivery:</b> ≤ 37 weeks	Any history I	Most recent pregnancy with one or more infants living I	Most recent pregnancy III		
<b>C21</b> (312)	<b>History of Low Birth Weight:</b> < 2500 g or 5 lbs. 8 oz	Any history I	Most recent pregnancy I	Most recent pregnancy III		
<b>C30*</b> (321)	<b>Fetal Death:</b> > 20 weeks gestation	Any history I	Most recent pregnancy I	Most recent pregnancy VI		
<b>C31*</b> (321)	<b>Neonatal Death:</b> < 28 days after birth	Any history I	Most recent pregnancy I	Most recent pregnancy VI		
<b>C33*</b> (321)	<b>History of Spontaneous Abortion, Fetal or Neonatal Loss:</b> termination at <20 weeks gestation or <500 g. History of fetal/neonatal death or 2 or more spontaneous abortions	I	Most recent pregnancy I	Most recent pregnancy only III		
<b>C44</b> (332)	<b>Closely Spaced Pregnancy:</b> conception before 16 months postpartum	Current pregnancy I	Most recent pregnancy I	Most recent pregnancy VI		
<b>C45</b> (335)	<b>Multi-fetal Gestation</b>	Current pregnancy I	Most recent pregnancy I	Most recent pregnancy III		

## CLINICAL RISK CRITERIA

WIC 210-12

ISIS CODE (USDA CODE)	INDICATOR OF NUTRITIONAL NEED DEFINITION	PRIORITIES				
		PREGNANT WOMEN	BREASTFEEDING WOMEN	NON-BREASTFEEDING WOMEN	INFANTS	CHILDREN
<b>C47*</b> (337)	<b>History of a Large for Gestational Age Infant:</b> history of giving birth to an infant weighing $\geq 9$ lbs. ( $\geq 4,000$ g)	Any history I	Most recent pregnancy I	Most recent pregnancy VI		
<b>C48*</b> (339)	<b>History of Birth with Nutrition Related Congenital or Birth Defect:</b> linked to inappropriate nutritional intake (e.g., inadequate zinc, folic acid, excess vitamin A)	Any history I	Most recent pregnancy I	Most recent pregnancy III		
<b>C49*</b> (363)	<b>Pre-Diabetes</b>		I N	III N		
<b>C50*</b> (343)	<b>Diabetes Mellitus: Type 1 or 2</b>	I N	I N	III N		
<b>C51*</b> (345)	<b>Hypertension/Prehypertension</b>	I N	I N	III N		
<b>C51*</b> (359)	<b>Recent Major Surgery or Trauma:</b> severity affects nutritional status within 2 months self-reported, or >2 months if nutritional support diagnosed by a physician				I	III
<b>C52*</b> (352)	<b>Active TB:</b> present within the past 6 months	I N	I N	III N	I N	III N

## CLINICAL RISK CRITERIA

WIC 210-12

ISIS CODE (USDA CODE)	INDICATOR OF NUTRITIONAL NEED DEFINITION	PRIORITIES				
		PREGNANT WOMEN	BREASTFEEDING WOMEN	NON-BREASTFEEDING WOMEN	INFANTS	CHILDREN
<b>C53*</b> (346)	<b>Current Renal (Kidney) Disease:</b> includes pyelonephritis, persistent proteinuria (excluding urinary tract infections involving the bladder)	I N	I N	III N		
<b>C53*</b> (352)	<b>Severe Acute Infections:</b> within the past 6 months which affect nutritional status. Includes: encephalitis, hepatitis, pneumonia, meningitis, and parasites				I	III
<b>C54</b> (359)	<b>Recent Burns:</b> severity affects nutritional status within the past 2 months self-reported, or > 2 months if there is continued need for nutritional support diagnosed by a physician				I	III
<b>C55*</b> (360)	<b>Current Cardio/Pulmonary Diseases:</b> severe enough to affect nutritional status	I N	I N	III N		
<b>C56*</b> (342, 354)	<b>Current Gastrointestinal Disorders:</b> conditions which interfere with intake or absorption of nutrients. They include but are not limited to post-bariatric surgery; stomach or intestinal ulcers; small bowel enterocolitis and syndrome; inflammatory bowel disease, ulcerative colitis - Crohn's disease; liver disease; pancreatitis; gall bladder disease; Celiac Sprue; Gluten enteropathy, Non-tropical Sprue	I N	I N	III N		

## CLINICAL RISK CRITERIA

WIC 210-12

ISIS CODE (USDA CODE)	INDICATOR OF NUTRITIONAL NEED DEFINITION	PRIORITIES				
		PREGNANT WOMEN	BREASTFEEDING WOMEN	NON-BREASTFEEDING WOMEN	INFANTS	CHILDREN
<b>C57*</b> (344)	<b>Current Thyroid Disorders:</b> hyperthyroid or hypothyroid conditions. <b>Postpartum Thyroiditis</b>	I N	I N	III N		
<b>C57*</b> (381)	<b>Oral Health Conditions:</b> diagnosis of oral health conditions by a physician, dentist or someone working under a physician's order or as self reported by applicant/participant/caregiver which impair the ability to ingest food in adequate quantity or quality. May include nursing or baby bottle caries; decay of maxillary anterior/primary molars; tooth decay; periodontal disease; and tooth loss.				I	III
<b>C58*</b> (347)	<b>Current Cancer:</b> treatment or condition affects nutritional status	I N	I N	III N		
<b>C59*</b> (348)	<b>Current Central Nervous System (CNS) Disorders:</b> conditions which affect nutritional status metabolically and/or mechanically. Includes epilepsy; cerebral palsy (CP); neural tube defects (NTD) (e.g., spina bifida or myelomeningocele); Parkinson's disease; and multiple sclerosis	I N	I N	III N		
<b>C60*</b> (382)	<b>Drug Exposure in Utero:</b> Fetal Alcohol Syndrome based on the presence of retarded growth, a pattern of facial and central nervous system abnormalities including mental retardation				I	III

## CLINICAL RISK CRITERIA

WIC 210-12

ISIS CODE (USDA CODE)	INDICATOR OF NUTRITIONAL NEED DEFINITION	PRIORITIES				
		PREGNANT WOMEN	BREASTFEEDING WOMEN	NON-BREASTFEEDING WOMEN	INFANTS	CHILDREN
<b>C60*</b> (703)	<b>Drug Exposure in Utero:</b> documentation or self-reported use of alcohol or illegal drugs during most recent pregnancy				I	
<b>C60</b> (371)	<b>Smoking 1:</b> 1-7 cigs/d	I	I	III		
<b>C62</b> (372)	<b>Drugs:</b> any illegal	I N	I N	III N		
<b>C63</b> (371)	<b>Maternal Smoking 2:</b> 7-14 cigs/d	I	I	III		
<b>C64</b> (371)	<b>Maternal Smoking 3:</b> 15-20 cigs/d	I	I	III		
<b>C65</b> (371)	<b>Maternal Smoking 4:</b> 21+ cigs/d	I	I	III		
<b>C70*</b> (348)	<b>Central Nervous System (CNS) Disorders:</b> conditions affecting energy requirements and an individual's ability to feed self, altering nutritional status metabolically, mechanically, or both. May include epilepsy; cerebral palsy; neural tube defects (e.g., spina bifida); Parkinson's disease; and multiple sclerosis				I N	III N

## CLINICAL RISK CRITERIA

WIC 210-12

ISIS CODE (USDA CODE)	INDICATOR OF NUTRITIONAL NEED DEFINITION	PRIORITIES				
		PREGNANT WOMEN	BREASTFEEDING WOMEN	NON-BREASTFEEDING WOMEN	INFANTS	CHILDREN
<b>C71*</b> (349)	<b>Down Syndrome:</b> condition must alter nutritional status metabolically, and/or mechanically				I N	III N
<b>C72</b> (359)	<b>C-Section in Most Recent Pregnancy:</b> within the past 2 months		I	VI		
<b>C72</b> (362)	<b>Developmental Delays, Sensory or Motor Delays:</b> interfering with the ability to chew or swallow food, or requiring a tube feeding. May include mental retardation; birth injury; head trauma; minimal brain function				I N	III N
<b>C73*</b> (360)	<b>Cystic Fibrosis:</b> current condition or treatment for the condition must be severe enough to affect nutritional status				I N	III N
<b>C74*</b> (360)	<b>Congenital Heart Disease:</b> current condition or treatment for the condition must be severe enough to affect nutritional status				I N	III N
<b>C75*</b> (342, 349)	<b>Congenital GI Disorders:</b> interfering with nutritional status. May include tracheal-esophageal fistula; esophageal or intestinal atresia; gastroschisis; short bowel syndrome				I N	III N
<b>C76*</b> (349)	<b>Congenital Facial Malformation:</b> causes physical abnormality that alters nutritional status (e.g., cleft lip or palate)				I N	III N

## CLINICAL RISK CRITERIA

WIC 210-12

ISIS CODE (USDA CODE)	INDICATOR OF NUTRITIONAL NEED DEFINITION	PRIORITIES				
		PREGNANT WOMEN	BREASTFEEDING WOMEN	NON-BREASTFEEDING WOMEN	INFANTS	CHILDREN
<b>C77*</b> (349)	<b>Other Genetic and Congenital Disorder(s):</b> causing physical or metabolic abnormality, which alters nutritional status. May include Down Syndrome; thalassemia; sickle cell anemia; or muscular dystrophy				I N	III N
<b>C80*</b> (342, 354)	<b>Gastro-Intestinal Disorders:</b> interfere with the intake or absorption of nutrients. May include stomach or intestinal disorders; small bowel enterocolitis; malabsorption syndrome; liver disease; or gallbladder disease, Celiac disease also known as: Celiac Sprue, Gluten-sensitive enteropathy, or Non-tropical sprue.				I N	III N
<b>C80*</b> (352)	<b>Severe Acute Infections:</b> within last 6 months which affect nutritional status. Includes: pneumonia; meningitis; parasitic infection; bronchiolitis (3 episodes/6 months); Rubella; or listeriosis.	I	I	III		
<b>C81*</b> (360)	<b>Cardiovascular Diseases:</b> condition and/or treatment severe enough to affect nutritional status (e.g., cardiorespiratory diseases)				I N	III N
<b>C82*</b> (352)	<b>Immunodeficiency or Chronic Infections:</b> condition severe enough to affect nutritional status. Includes hepatitis; HIV (Human Immunodeficiency Virus infection; AIDS (Acquired Immunodeficiency Disease).	I Present within past 6 months N	I Present within the past 6 months N	III Present within the past 6 months N	I N	III N

## CLINICAL RISK CRITERIA

WIC 210-12

ISIS CODE (USDA CODE)	INDICATOR OF NUTRITIONAL NEED DEFINITION	PRIORITIES				
		PREGNANT WOMEN	BREASTFEEDING WOMEN	NON-BREASTFEEDING WOMEN	INFANTS	CHILDREN
<b>C83*</b> (347)	<b>Cancer:</b> for which condition or treatment severe enough to affect nutritional status				I	III
<b>C84</b> (338)	<b>Pregnant Woman Currently Breastfeeding</b>	I				
<b>C85*</b> (343)	<b>Diabetes Mellitus:</b> Type 1 or 2				I N	III N
<b>C86*</b> (346)	<b>Current Renal (Kidney) Disease:</b> includes pyelonephritis and persistent proteinuria, but excludes urinary tract infections involving the bladder				I N	III N
<b>C86*</b> (351)	<b>Inborn Errors of Metabolism:</b> gene mutations or gene deletions that alter metabolism. Includes: <b>Amino Acid Disorders, Organic Acid Metabolism Disorders, Fatty Acid Oxidation Disorders, Lysosomal Storage Diseases, Urea Cycle Disorders, Carbohydrate Disorders, Peroxisomal disorders, and Mitochondrial Disorders.</b>  <b>See WPM Appendix: #960-04</b>	I  N	I  N	III  N		
<b>C87*</b> (351)	<b>Inborn Errors of Metabolism:</b> same as above				I N	III N

## CLINICAL RISK CRITERIA

WIC 210-12

ISIS CODE (USDA CODE)	INDICATOR OF NUTRITIONAL NEED DEFINITION	PRIORITIES				
		PREGNANT WOMEN	BREASTFEEDING WOMEN	NON-BREASTFEEDING WOMEN	INFANTS	CHILDREN
<b>C89*</b> (353)	<b>Food Allergies:</b> an adverse immune response to a food or a hypersensitivity causing adverse immunologic reaction				I N	III N
<b>C90*</b> (703)	<b>Infant Born of Woman with Mental Retardation or Developmental Delay</b>				I	
<b>C90</b> (362)	<b>Developmental Delays, Sensory or Motor Delays:</b> interfering with ability to eat, restricting the ability to chew or swallow food, or requiring tube feeding. Includes minimal brain function; feeding problems due to developmental delay; head trauma; brain damage or birth injury	I N	I N	III N		
<b>C91</b> (801)	<b>Homeless:</b> a woman, infant, or child who lacks a fixed and regular nighttime residence; or whose primary nighttime residence is: a supervised publicly or privately operated shelter (e.g., welfare hotel, a congregate shelter, or a shelter for victims of domestic violence) designed to provide temporary living accommodations; an institution that provides a temporary residence for individuals intended to be institutionalized; temporary accommodation of not more than 365 days in the residence of another individual; or a public or private place not designed for use as a regular sleeping accommodation for human beings	IV	IV	VI	IV	V

## CLINICAL RISK CRITERIA

WIC 210-12

ISIS CODE (USDA CODE)	INDICATOR OF NUTRITIONAL NEED DEFINITION	PRIORITIES				
		PREGNANT WOMEN	BREASTFEEDING WOMEN	NON-BREASTFEEDING WOMEN	INFANTS	CHILDREN
<b>C93*</b> (901)	<b>Recipient of Abuse:</b> within the past 6 months, includes any recent act or failure to act resulting in imminent risk of serious harm, death, serious physical or emotional harm or exploitation of an infant or child by a parent or caregiver, and battering or violent physical assaults on women	IV	IV	VI	IV	V
<b>C96*</b> (358)	<b>Eating Disorder:</b> (e.g. anorexia, bulimia) characterized by a disturbed sense of body image and a morbid fear of becoming fat. Symptoms may include abnormal eating patterns; self-induced vomiting; laxative or enema use; periods of starvation; or use of drugs(i.e., appetite suppressants or diuretics)	I  N	I  N	III  N		
<b>C98*</b> (802)	<b>Migrant:</b> all categories are eligible who are members of families which contain at least one individual whose principal employment is in agriculture on a seasonal basis, who has been so employed within the last 24 months, and who establishes, for the purposes of such employment, a temporary abode	IV	IV	VI	IV	V
<b>C99*</b> (903)	<b>Foster Care:</b> entering foster care system within past 6 months, or moving from one foster care home to another in the past 6 months. <u>Note:</u> C99 cannot be used for consecutive certifications while the child remains in the same foster home	IV	IV	VI	IV	V

## CLINICAL RISK CRITERIA

WIC 210-12

ISIS CODE (USDA CODE)	INDICATOR OF NUTRITIONAL NEED DEFINITION	PRIORITIES				
		PREGNANT WOMEN	BREASTFEEDING WOMEN	NON-BREASTFEEDING WOMEN	INFANTS	CHILDREN
<b>C100*</b> (903)	<b>Nutrient Deficiency Disease:</b> diagnosis of a nutritional deficiency or a disease caused by insufficient dietary intake of macro and micro nutrients. May include Protein Energy Malnutrition (PEM); scurvy; rickets; beriberi; hypocalcemia; Osteomalacia; vitamin K deficiency; pellagra; cheilosis; Menkes Disease, and xerophthalmia	I  N	I  N	III  N	I  N	III  N
<b>C101</b> (331)	<b>Young Teenagers:</b> age ≤15 years at conception	Current pregnancy I	Most recent pregnancy I	Most recent pregnancy VI		
<b>C102</b> (331)	<b>Teenagers:</b> age 16-17 years at conception	Current pregnancy I	Most recent pregnancy I	Most recent pregnancy VI		
<b>C102*</b> (344)	<b>Thyroid Disorders:</b> hypothyroid or hyperthyroid conditions. (includes congenital hyperthyroidism and congenital hypothyroidism)				I  N	III  N
<b>C103</b> (333)	<b>High Parity and Young Age:</b> under age 20 at conception, and 3 or more previous pregnancies (≥ 20 weeks gestation)	Current pregnancy I	Most recent pregnancy I	Most recent pregnancy III		
<b>C104*</b> (356)	<b>Hypoglycemia</b>	I  N	I  N	III  N	I  N	III  N

## CLINICAL RISK CRITERIA

WIC 210-12

ISIS CODE (USDA CODE)	INDICATOR OF NUTRITIONAL NEED DEFINITION	PRIORITIES				
		PREGNANT WOMEN	BREASTFEEDING WOMEN	NON-BREASTFEEDING WOMEN	INFANTS	CHILDREN
<b>C105*</b> (360)	<b>Other Medical Diseases and Conditions:</b> with nutritional implications severe enough to affect nutritional status. May include juvenile rheumatoid arthritis (JRA); lupus erythematosus or cystic fibrosis	I N	I N	III N		
<b>C106*</b> (349)	<b>Genetic and Congenital Disorders:</b> which alter nutritional status metabolically and/or mechanically. Includes: Down's syndrome; cleft lip or palate; thalassemia major; sickle cell anemia ;muscular dystrophy	I N	I N	III N		
<b>C107</b> (357)	<b>Drug Nutrient Interactions:</b> any use of prescription or over the counter medications that may interfere with nutritional status	I N	I N	III N	I N	III N
<b>C110*</b> (134)	<b>Failure to Thrive</b>				I N	III N
<b>C110*</b> (353)	<b>Food Allergies:</b> an adverse immune response to a food or a hypersensitivity that causes adverse immunologic reaction	I N	I N	III N		
<b>C111*</b> (355)	<b>Lactose Intolerance:</b> insufficient production of the enzyme lactase which causes an inability to digest lactose, producing any or all of the following GI disturbances: nausea; diarrhea; abdominal bloating; cramps	I	I	III	I	III

## CLINICAL RISK CRITERIA

WIC 210-12

ISIS CODE (USDA CODE)	INDICATOR OF NUTRITIONAL NEED DEFINITION	PRIORITIES				
		PREGNANT WOMEN	BREASTFEEDING WOMEN	NON-BREASTFEEDING WOMEN	INFANTS	CHILDREN
C112 (359)	<b>Recent Major Surgery, Trauma, Burns:</b> severity compromises nutritional status, or if nutrition support is diagnosed by a physician	I	I	III		
C113* (361)	<b>Depression</b> Presence of clinical depression, including postpartum depression. No longer a risk for children.	I	I	III		
C114* (381)	<b>Oral Health Conditions:</b> diagnosis of oral health conditions by a physician, dentist or someone working under a physician's order or as self reported by applicant/participant/caregiver. Includes but not limited to tooth decay; periodontal disease; gingivitis, tooth loss; ineffectively replaced teeth or oral infections which impair the ability to ingest food in adequate quantity or quality.	I	I	VI		
C115 (372)	<b>Alcohol Use:</b> any use in pregnancy	I				
C115 (372)	<b>Alcohol Use:</b> regular use of ≥ 2 drinks/day; or binge drinking ≥ 5 drinks on at least one day in past 30 days; or heavy drinking ≥ 5 drinks on same occasion on 5 or more days in the past 30 days. (One serving = 1 can of beer, 5 oz. wine, or 1 ½ oz. liquor)		I	III		

## CLINICAL RISK CRITERIA

WIC 210-12

ISIS CODE  (USDA CODE)	INDICATOR OF NUTRITIONAL NEED DEFINITION	<u>PRIORITIES</u>																
		PREGNANT WOMEN	BREASTFEEDING WOMEN	NON-BREASTFEEDING WOMEN	INFANTS	CHILDREN												
<b>C119</b> (334)	<b>Lack of or Inadequate Prenatal Care:</b> beginning after 1 <sup>st</sup> trimester or based on the Inadequate Prenatal Index (Kessner et al):	I	N/A	N/A														
	<table border="1" style="margin-left: auto; margin-right: auto; border-collapse: collapse;"> <thead> <tr> <th style="text-align: center;">Weeks Gestation</th> <th style="text-align: center;">Prenatal Visits</th> </tr> </thead> <tbody> <tr> <td style="text-align: center;">14-21</td> <td style="text-align: center;">0/unknown</td> </tr> <tr> <td style="text-align: center;">22-29</td> <td style="text-align: center;">1 or fewer</td> </tr> <tr> <td style="text-align: center;">30-31</td> <td style="text-align: center;">2 or fewer</td> </tr> <tr> <td style="text-align: center;">32-33</td> <td style="text-align: center;">3 or fewer</td> </tr> <tr> <td style="text-align: center;">34 or more</td> <td style="text-align: center;">4 or fewer</td> </tr> </tbody> </table>	Weeks Gestation	Prenatal Visits	14-21	0/unknown	22-29	1 or fewer	30-31	2 or fewer	32-33	3 or fewer	34 or more	4 or fewer					
Weeks Gestation	Prenatal Visits																	
14-21	0/unknown																	
22-29	1 or fewer																	
30-31	2 or fewer																	
32-33	3 or fewer																	
34 or more	4 or fewer																	
<b>C121</b> (902)	<b>Woman, Infant/Child of Primary Caregiver with Limited Ability to Make Feeding Decisions and/or Prepare Food:</b> individuals who are ≤ 17 years of age; mentally disabled/delayed and/or have a mental illness such as clinical depression (diagnosed by a physician or licensed psychologist); current use or history of alcohol or drug abuse; or physical disability which limits food preparation abilities	IV	IV	VI	IV	V												
<b>C122*</b> (360)	<b>Persistent Asthma, moderate or severe:</b> current requiring daily medications and severe enough to affect nutritional status	I	I	III	I	III												

## CLINICAL RISK CRITERIA

WIC 210-12

ISIS CODE (USDA CODE)	INDICATOR OF NUTRITIONAL NEED DEFINITION	PRIORITIES				
		PREGNANT WOMEN	BREASTFEEDING WOMEN	NON-BREASTFEEDING WOMEN	INFANTS	CHILDREN
<b>C123*</b> (360)	<b>Other Medical Diseases or Conditions:</b> nutritional implications severe enough to affect nutritional status. Includes: juvenile rheumatoid arthritis; lupus erythematosus				I N	III N
<b>C201</b> (603)	<b>Breastfed Infant with Suck Problem:</b> ineffective or weak suck				I N	
<b>C201</b> (602)	<b>Cracked &amp; Bleeding Nipples</b>		I			
<b>C202</b> (602)	<b>Persistent Sore Nipples</b>		I			
<b>C203</b> (602)	<b>Mastitis</b> (breast infection)		I N			
<b>C204</b> (603)	<b>Jaundice While Breastfeeding</b>				I N	
<b>C204</b> (602)	<b>Flat or Inverted Nipples</b>		I			
<b>C205</b> (602)	<b>Recurrent Plugged Ducts</b>		I			

## CLINICAL RISK CRITERIA

WIC 210-12

ISIS CODE (USDA CODE)	INDICATOR OF NUTRITIONAL NEED DEFINITION	PRIORITIES				
		PREGNANT WOMEN	BREASTFEEDING WOMEN	NON-BREASTFEEDING WOMEN	INFANTS	CHILDREN
<b>C206</b> (602)	<b>Failure of Milk to Come in by 4 Days Postpartum</b>		I			
<b>C208</b> (602)	<b>Severe Breast Engorgement</b>		I			
<b>C209</b> (602)	<b>Tandem Nursing</b>		I			
<b>C209</b> (603)	<b>Infant with Difficulty Latching onto Mother's Breast</b>				I	
<b>C210</b> (602)	<b>Breastfeeding Woman <math>\geq</math> 40 Years Old</b>		I			
<b>C904</b> (904)	<b>Environmental Tobacco Smoke (ETS) (i.e., second hand smoke) Exposure in the Home</b>	I	I	III	I	III
<b>N32</b> (601, 702)	<b>Breastfeeding a WIC Infant and Breastfeeding Infant of Woman at Nutritional Risk:</b> both the breastfeeding woman and her breastfed infant must be assigned the same, highest priority for which either is eligible		I, II or IV		I, II or IV	

**CLINICAL RISK CRITERIA**

**WIC 210-12**

ISIS CODE (USDA CODE)	INDICATOR OF NUTRITIONAL NEED DEFINITION	PRIORITIES				
		PREGNANT WOMEN	BREASTFEEDING WOMEN	NON-BREASTFEEDING WOMEN	INFANTS	CHILDREN
N40 (701)	Infant Born to Woman on WIC During Pregnancy				Only 6 months II	
N41 (701)	Infant Born to WIC Eligible Woman During Her Pregnancy				Only 6 months II	
*Condition diagnosed by a physician and self-reported by the applicant/participant/caregiver, or as reported or documented by a physician or someone working under a physician's orders.						