Home (/) Asthma Data County Asthma Profiles (/asthma-data/county-asthma-profiles) Los Angeles County Asthma Profile

Los Angeles County Asthma Profile

modified on: Thursday, 01 September 2016

In Los Angeles County, approximately 1,221,000 children and adults have been diagnosed with asthma.

Scroll down to browse the full County Profile. Or, download a printer-friendly PDF.

COUNTY PROFILE INDICATORS

Lifetime Asthma Prevalence	Emergency Department Visits
Active Asthma Prevalence	Asthma Hospitalizations
Work-Related Asthma	Asthma Deaths
Asthma Management Plans	Outdoor Air Quality ¹⁵
Asthma Risk Factors	Notes

	County Population			County Population		
Age Population %		Race/Ethnicity ¹ %				
	0-4	670,558	6.7	Asian/PI	13.9	
	5-17	1,651,655	16.5	Black	7.9	
	18-64	6,497,777	64.8	Hispanic	49.5	
	65+	1,205,588	12.0	White	26.5	
	Total	10,025,579		Other	2.2	

DATASOURCE:California Department of Finance, 2014 (2013 release)

LIFETIME ASTHMA PREVALENCE,2 2011-2012

Percent with Lifetime Asthma (95% Confidence Interval³)

	Age	Los Angeles County	California
Children	0-4		9.7 (7.2-12.3)
	5-17	15.6 (12.3-19.0)	17.1 (15.5-18.7)
Adults	18-64	12.4 (11.2-13.6)	14.0 (13.4-14.7)
	65+	10.4 (8.6-12.1)	12.0 (11.1-12.8)
Totals:	0-17	14.5 (11.5-17.5)	15.4 (14.0-16.7)
	18+	12.1 (11.1-13.1)	13.7 (13.1-14.3)
	All Ages	12.7 (11.6-13.7)	14.1 (13.6-14.6)

datasource: California Health Interview Survey (CHIS), 2014

ACTIVE ASTHMA PREVALENCE,4 2014

Percent with Active Asthma (95% Confidence Interval³)

	Age	Los Angeles County	California
Children	0-4	0.2 (0.0-0.4)	6.3 (4.6-8.0)
	5-17	10.2 (5.7-14.8)	11.6 (10.2-13.0)
Adults	18-64	6.2 (4.6-7.9)	7.6 (7.1-8.1)
	65+	7.1 (4.0-10.2)	8.2 (7.5-8.9)
Totals:	0-17	7.3 (4.0-10.5)	10.1 (9.0-11.3)
	18+	6.4 (4.9-7.9)	7.7 (7.3-8.2)
	All Ages	7.8 (6.9-8.7)	8.3 (7.9-8.7)

datasource: California Health Interview Survey (CHIS), 2014

WORK-RELATED ASTHMA5

Studies show that asthma is commonly caused or triggered by workplace exposures, but work-related asthma

(http://www.cdph.ca.gov/programs/ohsep/Pages/Asthma.aspx) is under-recognized and under-diagnosed. Research confirms that 15-30% of current adult asthma was initiated by work exposures, meaning that an estimated 71,400-142,800 adults in Los Angeles County have asthma caused by work.6

datasource: CHIS, 2014

ASTHMA MANAGEMENT PLANS

National guidelines recommend that health care providers give all patients with asthma a written self-management plan. In Los Angeles County, 84.4% (95% CI 80.2-88.7) of people with asthma have NOT received a written asthma management plan from a health care provider.

datasource: CHIS, 2014

ASTHMA RISK FACTORS, 2014

Los Angeles County Risk Factors

Interval)
13.2 (12.0-14.3)
6.7 (6.0-7.4)
24.7 (23.3-26.1)
15.4 (NA)
11.6 (NA)

ASTHMA DEATHS,9 2008-2010

Number of Deaths Due to Asthma (N) and Age-Adjusted Rate⁹ (per 1,000,000

		Los Ange	eles County	Califo	ornia
	Age	N	Rate	N	Rate
Children	0-17	14	1.7	57	1.9
Adults	18+	318	14.3	1,198	14.3
Totals	All Ages	332	11.0	1,255	11.1

datasource: California Death Statistical Master Files, 2008-2010

ASTHMA EMERGENCY DEPARTMENT VISITS,10 2014

Number of ED Visits Due to Asthma (N) and Age-Adjusted Rate¹¹ (per 10,000 residents)

		Los Angele	s County	Califo	rnia
	Age	N	Rate	N	Rate
Children	0-4	7,079	106.9	26,268	103.4
	5-17	13,688	83.4	47,613	72.4
Adults	18-64	25,597	39.3	95,689	39.2
	65+	4,573	38.3	17,374	36.1
Totals:	0-17	20,767	89.7	73,881	80.7
	18+	30,170	39.1	113,063	38.6
	All Ages	50,937	52.2	186,944	49.5

datasource: Office of Statewide Health Planning and Development (OSHPD), 2014 Expected Source of Payment for Asthma ED Visits

Payment Source	Los Angeles County	California
Medicare	10.97%	13.29%
Medi-Cal	50.06%	48.56%
Private	26.86%	26.32%
Other	12.10%	11.84%

datasource: Office of Statewide Health Planning and Development (OSHPD), 2014

ASTHMA HOSPITALIZATIONS,12 2014

Number of Hospitalizations Due to Asthma (N) and Age-Adjusted Rate¹¹ (per 10,000 residents)

		Los Angel	es County	Califo	rnia
	Age	N	Rate	N	Rate
Children	0-4	1,333	20.0	4,994	19.6
	5-17	1,534	9.4	5,037	7.7
Adults	18-64	3,876	5.7	12,134	4.7
	65+	2,447	20.7	7,208	15.2
Totals:	0-17	2,867	12.2	10,031	10.9
	18+	6,323	8.3	19,342	6.5
	All Ages	9,190	9.3	29,373	7.6

datasource: Office of Statewide Health Planning and Development (OSHPD), 2014 Average Charges¹³ Per Asthma Hospitalization

Age	Los Angeles County	California
0-17	\$23,254.06	\$27,127.04

Children Adults \$42,843.40 \$46,783.17 Total All Ages \$36,535.98 \$39,860.44 datasource: Office of Statewide Health Planning and Development (OSHPD), 2014


Expected Source of Payment for Asthma Hospitalizations

Payment Source	Los Angeles County	California
Medicare	28.78%	28.24%
Medi-Cal	48.28%	46.46%
Private	18.99%	20.34%
Other	3.95%	4.96%

datasource: Office of Statewide Health Planning and Development (OSHPD), 2014

ASTHMA DISPARITIES


Age-Adjusted Asthma Hospitalizations and ED Visits per 10,000 Los Angeles County Residents by Race/Ethnicity, 2014



Data Source: datasource: Office of Statewide Health Planning and Development (OSHPD), 2014

HOSPITALIZATION RATES OVER TIME


Age-Adjusted Asthma Hospitalizations per 10,000 Residents, Los Angeles County and California, 2004-2014




Data Source: datasource: Office of Statewide Health Planning and Development (OSHPD), 2004-2014

HEALTHY PEOPLE 2010¹⁴

Asthma Hospitalizations per 10,000 Residents by Age, Compared to HP2020 Targets, California and Los Angeles County, 2014



Asthma ED Visits per 10,000 Residents by Age, Compared to HP2020 Targets, California and Los Angeles County, 2014



Data Source: datasource: Office of Statewide Health Pla

NOTES


Estimates marked with an asterisk () have large confidence intervals, interpret with caution. Estimates marked with a dash (-) are statistically unstable, so are unreported.

- 1. PI = Pacific Islander; Please see technical notes for more information on race/ethnicity categorizations
- Lifetime asthma prevalence is the proportion of people in the population who have ever been diagnosed with asthma by a health provider.
 The 95% confidence interval (CI) is a range that expresses a level of certainty about an estimate based on the margin of error.
- The 95% CI means that we are 95 percent confident that this range contains the true population percent. A narrow CI means that there is less variability in the estimate and/or there is a larger sample size. A wide CI indicates more variability and/or a smaller sample size.

 4. Active asthma prevalence is the proportion of people in the population who have ever been diagnosed with asthma by a health provider and report that they still have asthma
- and/or report that they had an episode or attack within the past 12 months.
- 5. Work-related asthma is asthma that is caused or triggered by conditions or substances in the workplace.
- 6. Balmes J, Becklake M, Blanc P, et al. Environmental and Occupational Health Assembly, American Thoracic Society. American Thoracic Society Statement: Occupational Contribution to the Burden of Airway Disease. Am J Respir Crit Care Med. 2003;167:787-797; Lutzker L, Rafferty A, Brunner W, et al. Prevalence of Work-related Asthma in Michigan, Minnesota, and Oregon. Journal of Asthma. 2010;47:156-161.
- Obesity is defined as a body mass index (BMI) of 30 or greater.
- Data Sources for Asthma Risk Factors: Smoking CHIS, 2009; Obesity CHIS, 2009; Poverty Level American Community Survey, 2007-2009; Unemployment Rate State of California Employment Development Department, 2009
- 9. An asthma death is a death where asthma was indicated as the underlying cause on the death certificate. The rate of asthma deaths is the number of deaths per 1,000,000 residents, age-adjusted to the 2000 U.S. population.
- 10. An asthma ED visit is an admission to a licensed ED in California with the primary diagnosis of asthma. The rate of asthma ED visits is the number of visits per 10,000 residents, age-adjusted to the 2000 U.S. population.
- Population denominators for rates are from the California Department of Finance. All rates are age-adjusted to the 2000 U.S. population. Age-adjusted rates are modified to eliminate the effect of different age distributions in different populations. Rates based on numbers <20 are not reported.
- 12. An asthma hospitalization is a discharge from a licensed acute care hospital in California with the primary diagnosis of asthma. The rate of asthma hospitalizations is the number of hospitalizations per 10,000 residents, age-adjusted to the 2000 U.S. population.
- 13. Charges for asthma hospitalizations are the only type of data available to assess the costs of asthma in California counties. However, there are many other costs associated with asthma, including other types of health care utilization, medications, and indirect costs due to factors such as school and work missed.
- 14. Healthy People 2010 (HP2010) is a set of national benchmarks for a wide range of health topics, including asthma. For more information on HP2010, visit
- www.healthypeople.gov (/www.healthypeople.gov).

 15. Outdoor air quality data—including exposures such as PM2.5, PM10, ozone, and traffic pollution—can be found online through the California Environmental Health Tracking Program's Air Quality Data Query (http://www.ehib.org/page_isp?page_key=80) or on the California Air Resources Board (http://www.arb.ca.gov/aqd/aqdpage.htm) website.

Further details about the data presented in this report can be found in the accompanying Technical Notes document.



Asthma Data The Asthma in California Report (/asthma-data/cal-asthma-report) County Asthma Profiles (/asthma-data/county-asthma-profiles) County Comparisons (/asthma-data/county-comparisons) Custom Data Request (/asthma-data/datarequest) Fact Sheets

(/asthma-data/factsheets)

Projects California Asthma Quality Improvement Initiative Resources Publications (/resources/publications) Childcare Training Video Collaborations CAP (/collaborations/cap) SEHAC (Schools) (/collaborations/sehac) CHHC (Housing)

About Us Contact Us (/about-us/contact-us) (/about-us/our-vision) (/about-us/our-team)