Home (/) Asthma Data County Asthma Profiles (/asthma-data/county-asthma-profiles) Amador County Asthma Profile

# Amador County Asthma Profile

modified on: Thursday, 01 September 2016

This county profile may be missing graphs due to insufficient data. In addition, some data are combined into the **Eastern Counties** group, which includes Alpine, Amador, Calaveras, Inyo, Mariposa, Mono, and Tuolumne

### In Eastern Counties, approximately 19,000 children and adults have been diagnosed with asthma.

Scroll down to browse the full County Profile. Or, download a printer-friendly PDF.

# **COUNTY PROFILE INDICATORS**

Lifetime Asthma Prevalence	Emergency Department Visits
Active Asthma Prevalence	Asthma Hospitalizations
Work-Related Asthma	Asthma Deaths
Asthma Management Plans	Outdoor Air Quality <sup>15</sup>
Asthma Risk Factors	Notes

	County Population			County Popula	ation
Age		Population	%	Race/Ethnicity <sup>1</sup>	%
	0-4	1,341	3.6	Asian/PI	1.4
	5-17	4,466	12.1	Black	2.6
	18-64	22,016	59.8	Hispanic	13.8
	65+	9,011	24.5	White	77.9
	Total	36,833		Other	4.2

DATASOURCE:California Department of Finance, 2014 (2013 release)

### LIFETIME ASTHMA PREVALENCE,2 2011-2012

#### Percent with Lifetime Asthma (95% Confidence Interval3)

	Age	Eastern Counties	California
Children	0-4		9.7 (7.2-12.3)
	5-17		17.1 (15.5-18.7)
Adults	18-64	10.5 (5.8-15.2)	14.0 (13.4-14.7)
	65+		12.0 (11.1-12.8)
Totals:	0-17		15.4 (14.0-16.7)
	18+	10.8 (6.3-15.3)	13.7 (13.1-14.3)
	All Ages	10.5 (6.5-14.6)	14.1 (13.6-14.6)

datasource: California Health Interview Survey (CHIS), 2014

# ACTIVE ASTHMA PREVALENCE,4 2014

#### Percent with Active Asthma (95% Confidence Interval3)

	Age	Eastern Counties	California
Children 0-4			6.3 (4.6-8.0)
	5-17	10.4* (2.4-18.5)	11.6 (10.2-13.0)
Adults	18-64	10.3* (2.1-18.5)	7.6 (7.1-8.1)
	65+	7.8 (3.9-11.6)	8.2 (7.5-8.9)
Totals:	0-17	7.9* (1.9-13.9)	10.1 (9.0-11.3)
	18+	9.4 (3.9-15.0)	7.7 (7.3-8.2)
	All Ages	5.0 (2.2-7.7)	8.3 (7.9-8.7)

datasource: California Health Interview Survey (CHIS), 2014

# WORK-RELATED ASTHMA5

Studies show that asthma is commonly caused or triggered by workplace exposures, but work-related asthma

(http://www.cdph.ca.gov/programs/ohsep/Pages/Asthma.aspx) is under-recognized and under-diagnosed. Research confirms that 15-30% of current adult asthma was initiated by work exposures, meaning that an estimated 1,650-3,300 adults in Eastern Counties have asthma caused by work.<sup>6</sup>

datasource: CHIS, 2007

# ASTHMA MANAGEMENT PLANS

National guidelines recommend that health care providers give all patients with asthma a written self-management plan. In Eastern Counties, 78.4%\* (95% CI 61.4-95.4) of people with asthma have NOT received a written asthma management plan from a health care provider.

datasource: CHIS, 2014

## ASTHMA RISK FACTORS, 2014

### Eastern Counties Risk Factors

Risk Factor	Percent (95% Confidence Interval)
Percent of adults who are current smokers	16.9 (10.4-23.4)
Percent of adults and children exposed to second- hand smoke in the home	8.0 (3.8-12.3)
Percent of adults who are obese <sup>7</sup> (BMI>=30)	18.7 (13.9-23.5)
Percent of people below the Federal Poverty Level (Amador County only)	8.3 (NA)
Unemployment Rate (Amador County only)	11.7 (NA)

datasource: CHIS, 2011-2012

## ASTHMA DEATHS,9 2008-2010

# Number of Deaths Due to Asthma (N) and Age-Adjusted Rate9(per 1,000,000 residents)

	Amador County California		Amador County		ornia
	Age	N	Rate	N	Rate
Children	0-17	0		57	1.9
Adults	18+	0		1,198	14.3
Totals	All Ages	0		1,255	11.1

datasource: California Death Statistical Master Files, 2008-2010

### ASTHMA EMERGENCY DEPARTMENT VISITS, 10 2014

# Number of ED Visits Due to Asthma (N) and Age-Adjusted Rate11(per 10,000 residents)

		Amado	Amador County		California	
	Age	N	Rate	N	Rate	
Children	0-4	11		4,994	19.6	
	5-17	35	81.0	5,037	7.7	
Adults	18-64	135	60.2	12,134	4.7	
	65+	46	51.0	7,208	15.2	
Totals:	0-17	46	81.1	10,031	10.9	
	18+	181	58.7	19,342	6.5	
	All Ages	227	64.4	29,373	7.6	

datasource: Office of Statewide Health Planning and Development (OSHPD), 2014

### Expected Source of Payment for Asthma ED Visits

Payment Source	Amador County	California
Medicare	28.19%	13.29%
Medi-Cal	23.35%	48.56%
Private	40.97%	26.32%
Other	7.49%	11.84%

datasource: Office of Statewide Health Planning and Development (OSHPD), 2014

# ASTHMA HOSPITALIZATIONS,12 2014

# Number of Hospitalizations Due to Asthma (N) and Age-Adjusted Rate11(per

10,000 residents)						
		Amador County		Califo	California	
	Age	N	Rate	N	Rate	
Children	0-4	0		5,539	22.1	
	5-17	0		5,180	7.8	
Adults	18-64	11		13,395	5.4	
	65+	12	13.7	8,119	18.1	
Totals:	0-17	0	-	10,719	11.7	
	18+	23	5.7	21,514	7.5	
	All Ages	23	4.2	32,233	8.6	

datasource: Office of Statewide Health Planning and Development (OSHPD), 2014

### Average Charges13Per Asthma Hospitalization

	Age	Amador County	California
Children	0-17		\$27,127.04
Adults	18+	\$39,202.90	\$46,783.17
Total	All Ages	\$39,202.90	\$39,860.44

datasource: Office of Statewide Health Planning and Development (OSHPD), 2014


### **Expected Source of Payment for Asthma Hospitalizations**

Payment Source	Amador County	California
Medicare	56.52%	28.24%
Medi-Cal	26.09%	46.46%
Private	4.35%	20.34%
Other	13.04%	4.96%

datasource: Office of Statewide Health Planning and Development (OSHPD), 2014

# HOSPITALIZATION RATES OVER TIME

### Age-Adjusted Asthma Hospitalizations per 10,000 Residents, Alameda County and California, 2004-2014



Data Source: datasource: Office of Statewide Health Planning and Development (OSHPD) , 2004- 2014.

#### NOTES


\*Estimates marked with an asterisk (\*) have large confidence intervals, interpret with caution. Estimates marked with a dash (-) are statistically unstable, so are unreported.

- 1. PI = Pacific Islander; Please see technical notes for more information on race/ethnicity categorizations
- Lifetime asthma prevalence is the proportion of people in the population who have ever been diagnosed with asthma by a health provider.
   The 95% confidence interval (CI) is a range that expresses a level of certainty about an estimate based on the margin of error.
- The 95% CI means that we are 95 percent confident that this range contains the true population percent. A narrow CI means that there is less variability in the estimate and/or there is a larger sample size. A wide CI indicates more variability and/or a smaller sample size.

  4. Active asthma prevalence is the proportion of people in the population who have ever been diagnosed with asthma by a health provider and report that they still have asthma
- and/or report that they had an episode or attack within the past 12 months.
- 5. Work-related asthma is asthma that is caused or triggered by conditions or substances in the workplace.
- 6. Balmes J, Becklake M, Blanc P, et al. Environmental and Occupational Health Assembly, American Thoracic Society. American Thoracic Society Statement: Occupational Contribution to the Burden of Airway Disease. Am J Respir Crit Care Med. 2003;167:787-797; Lutzker L, Rafferty A, Brunner W, et al. Prevalence of Work-related Asthma in Michigan, Minnesota, and Oregon. Journal of Asthma. 2010;47:156-161.
- Obesity is defined as a body mass index (BMI) of 30 or greater.
- Data Sources for Asthma Risk Factors: Smoking CHIS, 2009; Obesity CHIS, 2009; Poverty Level American Community Survey, 2007-2009; Unemployment Rate State of California Employment Development Department, 2009
- 9. An asthma death is a death where asthma was indicated as the underlying cause on the death certificate. The rate of asthma deaths is the number of deaths per 1,000,000 residents, age-adjusted to the 2000 U.S. population.
- 10. An asthma ED visit is an admission to a licensed ED in California with the primary diagnosis of asthma. The rate of asthma ED visits is the number of visits per 10,000 residents, age-adjusted to the 2000 U.S. population.
- Population denominators for rates are from the California Department of Finance. All rates are age-adjusted to the 2000 U.S. population. Age-adjusted rates are modified to eliminate the effect of different age distributions in different populations. Rates based on numbers <20 are not reported.
- 12. An asthma hospitalization is a discharge from a licensed acute care hospital in California with the primary diagnosis of asthma. The rate of asthma hospitalizations is the number of hospitalizations per 10,000 residents, age-adjusted to the 2000 U.S. population.
- 13. Charges for asthma hospitalizations are the only type of data available to assess the costs of asthma in California counties. However, there are many other costs associated with asthma, including other types of health care utilization, medications, and indirect costs due to factors such as school and work missed.
- 14. Healthy People 2010 (HP2010) is a set of national benchmarks for a wide range of health topics, including asthma. For more information on HP2010, visit
- www.healthypeople.gov (/www.healthypeople.gov).

  15. Outdoor air quality data—including exposures such as PM2.5, PM10, ozone, and traffic pollution—can be found online through the California Environmental Health Tracking Program's Air Quality Data Query (http://www.ehib.org/page\_jsp?page\_key=80) or on the California Air Resources Board (http://www.arb.ca.gov/aqd/aqdpage.htm) website.

Further details about the data presented in this report can be found in the accompanying Technical Notes document.



Asthma Data The Asthma in California Report (/asthma-data/cal-asthma-report) County Asthma Profiles (/asthma-data/county-asthma-profiles) County Comparisons (/asthma-data/county-comparisons) Custom Data Request (/asthma-data/datarequest) Fact Sheets (/asthma-data/factsheets)

Projects California Asthma Quality Improvement Initiative Resources Publications (/resources/publications) Childcare Training Video Collaborations CAP (/collaborations/cap) SEHAC (Schools) (/collaborations/sehac) CHHC (Housing)

About Us Contact Us (/about-us/contact-us) (/about-us/our-vision) (/about-us/our-team)