

Home   Asthma Data   County Asthma Profiles   **Modoc County Asthma Profile**

# Modoc County Asthma Profile

modified on: Thursday, 01 September 2016

This county profile may be missing graphs due to insufficient data. In addition, some data are combined into the **Northern Counties** group, which includes Lassen, Del Norte, Modoc, Plumas, Sierra, Siskiyou, and Trinity.

**In Northern Counties, approximately 19,000 children and adults have been diagnosed with asthma.**

Scroll down to browse the full County Profile. Or, download a printer-friendly PDF.

## COUNTY PROFILE INDICATORS

Lifetime Asthma Prevalence	Emergency Department Visits
Active Asthma Prevalence	Asthma Hospitalizations
Work-Related Asthma	Asthma Deaths
Asthma Management Plans	Outdoor Air Quality <sup>15</sup>
Asthma Risk Factors	Notes

County Population			County Population	
Age	Population	%	Race/Ethnicity <sup>1</sup>	%
0-4	466	4.9	Asian/PI	0.5
5-17	1,439	15.1	Black	0.7
18-64	5,456	57.3	Hispanic	14.6
65+	2,154	22.6	White	79.0
<b>Total</b>	<b>9,515</b>		<b>Other</b>	<b>5.2</b>

DATASOURCE: California Department of Finance, 2014 (2013 release)

## LIFETIME ASTHMA PREVALENCE,<sup>2</sup> 2011-2012

Percent with Lifetime Asthma (95% Confidence Interval <sup>3</sup> )			
	Age	Northern Counties	California
<b>Children</b>	0-4	--	9.7 (7.2-12.3)
	5-17	--	17.1 (15.5-18.7)
<b>Adults</b>	18-64	15.1 (9.0-21.3)	14.0 (13.4-14.7)
	65+	--	12.0 (11.1-12.8)
	0-17	--	<b>15.4 (14.0-16.7)</b>
<b>Totals:</b>	18+	<b>13.3 (8.4-18.1)</b>	<b>13.7 (13.1-14.3)</b>
	<b>All Ages</b>	<b>13.7 (9.1-18.4)</b>	<b>14.1 (13.6-14.6)</b>

datasource: California Health Interview Survey (CHIS), 2014

## ACTIVE ASTHMA PREVALENCE,<sup>4</sup> 2014

Percent with Active Asthma (95% Confidence Interval <sup>3</sup> )			
	Age	Northern Counties	California
<b>Children</b>	0-4	--	6.3 (4.6-8.0)
	5-17	--	11.6 (10.2-13.0)
<b>Adults</b>	18-64	6.4* (1.2-11.7)	7.6 (7.1-8.1)
	65+	6.2* (1.8-10.6)	8.2 (7.5-8.9)
	0-17	--	<b>10.1 (9.0-11.3)</b>
<b>Totals:</b>	18+	<b>6.4* (2.0-10.8)</b>	<b>7.7 (7.3-8.2)</b>
	<b>All Ages</b>	<b>8.5 (5.3-11.7)</b>	<b>8.3 (7.9-8.7)</b>

datasource: California Health Interview Survey (CHIS), 2014

## WORK-RELATED ASTHMA<sup>5</sup>

Studies show that asthma is commonly caused or triggered by workplace exposures, but work-related asthma is under-recognized and under-diagnosed. Research confirms that 15-30% of current adult asthma was initiated by work exposures, meaning that **an estimated 1,350-2,700 adults in Northern Counties have asthma caused by work.**<sup>6</sup>

datasource: CHIS, 2007

## ASTHMA MANAGEMENT PLANS

National guidelines recommend that health care providers give all patients with asthma a written self-management plan. **In Northern Counties, 91.0%\* (95% CI 83.2-98.8) of people with asthma have NOT received a written asthma management plan from a health care provider.**

datasource: CHIS, 2013-2014

## ASTHMA RISK FACTORS, 2014

Northern Counties Risk Factors	
Risk Factor	Percent (95% Confidence Interval)
Percent of adults who are current smokers	15.2(10.8-19.6)
Percent of adults and children exposed to second-hand smoke in the home	11.5 (7.4-15.5)
Percent of adults who are obese <sup>7</sup> (BMI>=30)	31.4 (25.6-37.3)
Percent of people below the Federal Poverty Level (Modoc County only)	--
Unemployment Rate (Modoc County only)	12.8 (NA)

datasource: CHIS, 2011-2012

## ASTHMA DEATHS,<sup>9</sup> 2008-2010

Number of Deaths Due to Asthma (N) and Age-Adjusted Rate <sup>9</sup> (per 1,000,000 residents)					
	Modoc County			California	
	Age	N	Rate	N	Rate
<b>Children</b>	0-17	0	--	57	1.9
<b>Adults</b>	18+	1	--	1,198	14.3
<b>Totals</b>	<b>All Ages</b>	<b>1</b>	<b>--</b>	<b>1,255</b>	<b>11.1</b>

datasource: California Death Statistical Master Files, 2008-2010

ASTHMA EMERGENCY DEPARTMENT VISITS,<sup>10</sup> 2014

Number of ED Visits Due to Asthma (N) and Age-Adjusted Rate <sup>11</sup> (per 10,000 residents)					
	Age	Modoc County		California	
		N	Rate	N	Rate
Children	0-4	<5	--	26,268	103.4
	5-17	<15	93.7	47,613	72.4
Adults	18-64	<45	92.3	95,689	39.2
	65+	<5	--	17,374	36.1
Totals:	0-17	15	74.9	73,881	80.7
	18+	45	79.1	113,063	38.6
	All Ages	60	78.0	186,944	49.5

datasource: Office of Statewide Health Planning and Development (OSHDP), 2014

## Expected Source of Payment for Asthma ED Visits

Payment Source	Modoc County	California
Medicare	10.00%	13.29%
Medi-Cal	53.33%	48.56%
Private	21.67%	26.32%
Other	15.00%	11.84%

datasource: Office of Statewide Health Planning and Development (OSHDP), 2014

ASTHMA HOSPITALIZATIONS,<sup>12</sup> 2014

Number of Hospitalizations Due to Asthma (N) and Age-Adjusted Rate <sup>11</sup> (per 10,000 residents)					
	Age	Modoc County		California	
		N	Rate	N	Rate
Children	0-4	0	--	4,994	19.6
	5-17	0	--	5,037	7.7
Adults	18-64	<8	--	12,134	4.7
	65+	<5	--	7,208	15.2
		0-17	0	10,031	10.9
Totals:	18+	8	--	19,342	6.5
		All Ages	8	29,373	7.6

datasource: Office of Statewide Health Planning and Development (OSHDP), 2014

Average Charges<sup>13</sup> Per Asthma Hospitalization

	Age	Modoc County	California
Children	0-17	--	\$27,127.04
Adults	18+	\$8,379.63	\$46,783.17
Total	All Ages	\$8,379.63	\$39,860.44

datasource: Office of Statewide Health Planning and Development (OSHDP), 2014

## Expected Source of Payment for Asthma Hospitalizations

Payment Source	Modoc County	California
Medicare	12.50%	28.24%
Medi-Cal	62.50%	46.46%
Private	--	20.34%
Other	25.00%	4.96%

datasource: Office of Statewide Health Planning and Development (OSHDP), 2014


## NOTES

\*Estimates marked with an asterisk (\*) have large confidence intervals, interpret with caution. Estimates marked with a dash (-) are statistically unstable, so are unreported.

1. PI = Pacific Islander; Please see technical notes for more information on race/ethnicity categorizations.
2. Lifetime asthma prevalence is the proportion of people in the population who have ever been diagnosed with asthma by a health provider.
3. The 95% confidence interval (CI) is a range that expresses a level of certainty about an estimate based on the margin of error.  
The 95% CI means that we are 95 percent confident that this range contains the true population percent. A narrow CI means that there is less variability in the estimate and/or there is a larger sample size. A wide CI indicates more variability and/or a smaller sample size.
4. Active asthma prevalence is the proportion of people in the population who have ever been diagnosed with asthma by a health provider and report that they still have asthma and/or report that they had an episode or attack within the past 12 months.
5. Work-related asthma is asthma that is caused or triggered by conditions or substances in the workplace.
6. Balmes J, Becklake M, Blanc P, et al. Environmental and Occupational Health Assembly, American Thoracic Society. American Thoracic Society Statement: Occupational Contribution to the Burden of Airway Disease. Am J Respir Crit Care Med. 2003;167:787-797; Lutzker L, Rafferty A, Brunner W, et al. Prevalence of Work-related Asthma in Michigan, Minnesota, and Oregon. Journal of Asthma. 2010;47:156-161.
7. Obesity is defined as a body mass index (BMI) of 30 or greater.
8. Data Sources for Asthma Risk Factors: Smoking — CHIS, 2009; Obesity — CHIS, 2009; Poverty Level — American Community Survey, 2007-2009; Unemployment Rate — State of California Employment Development Department, 2009
9. An asthma death is a death where asthma was indicated as the underlying cause on the death certificate. The rate of asthma deaths is the number of deaths per 1,000,000 residents, age-adjusted to the 2000 U.S. population.
10. An asthma ED visit is an admission to a licensed ED in California with the primary diagnosis of asthma. The rate of asthma ED visits is the number of visits per 10,000 residents, age-adjusted to the 2000 U.S. population.
11. Population denominators for rates are from the California Department of Finance. All rates are age-adjusted to the 2000 U.S. population. Age-adjusted rates are modified to eliminate the effect of different age distributions in different populations. Rates based on numbers <20 are not reported.
12. An asthma hospitalization is a discharge from a licensed acute care hospital in California with the primary diagnosis of asthma. The rate of asthma hospitalizations is the number of hospitalizations per 10,000 residents, age-adjusted to the 2000 U.S. population.
13. Charges for asthma hospitalizations are the only type of data available to assess the costs of asthma in California counties. However, there are many other costs associated with asthma, including other types of health care utilization, medications, and indirect costs due to factors such as school and work missed.
14. Healthy People 2010 (HP2010) is a set of national benchmarks for a wide range of health topics, including asthma. For more information on HP2010, visit [www.healthypeople.gov](http://www.healthypeople.gov).
15. Outdoor air quality data—including exposures such as PM2.5, PM10, ozone, and traffic pollution—can be found online through the California Environmental Health Tracking Program's Air Quality Data Query or on the California Air Resources Board website.

Further details about the data presented in this report can be found in the accompanying Technical Notes document.

## AIR QUALITY MAP



## Asthma Data

The Asthma in California Report  
County Asthma Profiles  
County Comparisons  
Custom Data Request  
Fact Sheets

## Projects

California Asthma Quality Improvement Initiative

## Resources

Publications  
Childcare Training Video

## Collaborations

CAP  
SEHAC (Schools)  
CHHC (Housing)

## About Us

Contact Us  
Our Vision  
Our Team

Subscribe