Home (/) Asthma Data County Asthma Profiles (/asthma-data/county-asthma-profiles) San Luis Obispo County Asthma Profile

San Luis Obispo County Asthma Profile

modified on: Thursday, 01 September 2016

In San Luis Obispo County, approximately 27,000 children and adults have been diagnosed with asthma.

Scroll down to browse the full County Profile. Or, download a printer-friendly PDF.

COUNTY PROFILE INDICATORS

Lifetime Asthma Prevalence	Emergency Department Visits
Active Asthma Prevalence	Asthma Hospitalizations
Work-Related Asthma	Asthma Deaths
Asthma Management Plans	Outdoor Air Quality ¹⁵
Asthma Risk Factors	Notes

County Population			County Population		
Age Population %		%	Race/Ethnicity ¹	%	
0-4	13,043	4.8	Asian/PI	3.4	
5-17	36,847	13.5	Black	2.0	
18-64	176,281	64.7	Hispanic	22.2	
65+	46,397	17.0	White	69.5	
Total	272,568		Other	2.9	

DATASOURCE:California Department of Finance, 2014 (2013 release)

LIFETIME ASTHMA PREVALENCE,2 2011-2012

Percent with Lifetime Asthma (95% Confidence Interval³)

	Age	San Luis Obispo County	California
Children	0-4		9.7 (7.2-12.3)
	5-17		17.1 (15.5-18.7)
Adults	18-64		14.0 (13.4-14.7)
	65+	7.6* (1.9-13.2)	12.0 (11.1-12.8)
Totals:	0-17	-	15.4 (14.0-16.7)
	18+	7.9* (0.9-14.9)	13.7 (13.1-14.3)
	All Ages	10.9 (6.8-14.9)	14.1 (13.6-14.6)

datasource: California Health Interview Survey (CHIS), 2014

ACTIVE ASTHMA PREVALENCE,4 2014

Percent with Active Asthma (95% Confidence Interval³)

	Age	San Luis Obispo County	California
Children	0-4		6.3 (4.6-8.0)
	5-17		11.6 (10.2-13.0)
Adults	18-64		7.6 (7.1-8.1)
	65+	7.6* (1.9-13.2)	8.2 (7.5-8.9)
Totals:	0-17	-	10.1 (9.0-11.3)
	18+	7.9* (0.9-14.9)	7.7 (7.3-8.2)
	All Ages	5.2 (3.5-6.9)	8.3 (7.9-8.7)

datasource: California Health Interview Survey (CHIS), 2014

WORK-RELATED ASTHMA5

Studies show that asthma is commonly caused or triggered by workplace exposures, but work-related asthma

(http://www.cdph.ca.gov/programs/ohsep/Pages/Asthma.aspx) is under-recognized and under-diagnosed. Research confirms that 15-30% of current adult asthma was initiated by work exposures, meaning that an estimated 2,100-4,200 adults in San Luis Obispo County have asthma caused by work.6

datasource: CHIS, 2014

ASTHMA MANAGEMENT PLANS

National guidelines recommend that health care providers give all patients with asthma a written self-management plan. In San Luis Obispo County, - of people with asthma have NOT received a written asthma management plan from a health care provider.

datasource: CHIS, 2014

ASTHMA RISK FACTORS, 2014

San Luis Obispo County Risk Factors

10.6 (6.7-14.5) 4.6 (2.1-7.2)
46 (24 7 2)
4.0 (2.1-7.2)
12.6 (9.2-15.9)
12.7 (NA)
9.0 (NA)

ASTHMA DEATHS,9 2008-2010

Number of Deaths Due to Asthma (N) and Age-Adjusted Rate⁹ (per 1,000,000 residents)

		San Luis	San Luis Obispo County		California	
	Age	N	Rate	N	Rate	
Children	0-17	1		57	1.9	
Adults	18+	5		1,198	14.3	
Totals	All Ages	6		1,255	11.1	

datasource: California Death Statistical Master Files, 2008-2010

ASTHMA EMERGENCY DEPARTMENT VISITS,10 2014

Number of ED Visits Due to Asthma (N) and Age-Adjusted Rate¹¹ (per 10,000 residents)

		San Luis O	bispo County	Califo	rnia
	Age	N	Rate	N	Rate
Children	0-4	95	72.8	26,268	103.4
	5-17	176	48.0	47,613	72.4
Adults	18-64	494	31.4	95,689	39.2
	65+	107	22.6	17,374	36.1
Totals:	0-17	271	54.7	73,881	80.7
	18+	601	29.9	113,063	38.6
	All Ages	872	36.3	186,944	49.5

datasource: Office of Statewide Health Planning and Development (OSHPD), 2014 Expected Source of Payment for Asthma ED Visits

Payment Source	San Luis Obispo County	California
Medicare	17.09%	13.29%
Medi-Cal	24.20%	48.56%
Private	45.87%	26.32%
Other	12.84%	11.84%

datasource: Office of Statewide Health Planning and Development (OSHPD), 2014

ASTHMA HOSPITALIZATIONS,12 2014

Number of Hospitalizations Due to Asthma (N) and Age-Adjusted Rate¹¹ (per 10,000 residents)

		San Luis O	bispo County	Califo	rnia
	Age	N	Rate	N	Rate
Children	0-4	29	22.2	4,994	19.6
	5-17	25	6.9	5,037	7.7
Adults	18-64	37	2.5	12,134	4.7
	65+	26	5.9	7,208	15.2
Totals:	0-17	54	11.0	10,031	10.9
	18+	63	3.0	19,342	6.5
	All Ages	117	5.1	29,373	7.6

datasource: Office of Statewide Health Planning and Development (OSHPD), 2014

Average	Cnarges 13	Per Astnma	Hospitalization

	Age	San Luis Obispo County	California
Children	0-17	\$21,305.30	\$27,127.04
Adults	18+	\$83,914.40	\$46,783.17
Total	All Ages	\$55,017.89	\$39,860.44


Expected Source of Payment for Asthma Hospitalizations

Payment Source	San Luis Obispo County	California
Medicare	28.21%	28.24%
Medi-Cal	42.74%	46.46%
Private	20.51%	20.34%
Other	8.55%	4.96%

datasource: Office of Statewide Health Planning and Development (OSHPD), 2014

ASTHMA DISPARITIES


Age-Adjusted Asthma Hospitalizations and ED Visits per 10,000 San Luis Obispo County Residents by Race/Ethnicity, 2014



Data Source: datasource: Office of Statewide Health Planning and Development (OSHPD), 2014

HOSPITALIZATION RATES OVER TIME


Age-Adjusted Asthma Hospitalizations per 10,000 Residents, San Luis Obispo County and California, 2004-2014



Data Source: datasource: Office of Statewide Health Planning and Development (OSHPD), 2004-2014

HEALTHY PEOPLE 2010¹⁴

Asthma Hospitalizations per 10,000 Residents by Age, Compared to HP2020 Targets, California and San Luis Obispo County, 2014



Asthma ED Visits per 10,000 Residents by Age, Compared to HP2020 Targets, California and San Luis Obispo County, 2014

150 San Luis Obispo County California HP2020 103.4 100 95.7 rate 72.8 47.0 49.6 50 35.3 36.1 13.7 Age (years) Planning and Development (OSHPD), 2014. Data Source: datasource: Office of Statewide Health Pla

NOTES


Estimates marked with an asterisk () have large confidence intervals, interpret with caution. Estimates marked with a dash (-) are statistically unstable, so are unreported.

- 1. PI = Pacific Islander; Please see technical notes for more information on race/ethnicity categorizations
- Lifetime asthma prevalence is the proportion of people in the population who have ever been diagnosed with asthma by a health provider.
 The 95% confidence interval (CI) is a range that expresses a level of certainty about an estimate based on the margin of error.
- The 95% CI means that we are 95 percent confident that this range contains the true population percent. A narrow CI means that there is less variability in the estimate and/or there is a larger sample size. A wide CI indicates more variability and/or a smaller sample size.

 4. Active asthma prevalence is the proportion of people in the population who have ever been diagnosed with asthma by a health provider and report that they still have asthma
- and/or report that they had an episode or attack within the past 12 months.
- 5. Work-related asthma is asthma that is caused or triggered by conditions or substances in the workplace.
- 6. Balmes J, Becklake M, Blanc P, et al. Environmental and Occupational Health Assembly, American Thoracic Society. American Thoracic Society Statement: Occupational Contribution to the Burden of Airway Disease. Am J Respir Crit Care Med. 2003;167:787-797; Lutzker L, Rafferty A, Brunner W, et al. Prevalence of Work-related Asthma in Michigan, Minnesota, and Oregon. Journal of Asthma. 2010;47:156-161.
- Obesity is defined as a body mass index (BMI) of 30 or greater.
- Data Sources for Asthma Risk Factors: Smoking CHIS, 2009; Obesity CHIS, 2009; Poverty Level American Community Survey, 2007-2009; Unemployment Rate State of California Employment Development Department, 2009
- 9. An asthma death is a death where asthma was indicated as the underlying cause on the death certificate. The rate of asthma deaths is the number of deaths per 1,000,000 residents, age-adjusted to the 2000 U.S. population.
- 10. An asthma ED visit is an admission to a licensed ED in California with the primary diagnosis of asthma. The rate of asthma ED visits is the number of visits per 10,000 residents, age-adjusted to the 2000 U.S. population.
- Population denominators for rates are from the California Department of Finance. All rates are age-adjusted to the 2000 U.S. population. Age-adjusted rates are modified to eliminate the effect of different age distributions in different populations. Rates based on numbers <20 are not reported.
- 12. An asthma hospitalization is a discharge from a licensed acute care hospital in California with the primary diagnosis of asthma. The rate of asthma hospitalizations is the number of hospitalizations per 10,000 residents, age-adjusted to the 2000 U.S. population.
- 13. Charges for asthma hospitalizations are the only type of data available to assess the costs of asthma in California counties. However, there are many other costs associated with asthma, including other types of health care utilization, medications, and indirect costs due to factors such as school and work missed.
- 14. Healthy People 2010 (HP2010) is a set of national benchmarks for a wide range of health topics, including asthma. For more information on HP2010, visit
- www.healthypeople.gov (/www.healthypeople.gov).

 15. Outdoor air quality data—including exposures such as PM2.5, PM10, ozone, and traffic pollution—can be found online through the California Environmental Health Tracking Program's Air Quality Data Query (http://www.ehib.org/page_isp?page_key=80) or on the California Air Resources Board (http://www.arb.ca.gov/aqd/aqdpage.htm) website.

Further details about the data presented in this report can be found in the accompanying Technical Notes document.



Asthma Data The Asthma in California Report (/asthma-data/cal-asthma-report) County Asthma Profiles (/asthma-data/county-asthma-profiles) County Comparisons (/asthma-data/county-comparisons) Custom Data Request (/asthma-data/datarequest) Fact Sheets

(/asthma-data/factsheets)

Projects California Asthma Quality Improvement Initiative Resources Publications (/resources/publications) Childcare Training Video Collaborations CAP (/collaborations/cap) SEHAC (Schools) (/collaborations/sehac) CHHC (Housing)

About Us Contact Us (/about-us/contact-us) (/about-us/our-vision) (/about-us/our-team)