

Protect Yourself and Your Baby from Syphilis

What is Congenital Syphilis?

Syphilis is a sexually transmitted disease (STD).

Congenital syphilis occurs when a pregnant woman with syphilis passes the infection to her unborn child.

This can cause serious problems like premature birth, low birth weight, birth defects and stillbirth.

What are Symptoms of Syphilis?

Most people with syphilis have symptoms such as a sore or rash. Even if they do, they may not notice them.

The only way to know for sure is to get tested!

Getting tested for syphilis is part of routine prenatal care.

Who Should Get Tested?

If you are pregnant or might get pregnant, it is important to get routine prenatal care.

Getting tested for syphilis and other STDs is part of routine prenatal care.

Pregnant women should get syphilis testing at the first prenatal visit. Syphilis testing should be repeated during the third trimester (28-32 weeks gestational age) and at delivery in women who are at high risk for syphilis or live in areas with high rates of syphilis.

Be sure to get your syphilis test results and follow any medical advice at that time.

These clinics offer FREE or LOW-COST STD testing and treatment and pregnancy planning services.

Clinic Name1

Street Address

City

Phone: xxx.xxx.xxxx

Clinic hours (M-F, xx-xx)

Clinic Name4

Street Address

City

Phone: xxx.xxx.xxxx

Clinic hours (M-F, xx-xx)

Clinic Name2

Street Address

City

Phone: xxx.xxx.xxxx

Clinic hours (M-F, xx-xx)

Clinic Name5

Street Address

City

Phone: xxx.xxx.xxxx

Clinic hours (M-F, xx-xx)

Clinic Name3

Street Address

City

Phone: xxx.xxx.xxxx

Clinic hours (M-F, xx-xx)

Clinic Name6

Street Address

City

Phone: xxx.xxx.xxxx

Clinic hours (M-F, xx-xx)

For a complete list of free or low-cost clinics near you, visit <https://gettested.cdc.gov/> or call Public Health at xxx-xxx-xxxx.

County
Health Department
logo here

