

## **California Influenza and Other Respiratory Disease Surveillance for Week 12**

**(March 19, 2017 to March 25, 2017)**

Note: This report includes data from many sources of influenza surveillance and it should be viewed as a preliminary “snapshot” of influenza activity for each surveillance week. Because data are preliminary, the information may be updated in later reports as additional data are received. These data should not be considered population-based or representative of all California public health jurisdictions.

**Overall influenza activity in California was downgraded to “regional\*” during Week 12**

### **Influenza Report Highlights**

- Influenza activity in California remains elevated
- Outpatient influenza-like illness (ILI)
  - 2.0% of patient visits during Week 12 were for ILI, which is within expected levels for this time of year
- Hospitalization data
  - 5.4% of Kaiser patients hospitalized during Week 12 were admitted with a pneumonia or influenza (P&I) diagnosis, which is slightly above expected levels for this time of year
- Influenza virus detections by Respiratory Laboratory Network and Sentinel Laboratories
  - 314 (8.6%) of 3,663 specimens tested were positive for influenza during Week 12
- Influenza-associated deaths among patients 0–64 years of age
  - 5 laboratory-confirmed influenza deaths were reported during Week 12
- Influenza-associated outbreaks
  - 4 laboratory-confirmed influenza outbreaks were reported during Week 12

**\*Regional:** Outbreaks of influenza or increases in ILI and recent laboratory confirmed influenza in at least two but less than half the regions of the state with recent laboratory evidence of influenza in those regions.


## A. Outpatient and Inpatient Data

### 1. Influenza Sentinel Providers

Sentinel providers (physicians, nurse practitioners, and physician assistants) situated throughout California report on a weekly basis the number of patients seen with influenza-like illness (ILI) and the total number of patients seen for any reason. ILI is defined as any illness with fever ( $\geq 100^{\circ}\text{F}$  or  $37.8^{\circ}\text{C}$ ) AND cough and/or sore throat (in the absence of a known cause other than influenza).

A total of 73 enrolled sentinel providers have reported data for Week 12. Based on available data, the percentage of visits for ILI during Week 12 was 2.0% compared to Week 11 (2.3%) and is within expected levels for this time of year (Figure 1).

**Figure 1. Percentage of Influenza-like Illness Visits Among Patients Seen by California Sentinel Providers, 2012–2017**




### 2. Kaiser Permanente Hospitalization Data

Inpatients at Kaiser Permanente facilities with an admission diagnosis including the keywords “flu,” “influenza,” “pneumonia,” or variants of the keywords are defined as pneumonia and influenza (P&I) related admissions. The number of P&I admissions is divided by the total number of hospital admissions occurring in the same time period to estimate the percentage of P&I admissions. Admissions for pregnancy, labor and delivery, birth, and outpatient procedures are excluded from the denominator.

The percentage of hospitalizations for pneumonia and influenza (P&I) in Kaiser Permanente facilities in northern California during Week 12 was 5.4% compared to Week 11 (5.3%) and is slightly above expected levels for this time of the year (Figure 2).

**Figure 2. Percentage of P&I Hospitalizations in Kaiser Permanente Northern California Hospitals, 2012–2017**




The seasonal baseline was calculated using a regression model applied to data from the previous five years. The epidemic threshold is two standard deviations above the seasonal baseline and is the point at which the observed percentage of pneumonia and influenza hospitalizations in Kaiser Permanente hospitals in northern California is significantly higher than would be expected at that time of the year.

### 3. Influenza-Associated Hospitalizations, California Emerging Infections Program

The California Emerging Infections Program (CEIP), Influenza Surveillance Network (FluSurv-NET) conducts population-based surveillance for laboratory-confirmed influenza-associated hospitalizations among patients of all ages in Alameda, Contra Costa, and San Francisco counties.

The incidence of influenza-associated hospitalizations per 100,000 population was lower in Week 10 (0.75) compared to Week 9 (1.11) (Figure 3). Data for the most current two weeks are not presented because results are still being collected and are likely to change.

**Figure 3. Incidence of Influenza Hospitalizations in CEIP Counties, 2014–2017**



Note: The 2014-15 season contains a week 53. Data have been shifted so that week 1 aligns across years.

## **B. Laboratory Update - Influenza**


### 1. Respiratory Laboratory Network (RLN) and Sentinel Laboratory Surveillance Results

Laboratory surveillance for influenza and other respiratory viruses involves the use of data from hospital, academic, private and public health laboratories located throughout California. These laboratories report the number of laboratory-confirmed influenza and other respiratory virus detections and isolations on a weekly basis.


The percentage of influenza detections in the RLN and sentinel laboratories in Week 12 (8.6%) was lower than Week 11 (11.6%) (Figure 3). Additional details can be found in Figures 4 and 5 and Table 1.

Neither the RLN nor CDPH-VRDL has identified any influenza viruses by polymerase chain reaction (PCR) that are suggestive of a novel influenza virus.

**Figure 4. Percentage of Influenza Detections in Respiratory Laboratory Network and Sentinel Laboratories, 2012–2017**



**Figure 5. Number of Influenza Detections by Type and Subtype Detected in Respiratory Laboratory Network and Sentinel Laboratories, 2016–2017**



**Table 1. Respiratory Specimens Testing Positive for Influenza by Influenza Type and Subtype — Respiratory Laboratory Network and Sentinel Laboratories, Current Week and Season to Date**

	<b>Week 12 (Number)</b>	<b>Week 12 (Percent)</b>	<b>Season to Date (Number)</b>	<b>Season to Date (Percent)</b>
<b>Number of Specimens Tested</b>	3,663		106,337	
<b>Number of Specimens Positive for Influenza</b>	314	8.6*	19,105	18.0*
<b>Influenza Type/Subtype of Positive Specimens</b>				
A	185	58.9 <sup>†</sup>	18,207	95.3 <sup>†</sup>
2009 A (H1)	0	0.0 <sup>‡</sup>	142	0.8 <sup>‡</sup>
A (H3)	22	11.9 <sup>‡</sup>	3,996	21.9 <sup>‡</sup>
A, not subtyped	163	88.1 <sup>‡</sup>	14,069	77.3 <sup>‡</sup>
B	129	41.1 <sup>†</sup>	898	4.7 <sup>†</sup>

\* Percent of total specimens tested for influenza

<sup>†</sup> Percent of specimens positive for influenza

<sup>‡</sup> Percent of influenza A positives

## 2. Antiviral Resistance Testing

Of the influenza specimens tested by the CDPH-VRDL to date this season, none have been found to be resistant to Oseltamivir (Table 2).

**Table 2. Number of specimens tested for Oseltamivir resistance, 2016–2017**

	<b>Oseltamivir Resistance</b>
<b>Influenza 2009A (H1)</b>	0/26
<b>Influenza A (H3)</b>	0/69
<b>Influenza B</b>	0/19

## 3. Influenza Virus Strain Characterization

To date in California, all influenza 2009 A (H1) and A (H3) antigenically characterized viruses have matched the influenza 2009 A (H1) and A (H3) components included in the trivalent and quadrivalent influenza vaccines (Table 3). In addition, all influenza B antigenically characterized viruses in California have matched the influenza B Yamagata lineage virus included in the quadrivalent influenza vaccine.

**Table 3. Influenza virus antigenic characterization data — California and the United States, 2016– 2017**

Influenza Subtype/Lineage	Vaccine Strain	Match Vaccine Strain (California)	Match Vaccine Strain (United States)
<b>Influenza A (H1)</b>	A/California/7/2009-like (H1N1)	19/19	176/176
<b>Influenza A (H3)</b>	A/Hong Kong/4801/2014-like	66/66	535/547
<b>Influenza B Victoria*</b>	B/Brisbane/60/2008-like	3/3	154/167
<b>Influenza B Yamagata†</b>	B/Phuket/3073/2013-like	9/9	179/179

\*The influenza B Victoria lineage virus is included in only the 2016–2017 trivalent and quadrivalent influenza vaccine

†The influenza B Yamagata lineage virus is included in both the 2016–2017 quadrivalent influenza vaccines

### **C. Laboratory-Confirmed Severe Influenza Case Reports**

Currently, as mandated under Section 2500 of the California Code of Regulations, deaths among patients aged 0–64 years with laboratory-confirmed influenza are reportable to CDPH. The weekly influenza report includes confirmed deaths formally reported to CDPH as of March 25, 2017 (Week 12).

During Week 12, five laboratory-confirmed influenza fatalities were reported. To date, CDPH has received 80 reports of laboratory-confirmed influenza-associated deaths among patients <65 years of age during the 2016–2017 influenza season.

### **D. Influenza-Associated Outbreaks**

During Week 12, four laboratory-confirmed influenza outbreaks were reported. To date, 252 laboratory-confirmed influenza outbreaks have been reported to CDPH for the 2016–2017 season.

### **E. California Border Region Influenza Surveillance Network Data**


The border influenza surveillance network is comprised of outpatient provider sentinel sites whose geographical coverage extends approximately 100 kilometers (60 miles) north of the California-Baja California border and includes Imperial and San Diego Counties, as well as some parts of Riverside County.

#### 1. Syndromic Surveillance Update

A total of 11 border region sentinel providers reported data during Week 12, compared to 15 during Week 11 of 2017. The total number of patients screened by all sentinel sites for ILI during Week 12 was 10,017.

Outpatient ILI activity was the same in Week 12 (1.4%) as in Week 11 (1.4%). ILI activity for the California border region during Week 12 was lower when compared to activity for the same week during the 2014–2015 and 2015–2016 season (Figure 6). All influenza syndromic data summarized for the border region represents a subset of CDC influenza sentinel providers in California.


**Figure 6. Percentage of Influenza-like Illness Visits among Patients Seen by California Border Region Sentinel Providers, 2014–2017**



## 2. Virologic Surveillance Update

Cumulatively this season, a total of 8,323 respiratory specimens have been tested from border region clinics; of these, 644 (7.7%) tested positive for influenza. Of the 644 specimens that tested positive, 579 (89.9%) were influenza A and 65 (10.1%) were influenza B. Of the 579 specimens that tested positive for influenza A, 22 (3.8%) were subtyped as 2009 A (H1), 220 (38.0%) were subtyped as A (H3), and 337 (58.2%) had no further subtyping performed. None of the specimens that tested positive for influenza B were lineage typed. For Week 12, a total of 322 respiratory specimens were submitted for testing; 39 (12.1%) were positive for influenza. Of the 39 specimens that tested positive, 27 (69.2%) were influenza A and 12 (30.8%) were influenza B. Of the 27 specimens that tested positive for influenza A, 1 (3.7%) was subtyped as A (H3) and 26 (96.3%) had no further subtyping performed. The specimens that tested positive for influenza B were not lineage typed. Laboratory data summarized in Figure 7 includes data from influenza sentinel sites as well as laboratory data from other border region laboratories.

**Figure 7. Number of Influenza Detections by Type and Subtype Detected in California Border Region Respiratory Laboratory Network and Sentinel Laboratories, 2016–2017**



## F. Other Respiratory Viruses

### 1. Laboratory-Confirmed Severe Respiratory Syncytial Virus Case Reports


Currently, as mandated under Section 2500 of the California Code of Regulations, deaths among children aged 0–4 years with laboratory-confirmed respiratory syncytial virus (RSV) are reportable to CDPH. The weekly influenza report includes confirmed deaths formally reported to CDPH as of March 18, 2017 (Week 12).

During Week 12, no laboratory-confirmed RSV fatalities were reported. To date, CDPH has received four reports of laboratory-confirmed RSV-associated deaths among children <5 years of age during the 2016– 2017 influenza season.


### 2. Laboratory Update

During Week 12, 3,184 specimens were tested for RSV and 181 (5.7%) were positive, which was lower than Week 11 (8.0%) (Figure 8). Information on other respiratory viruses can be found in Figure 9.

**Figure 8. Percentage of RSV Detections in Respiratory Laboratory Network and Sentinel Laboratories, 2012–2017**



**Figure 9. Percentage of Other Respiratory Pathogen Detections in Respiratory Laboratory Network and Sentinel Laboratories, 2016–2017**



---

For questions regarding influenza surveillance and reporting in California, please email [InfluenzaSurveillance@cdph.ca.gov](mailto:InfluenzaSurveillance@cdph.ca.gov). This account is monitored daily by several epidemiologists.

For more information regarding the different influenza surveillance data sources, please visit the CDPH Influenza Surveillance Program at <http://www.cdph.ca.gov/programs/dcdc/Pages/CaliforniaInfluenzaSurveillanceProject.aspx>.

To obtain additional information regarding influenza, please visit the CDPH influenza website at [http://www.cdph.ca.gov/HealthInfo/discond/Pages/Influenza\(Flu\).aspx](http://www.cdph.ca.gov/HealthInfo/discond/Pages/Influenza(Flu).aspx).

A copy of the case report form for reporting any laboratory-confirmed influenza case that was either admitted to the ICU or died can be downloaded from [http://www.cdph.ca.gov/HealthInfo/discond/Pages/Influenza\(Flu\).aspx](http://www.cdph.ca.gov/HealthInfo/discond/Pages/Influenza(Flu).aspx).