






# Respiratory Virus Weekly Report

## Week 8: February 16, 2025 – February 22, 2025

This report gives information about SARS-CoV-2 (the virus that causes COVID-19) influenza (flu), Respiratory Syncytial Virus (RSV), and other respiratory viruses. It's a quick look at the activity for each week of surveillance and updates most Fridays. The data is early and might change in future reports as we have more information. This report doesn't cover all areas of California, so it might not represent the entire state's public health situation.

### - Report Highlights -

The 2024-2025 Respiratory Virus Season covers June 30, 2024 through June 28, 2025.

	COVID-19	FLU	RSV
 Test Positivity (change)	2.0% (-0.3)	20.3% (-3.0)	4.9% (-0.3)
 New Admissions Rate per 100,000 Persons (change)	1.7 (-0.5)	5.0 (-1.8)	N/A
 Percent of Total Deaths (change)	0.6% (-0.1)	1.9% (-0.4)	0.7% (0.2)
 Total Season Pediatric Deaths (new)	3 (0)	17 (2)	4 (1)
 Wastewater Concentrations (trend)	LOW (DECREASING)	N/A	N/A

### Key Messages

- Influenza activity remains high but continues to decrease. RSV activity remains low. COVID-19 activity is low.
- The majority of positive influenza specimens are type A with both H1 and H3 strains circulating. Influenza type B represents a small minority of positive specimens (<5%).
- CDPH data show that only a small percentage of eligible Californians have received the appropriate respiratory virus vaccines.

This year's respiratory virus season has been dominated by seasonal influenza. See CDC's [Preliminary Estimated Flu Disease Burden 2024-2025 Flu Season](#). With elevated respiratory virus activity, now is the time to take [added precautions](#) to prevent disease spread, especially if you have [risk factors for severe disease](#). Get vaccinated, protect yourself from serious illness, and reduce strain on our healthcare system. Updated COVID-19 and influenza vaccines are still available. Visit the [CDC's RSV vaccination](#) webpage for information about preventing RSV infection. Talk to your healthcare provider today.


For the most up to date information on bird flu, please see the [bird flu webpage](#).

# Laboratory Surveillance

## COVID-19

The overall percentage of SARS-CoV-2 detections from results received through electronic laboratory reporting (ELR) during the week ending February 22, 2025 was 2.0% compared to 2.3% during the week ending February 15, 2025.

**Figure 1. Percentage of SARS-CoV-2 Detections from Test Results Received through Electronic Laboratory Reporting, 2022/2023 Season to Date**




[Percentage of SARS-CoV-2 Detections from Test Results Received through Electronic Laboratory Reporting](#)

## Influenza

The overall percentage of influenza detections in clinical sentinel laboratories during the week ending February 22, 2025 was 20.3% compared to 23.3% during the week ending February 15, 2025.

**Figure 2. Percentage of Influenza Detections at Clinical Sentinel Laboratories, 2020/2021 Season to Date**




[Percentage of Influenza Detections at Clinical Sentinel Laboratories](#)

## Respiratory Syncytial Virus (RSV)

The overall percentage of RSV detections in clinical sentinel laboratories during the week ending February 22, 2025 was 4.9% compared to 5.2% during the week ending February 15, 2025.


**Figure 3. Percentage of RSV Detections at Clinical Sentinel Laboratories, 2020/2021 Season to Date**



[Percentage of RSV Detections at Clinical Sentinel Laboratories](#)

## Other Respiratory Viruses

Figure 4. Percentage of Other Respiratory Pathogen Detections at Clinical Sentinel Laboratories, June 30, 2024 to Date



[Percentage of Other Respiratory Pathogen Detections at Clinical Sentinel Laboratories](#)


[Lab Data Notes](#)

# Hospitalization Surveillance

## COVID-19

The overall rate of admissions for COVID-19 from NHSN during the week ending February 22, 2025 was 1.7 compared to 2.2 during the week ending February 15, 2025.

Figure 5. Rate of COVID-19 Admissions from NHSN from November 1, 2024 to Date




[Rate of COVID-19 Admissions from NHSN](#)

## Influenza

The overall rate of admissions for Influenza from NHSN during the week ending February 22, 2025 was 5.0 compared to 6.8 during the week ending February 15, 2025.

**Figure 6. Rate of Influenza Admissions from NHSN from November 1, 2024 to Date**




[Rate of Influenza Admissions from NHSN](#)

[NHSN Hospitalization Data Notes](#)

## Respiratory Syncytial Virus (RSV)

The overall percentage of admissions for RSV in Kaiser Permanente facilities in northern California during the week ending February 22, 2025 was 0.9% compared to 0.9% during the week ending February 15, 2025.

**Figure 7. Percentage of RSV Admissions at Kaiser Permanente Northern California Facilities, 2020/2021 Season to Date**



[Percentage of RSV Admissions at Kaiser Permanente Northern California Facilities](#)

[Kaiser Hospitalization Data Notes](#)




# Mortality Surveillance

## COVID-19

The overall percentage of deaths with COVID-19 listed anywhere on the death certificate during the week ending February 22, 2025 was 0.6% compared to 0.7% during the week ending February 15, 2025.


**Figure 8. Percentage of deaths attributed to COVID-19 in death certificates, 2022/2023 Season to Date**



## Influenza

The overall percentage of deaths with influenza listed anywhere on the death certificate during the week ending February 22, 2025 was 1.9% compared to 2.3% during the week ending February 15, 2025.

**Figure 9. Percentage of deaths attributed to Influenza in death certificates, 2020/2021 Season to Date**



## Respiratory Syncytial Virus (RSV)

The overall percentage of deaths with RSV listed anywhere on the death certificate during the week ending February 22, 2025 was 0.7% compared to 0.5% during the week ending February 15, 2025.

**Figure 10. Percentage of deaths attributed to RSV in death certificates, 2020/2021 Season to Date**





Figure 11. Age Distribution of COVID-19, Flu, and RSV-Coded Deaths from Death Certificates, 2024/2025 Season to Date



[Age Distribution of COVID-19, Flu, and RSV-Coded Deaths from Death Certificates](#)


[Death Data Notes](#)

# Wastewater Surveillance

## COVID-19

Statewide concentrations of SARS-CoV-2 RNA in wastewater are at a low level, and concentrations are decreasing compared to 21 days ago. Local and regional levels and trends may differ from state-wide estimates. Among individual sites, 71% are decreasing and 17% are increasing.

Figure 12. Aggregated SARS-CoV-2 RNA wastewater concentrations, from 2022/2023 Season to Date



[Aggregated SARS-CoV-2 RNA Wastewater Concentrations](#)

[Wastewater Data Notes](#)

## Resources

### Explore CDPH Data: COVID-19, Influenza, RSV

- COVID-19
  - [COVID-19 Resources](#)
  - [COVID-19 Vaccines](#)
  - [COVID-19 Vaccine Data](#)
  - [COVID-19 Health Equity](#)
  - [SARS-CoV-2 Variant Tracking](#)
  - [California Wastewater Surveillance Program](#)
  - [CDPH CalSuWers Wastewater Dashboard](#)
  - [CDPH Wastewater Surveillance Open Data Portal](#)
  - [Border Health Status Report](#)
- Influenza
  - [Influenza \(Flu\)](#)
  - [Novel Influenza](#)
  - [H5N1 Bird Flu](#)
- RSV
  - [RSV Resources](#)
- General Respiratory Virus Resources
  - [Respiratory Virus Report Datasets and Data Dictionaries on Open Data Portal](#)
  - [Respiratory Virus Toolkit](#)
  - [HAI Respiratory Virus Resources](#)

### Explore CDC and Other Resources

- COVID-19
  - [COVID-19 | CDC](#)
  - [COVID-19 Data Tracker for deaths, ED visits, and test positivity | CDC](#)
  - [COVID-19 Vaccine Resources | CDC](#)
- Influenza
  - [Influenza \(Flu\) | CDC](#)
  - [Weekly U.S. Influenza Surveillance Report \(FluView\) | CDC](#)
  - [FluView Interactive | CDC](#)
  - [FluVaxView- Flu Vaccine Coverage | CDC](#)
- RSV
  - [About RSV | CDC](#)
  - [RSVaxView- RSV Vaccine Coverage | CDC](#)
- General Respiratory Virus Resources
  - [Emergency Department \(ED\) Visits for Viral Respiratory Illnesses | CDC](#)
  - [Respiratory Virus Activity Levels | CDC](#)
  - [Respiratory Illness Data Channel | CDC](#)
  - [National Wastewater Surveillance System \(NWSS\) | CDC](#)
  - [WastewaterSCAN Dashboard](#)
  - [About Wastewater Data \(NWSS\) | CDC](#)

# About This Report

## More Information

- Accessible .csv files with data for all figures can be downloaded from the [Open Data Portal \(ODP\)](#).
- For questions regarding influenza and RSV surveillance and reporting in California, please email [InfluenzaSurveillance@cdph.ca.gov](mailto:InfluenzaSurveillance@cdph.ca.gov).
- For questions regarding COVID-19 surveillance and reporting in California, please email [COVIDepi@cdph.ca.gov](mailto:COVIDepi@cdph.ca.gov).

## Lab Data

- COVID-19
  - Data are received by the California Department of Public Health (CDPH) through electronic laboratory reporting of test results for COVID-19 among California residents.
  - COVID-19 testing data from Los Angeles County has an additional 7-day lag.
  - Test positivity is based on SARS-CoV-2 test results with a specimen collection date reported during a given week. Weeks are defined as Sunday through Saturday.
  - Test positivity is calculated by dividing the number of positive Nucleic Acid Amplification Test (NAAT) results by the total number of tests done.
- Influenza, Respiratory Syncytial Virus (RSV), and Other Respiratory Viruses (RVs)
  - CDPH gets data from volunteer labs in California called Sentinel Clinical Laboratories. These labs report how many tests were positive for influenza, RSV, and other RVs each week, and how many tests they did.
  - Volunteer labs include two large regional healthcare system laboratories covering counties in the Bay Area, Greater Sacramento, San Joaquin Valley, and Southern California regions of California, four children's hospital labs in the Bay Area (1), San Joaquin Valley (1) and Southern California (2), seven general acute care hospital labs in the Bay Area (2) and Southern California (5), and one clinic in Southern California.
  - These numbers don't include all the testing for influenza, RSV, and other RVs in California.
  - Test positivity is based on testing results reported during a given week. Weeks are defined as Sunday through Saturday. The data comes from labs that take part in the program.
  - Test positivity is the number of positive results divided by the total tests done.

## National Healthcare Safety Network (NHSN) Hospitalization Data

- COVID-19 and Influenza
  - On November 1, 2024, the new [NHSN rules](#) started. They require reporting COVID-19 and influenza hospital admissions.
  - CDPH pulls NHSN data from the Centers for Disease Control and Prevention (CDC) on the Wednesday prior to the publication of the report. Results may differ depending on which day data are pulled.
  - Admission rates are calculated using population estimates from the [P-3: Complete State and County Projections Dataset](#) provided by the state of California Department of Finance. Reported weekly admission rates for the entire season use the population estimates for the year the season started.
  - For more information on NHSN data including the protocol and data collection information, see the [CDC NHSN webpage](#).

## Kaiser Hospitalization Data

- Respiratory Syncytial Virus (RSV)
  - CDPH counts RSV-related admissions at Kaiser Permanente (KP) Northern California hospitals as those with diagnoses that include “RSV,” “syncytial,” or “bronchiolitis.” These numbers don’t always mean the admissions were confirmed by a lab.
  - To find the percentage of RSV admissions, divide the number of RSV admissions by the total number of admissions during the same period.
  - Admissions for pregnancy, labor and delivery, birth, and outpatient procedures are not included in the total number of admissions.

## Death Data

- COVID-19
  - COVID-19 death data come from CDPH’s Comprehensive Death File.
- Influenza and Respiratory Syncytial Virus (RSV)
  - Influenza and RSV death data come from CDPH. For 2020 to the present, the data is from the Comprehensive Death File (Dynamic). For 2018 and 2019, it’s from the Comprehensive Death File (Static).
- These numbers do not always mean the COVID-19, influenza or RSV deaths were confirmed by a laboratory.
- California counts deaths from COVID-19, influenza, and RSV for residents. It does so if the death certificate mentions any of these illnesses.

## Wastewater Surveillance Data for COVID-19

- These data represent wastewater data produced by multiple groups throughout California contributing to the California Surveillance of Wastewaters (Cal-SuWers) network. These groups include the California Department of Health Drinking Water and Radiation Laboratory (DWRL), [WastewaterSCAN](#) (through Verily laboratory), [CDC NWSS](#) (through Verily laboratory), and additional county, academic, and private programs in California. Not all wastewater surveillance programs in California participate in the Cal-SuWers network.
  - There is a special system for testing for the SARS-CoV-2 virus in wastewater. The testing locations are not evenly spread across the state. The number of sites, where they are, and the labs that work with them have changed a lot over time. For details about the sites involved, check the [CDPH Wastewater Surveillance](#) homepage.
- RNA is taken from wastewater samples and tested using PCR methods that look for a specific part of the N gene of SARS-CoV-2.
- The amount of SARS-CoV-2 in wastewater from all sites in California is combined to get a big picture of current trends and levels across the state. This is done using the [CDC’s Wastewater Viral Activity Level \(WWAL\)](#) method. The weekly state SARS-CoV-2 trend and level are only reported if at least 50% of sites in California measure at least one wastewater sample that week.
- Site specific wastewater data are available on the [CDPH Wastewater Surveillance Network Dashboard](#) and a downloadable wastewater dataset is available through the [California Health and Human Services Open Data Portal Wastewater Surveillance Data page](#).
- Wastewater trends are checked by looking at the average change in levels over the past three weeks. The changes are grouped into these categories:



- Decreasing: Less than -20% means levels are going down.
  - Plateau: Between -20% and +20% means levels are steady.
  - Increasing: More than 20% means levels are going up.
  - Strong increase: More than 100% means a big increase.
  - Very strong increase: More than 250% means a very big increase.
- Wastewater levels are evaluated by comparing the current concentration in wastewater to percentiles of historical concentrations over the past 365 days. If the current level is in the bottom third, compared to levels from the past year, it's considered low. If it's in the middle third, it's considered medium. If it's in the top third, it's considered high.

## Pediatric Deaths

- COVID-19
  - COVID-19 pediatric deaths require laboratory confirmation (positive PCR) and are sourced from the California Department of Public Health, California Comprehensive Death File (Dynamic).
- Influenza
  - Providers must report influenza-related deaths in children under 18. This is required by California law ([Title 17, Section 2500 of the California Code of Regulations \(PDF\)](#)). These deaths also need to be confirmed by a laboratory test.
- Respiratory Syncytial Virus (RSV)
  - Providers must report RSV-related deaths in children under 5. This is required by California law ([Title 17, Section 2500 of the California Code of Regulations \(PDF\)](#)). These deaths also need to be confirmed by a laboratory test.
- To find pediatric deaths from COVID-19, influenza, and RSV, the methods look at what's written on death certificates, and they do not use lab tests. This is different from how we identify these deaths when lab tests are required. Providers are required to report influenza and RSV pediatric deaths, following rules in [Title 17, Section 2500 \(PDF\)](#).