




# Figure 1. Number of Tuberculosis Cases: California, 1930-2012




# Figure 2. Number of Tuberculosis Cases and Case Rates: California, 2003-2012




# Figure 3. Tuberculosis Cases by Race/Ethnicity: California, 2012




# Figure 4. Tuberculosis Case Rates by Race/Ethnicity: California, 2003-2012




# Figure 5. Tuberculosis Cases in Foreign-born and U.S.-born Persons: California, 2003-2012




# Figure 6. Tuberculosis Cases by Country of Origin: California, 2012



# Figure 7. Tuberculosis Cases in Persons 0-4 Years of Age: California, 2003-2012





# Figure 8. Tuberculosis Cases by Verification Criteria: California, 2012




\*NAAT= Nucleic Acid Amplification Test

# Figure 9. Deaths in Persons with Tuberculosis: California, 2001-2010




# Figure 10. Tuberculosis Cases by HIV/AIDS Diagnosis\*: California, 2002-2011




\* Match found in HIV/AIDS Registry, California Office of AIDS

# Figure 11. HIV/AIDS-associated Tuberculosis\* by Race/Ethnicity: California, 2002-2011




\* Match found in HIV/AIDS Registry, California Office of AIDS

# Figure 12. Tuberculosis Cases with Multidrug Resistance (MDR) on Initial or Final Drug Susceptibility Testing\*: California, 2001-2010



\*Cases with resistance to at least isoniazid and rifampin reported on the Initial Drug Susceptibility Report (Follow-up 1) or on the Case Completion Report (Follow-up 2)


# Figure 13. Tuberculosis Cases with Initial Multidrug Resistance (MDR)\*: California, 2008-2012



98

\*Cases with initial drug resistance to at least isoniazid and rifampin


# Figure 14. Tuberculosis Cases<sup>1</sup> by Outcome of Therapy: California, 2010



<sup>1</sup> Patient was alive at diagnosis and started on an initial drug regimen of two or more drugs.

<sup>2</sup> Patient could not be located prior to the completion of treatment.


# Figure 15. Outcome in Tuberculosis Cases for Whom One Year or Less of Treatment was Indicated<sup>1</sup>: California, 2010



<sup>1</sup> Excludes cases with rifampin resistant disease, cases with meningeal disease, and cases less than 15 years of age with disseminated tuberculosis disease


<sup>2</sup> Patient could not be located prior to the completion of treatment.

# Figure 16. Outcome in Tuberculosis Cases for Whom Greater than One Year of Treatment was Indicated<sup>1</sup>: California, 2010



<sup>1</sup> Includes cases with rifampin resistant disease, cases with meningeal disease, and cases less than 15 years of age with disseminated tuberculosis disease

# Figure 17. Completion of Tuberculosis Therapy<sup>1</sup>: California, 2001-2010



<sup>1</sup> Excludes cases with rifampin resistant disease, cases with meningeal disease, cases less than 15 years of age with disseminated tuberculosis disease, and those that died during treatment