

CLABSI Table 5. Central Line-Associated Bloodstream Infection (CLABSI) Rates and Central Line Insertion Practices (CLIP) Adherence Percentages in Medical Critical Care Units, Reported by California Major Teaching Hospitals, January - December 2014

Hospitals highlighted in red did not participate in 2014 data validation; completeness of reported infections has not been evaluated

County	Hospital Name	CLABSI count	Central line-days	Rate per 1000*	95% Confidence Interval**	Statistical Interpretation***	CLIP Adherence %^
	STATE OF CALIFORNIA POOLED DATA	100	62875	1.59			97.6
Fresno	Community Regional Medical Center	4	2078	1.92	(0.52,4.93)	N	98.8
Los Angeles	Cedars-Sinai Medical Center	2	5464	0.37	(0.04,1.32)	L	95.1
Los Angeles	Kaiser Foundation Hospital, Los Angeles	0	2489	0	(0.00,1.20)	L	90.7
Los Angeles	Keck Hospital of USC	1	1536	0.65	(0.02,3.63)	N	99.1
Los Angeles	LAC+USC Medical Center	8	5048	1.58	(0.68,3.12)	N	99.3
<i>Los Angeles</i>	<i>LAC/Harbor-UCLA Medical Center</i>	<i>3</i>	<i>2325</i>	<i>1.29</i>	<i>(0.27,3.77)</i>	<i>N</i>	<i>97.8</i>
<i>Los Angeles</i>	<i>Los Angeles County Olive View-UCLA Medical Center</i>	<i>3</i>	<i>2105</i>	<i>1.43</i>	<i>(0.29,4.16)</i>	<i>N</i>	<i>99.6</i>
Los Angeles	Ronald Reagan UCLA Medical Center	36	13867	2.6	(1.82,3.59)	H	99.9
Orange	University of California Irvine Medical Center	1	4488	0.22	(0.01,1.24)	L	99.2
Sacramento	University of California Davis Medical Center	5	2424	2.06	(0.67,4.81)	N	98.6
San Bernardino	Arrowhead Regional Medical Center	1	2058	0.49	(0.01,2.71)	N	99.4
San Bernardino	Loma Linda University Medical Center	17	7080	2.4	(1.40,3.84)	N	94.3
San Diego	Scripps Mercy Hospital	1	1315	0.76	(0.02,4.24)	N	95
San Diego	University of California, San Diego Medical Center	13	3190	4.08	(2.17,6.97)	H	97
San Francisco	San Francisco General Hospital	2	2329	0.86	(0.10,3.10)	N	100
Santa Clara	Santa Clara Valley Medical Center	1	3077	0.32	(0.01,1.81)	N	95.4
Santa Clara	Stanford Hospital	2	2002	1	(0.12,3.61)	N	99.6

*Rate per 1000 central line days

**95% confidence interval calculated for rates based on exact Poisson distribution.

*** Statistical interpretation indicates hospital-specific rates that were statistically higher (H), lower (L), or no different (N) than California average rates, or rates that were not tested due to too few central line-days or too few hospitals reported (---).

^The CLIP bundle includes eight components: hand hygiene, recommend skin preparation agent, allowing the skin preparation agent to dry, and five maximal sterile barriers (cap, mask, gown, gloves, and drape).

Source: Central Line-Associated Bloodstream Infections and Central Line Insertion Practices in California Hospitals, January 2014 through December 2014 (Inclusive).