

CLABSI Table 35. Central Line-Associated Bloodstream Infection (CLABSI) Rates Related to Permanent-Type Central Lines in Bone Marrow Transplant Care Units, Reported by California Hospitals, January - December 2014

County	Hospital Name	CLABSI count	Central line-days	Rate per 1000*	95% Confidence Interval**	Statistical Interpretation***
	STATE OF CALIFORNIA POOLED DATA	70	23763	2.95		
Los Angeles	City of Hope Helford Clinical Research Hospital	39	11087	3.52	(2.50,4.81)	---
Los Angeles	Ronald Reagan UCLA Medical Center	5	455	10.99	(3.57,25.64)	---
Sacramento	Sutter General Hospital	6	3195	1.88	(0.69,4.09)	---
Sacramento	University of California Davis Medical Center	10	4390	2.28	(1.09,4.19)	---
San Diego	University of California, San Diego Medical Center	5	1235	4.05	(1.31,9.45)	---
Santa Clara	Stanford Hospital	5	3401	1.47	(0.48,3.43)	---

*Rate per 1000 central line days.

**95% confidence interval calculated for rates based on exact Poisson distribution.

*** Statistical interpretation indicates hospital-specific rates that were statistically higher (H), lower (L), or no different (N) than California average rates, or rates that were not tested due to too few central line-days or too few hospitals reported (---).

Source: Central Line-Associated Bloodstream Infections and Central Line Insertion Practices in California Hospitals, January 2014 through December 2014 (Inclusive).