

CLABSI Table 38. Central Line-Associated Bloodstream Infection (CLABSI) Rates Related to Permanent-Type Central Lines in Solid Organ Transplant Care Units, Reported by California Hospitals, January - December 2014

Hospitals highlighted in red did not participate in 2014 data validation; completeness of reported infections has not been evaluated.

County	Hospital Name	CLABSI count	Central line-days	Rate per 1000*	95% Confidence Interval**	Statistical Interpretation***
	STATE OF CALIFORNIA POOLED DATA	5	2099	2.38		
Los Angeles	Keck Hospital of USC	0	28	.	-----	---
<i>Los Angeles</i>	<i>LAC/Harbor-UCLA Medical Center</i>	<i>0</i>	<i>90</i>	<i>0</i>	<i>(0.00,33.29)</i>	<i>---</i>
Los Angeles	Ronald Reagan UCLA Medical Center	2	189	10.58	(1.28,38.23)	---
Los Angeles	Saint Vincent Medical Center	0	624	0	(0.00,4.80)	---
Riverside	Riverside Community Hospital	2	114	17.54	(2.12,63.37)	---
San Diego	Sharp Memorial Hospital	0	690	0	(0.00,4.34)	---
San Francisco	UCSF Medical Center	1	364	2.75	(0.07,15.31)	---

*Rate per 1000 central line days.

**95% confidence interval calculated for rates based on exact Poisson distribution.

*** Statistical interpretation indicates hospital-specific rates that were statistically higher (H), lower (L), or no different (N) than California average rates, or rates that were not tested due to too few central line-days or too few hospitals reported (---).

Source: Central Line-Associated Bloodstream Infections and Central Line Insertion Practices in California Hospitals, January 2014 through December 2014 (Inclusive).