

**CLABSI Table 42. Central Line-Associated Bloodstream Infection (CLABSI) Rates Related to Temporary-Type Central Lines in Bone Marrow Transplant Care Units, Reported by California Hospitals, January - December 2014**

County	Hospital Name	CLABSI count	Central line-days	Rate per 1000*	95% Confidence Interval**	Statistical Interpretation***
	<b>STATE OF CALIFORNIA POOLED DATA</b>	<b>139</b>	<b>43709</b>	<b>3.18</b>		
Los Angeles	City of Hope Helford Clinical Research Hospital	56	17583	3.18	(2.41,4.14)	---
Los Angeles	Ronald Reagan UCLA Medical Center	28	7950	3.52	(2.34,5.09)	---
Sacramento	Sutter General Hospital	4	3758	1.06	(0.29,2.73)	---
Sacramento	University of California Davis Medical Center	15	4031	3.72	(2.08,6.14)	---
San Diego	University of California, San Diego Medical Center	27	6751	4	(2.64,5.82)	---
Santa Clara	Stanford Hospital	9	3636	2.48	(1.13,4.70)	---

\*Rate per 1000 central line days.

\*\*95% confidence interval calculated for rates based on exact Poisson distribution.

\*\*\* Statistical interpretation indicates hospital-specific rates that were statistically higher (H), lower (L), or no different (N) than California average rates, or rates that were not tested due to too few central line-days or too few hospitals reported (---).

Source: Central Line-Associated Bloodstream Infections and Central Line Insertion Practices in California Hospitals, January 2014 through December 2014 (Inclusive).