

CLABSI Table 36. Central Line-Associated Bloodstream Infection (CLABSI) Rates Related to Permanent -Type Central Lines in Bone Marrow Transplant Care Units, Reported by California Hospitals, January-December 2015

| County | Hospital Name | CLABSI count | Central line days | Rate per 1000* | 95% Confidence Interval** | Statistical Interpretation*** |
|---------------|--|--------------|-------------------|----------------|---------------------------|-------------------------------|
| | STATE OF CALIFORNIA POOLED DATA | 87 | 26583 | 3.27 | | |
| Los Angeles | City of Hope Helford Clinical Research Hospital | 48 | 13440 | 3.57 | (2.63,4.74) | |
| Los Angeles | Ronald Reagan UCLA Medical Center | 3 | 1803 | 1.66 | (0.34,4.86) | |
| Sacramento | Sutter Medical Center, Sacramento | 6 | 2265 | 2.65 | (0.97,5.77) | |
| Sacramento | University of California Davis Medical Center | 12 | 4451 | 2.70 | (1.39,4.71) | |
| San Diego | University of California, San Diego Medical Center | 14 | 1088 | 12.87 | (7.03,21.59) | |
| San Francisco | UCSF Medical Center | 1 | 340 | 2.94 | (0.07,16.39) | |
| Santa Clara | Stanford Health Care | 3 | 3196 | 0.94 | (0.19,2.74) | |

*CLABSI rate per 1000 central line days.

**95% confidence interval calculated for rates based on exact Poisson distribution.

*** Statistical interpretation indicates hospital-specific rates that were statistically higher, lower, or no different than California average rates, or rates that were not tested due to too few central line days or too few hospitals reported .

Source: Central Line-Associated Bloodstream Infections and Central Line Insertion Practices in California Hospitals, January through December 2015. California Department of Public Health.