Table 19. Rates of central line-associated bloodstream infection in patient care locations providing pediatric step-down care, reported by California hospitals, January - December 2011 (inclusive)

Reporting hospital*	Cases	Central line-days	Rate	95% Confidence Interval	Statistical interpretation**
STATE OF CALIFORNIA POOLED DATA	**6**	**2609**	**2.30**		
LAC/HARBOR-UCLA MED CTR, TORRANCE	0	153	0.00	(0.00,19.58)	✦
RADY CHILDREN'S HOSP, SAN DIEGO	4	1897	2.11	(0.57,5.40)	✦
UNIVERSITY OF CALIFORNIA DAVIS MED CTR, SACRAMENTO	1	68	14.71	(0.37,81.94)	✦
† UNIVERSITY OF CALIFORNIA, SAN FRANCISCO	1	491	2.04	(0.05,11.35)	✦
UCSF MED CTR, SAN FRANCISCO					
UCSF MED CTR AT MOUNT ZION, SAN FRANCISCO					

* Hospitals with only one acute care campus associated with their license were listed by the campus name; hospitals with more than one campus associated with their license were listed by licensee names with acute care campus name(s) indented below

**Statistical interpretation indicates hospital-specific rates that were statistically higher (●), lower (○), or no different (⊙) than California average rates, or rates that were not tested due to too few central line-days or too few hospitals reported (✦)

† Hospital reported infection data aggregated over all acute care campuses

‡ Hospital reported infection data separately by acute care campus

Notes: Rate per 1000 central line days; 95% confidence interval calculated for rates based on exact Poisson distribution; ------ indicates data not shown due to too few central line-days reported; pediatric step down general care locations specialize in patients ≤18 years old that are hemodynamically stable who can benefit from close supervision and monitoring

Source: Central Line-Associated Bloodstream Infections in California Hospitals, January through December 2011. California Department of Public Health