

**Vaccines (other than influenza vaccine) in a multi-dose vial formulation  
that exceed the California legal limit for mercury content - July, 2006**

<b>Abbreviation</b>	<b>DT</b>	<b>TT</b>	<b>Td</b>	<b>MPSV4</b>
<b>Full name</b>	Diphtheria and Tetanus Toxoids Adsorbed USP (For Pediatric Use)	Tetanus Toxoid Adsorbed USP (5 Lf/dose)  Tetanus Toxoid For Booster Use Only (4 Lf/dose)	Tetanus and Diphtheria Toxoids Adsorbed For Adult Use	Meningococcal Polysaccharide Vaccine Groups A,C,Y and W-135 Combined (Trade name: <i>Menomune®-A/C/Y/W-135</i> )
<b>Manufacturer</b>	Sanofi Pasteur	Sanofi Pasteur	Massachusetts Biologic Laboratories	Sanofi Pasteur
<b>Formulation</b>	5 mL 10-dose vial	5 mL 10-dose vial	5 mL 10-dose vial	5 mL 10-dose vial
<b>Use in infants and toddlers?</b>	Infrequent	No; labeled for $\geq 7$ years	No; labeled for $\geq 7$ years	Rare; labeled for $\geq 2$ years
<b>Use in pregnant women?</b>	No; labeled up to 7 <sup>th</sup> birthday	Rare	Infrequent	Rare
<b>Alternatives which meet California law?</b>	0.5 ml single-dose vial of DT from same manufacturer	0.5 ml single-dose vial of TT from same manufacturer	0.5 ml single-dose vial of Td from Sanofi Pasteur	0.5 ml single-dose vial of MPSV4 from same manufacturer
<b>Additional comments</b>	Most infants and children should receive DTaP vaccine instead of DT.	Pregnant women should receive Td vaccine instead of TT.	This formulation of Td is seldom available outside of Massachusetts.	Pregnant women at high risk of meningococcal disease may receive meningococcal conjugate vaccine (MCV4) when available instead of MPSV4. As of June, 2006, FDA is reviewing a request for MCV4 to be re-labeled for use at $\geq 2$ years of age.