

SYPHILIS HEALTH CHECK™
RAPID POINT-OF-CARE SYPHILIS TESTING
Fact Sheet and Frequently Asked Questions (FAQ)



Syphilis Health Check™ (SHC) Facts

- Manufacturer: Trinity Biotech
- Qualitative rapid assay for detection of *Treponema pallidum* (syphilis) antibodies in human whole blood (fingerstick sample), serum or plasma
- 3rd generation enzyme immunoassay (EIA) format detecting both immunoglobulin G (IgG) and immunoglobulin M (IgM) antibodies
- Results available in 10 minutes
- 97-98% agreement with other reference treponemal assays, sensitivity may be reduced in primary disease
- Food and Drug Administration (FDA) cleared and Clinical Laboratory Improvement Amendments (CLIA) waived (may be performed in traditional clinical/laboratory settings and non-traditional settings such as mobile vans, health fairs, and other venues)
- Test kit contains: 20 test devices, pipettes, and 1 bottle of diluent. Shelf-stable at room temperature for 2 years from manufacture date.
- Control kit contains: 1 bottle each of positive/negative control. Controls must be refrigerated.
- Additional supplies needed: gloves, lancets, timer
- Price: \$400/kit, \$20/test; \$53/controls
- Billing and coding information: CPT: 86780; LOINC: 2411-09; HCPCS: 86592 (Medicare Part A); Reimbursement: \$18.06/test

How should SHC be used? Who can be tested with SHC?

SHC can be used for initial screening of asymptomatic patients, or SHC can be used in conjunction with a non-treponemal test (e.g. rapid plasma reagin [RPR]) and clinical findings to aid in diagnosis of symptomatic patients. Specific populations to target for screening include men who have sex with men, commercial sex workers and their clients, pregnant patients (especially with poor prenatal care), and injection drug users.

Who should NOT be tested with SHC?

Patients with a known prior history of syphilis should NOT receive testing with the SHC.

Treponemal antibodies often remain positive for life after previous infection, therefore the SHC cannot distinguish an old infection versus a new, current infection. For patients with a prior history of syphilis, a non-treponemal test (e.g. RPR or venereal disease research laboratory [VDRL]) should be performed instead of SHC.

Who can perform SHC in California?

In California, Business and Professions Code 1206.5 dictates which individuals can perform CLIA waived tests. Please refer to the code for a complete listing. **SHC cannot be performed** by disease intervention specialists, HIV test counselors or other field staff in California unless the person performing the test is covered under an included classification (refer to BPC 1206.5).

<http://law.onecle.com/california/business/1206.5.html>



What settings should be targeted for use of SHC?

SHC may be useful in 1) clinical settings such as emergency departments or urgent cares, sexually transmitted diseases (STD) clinics, HIV clinics, family planning clinics, practices providing prenatal care; 2) non-clinical settings offering limited services: corrections, drug treatment facilities; 3) community/outreach settings: mobile vans, commercial sex venues, needle exchange sites, health fairs.

How is the test performed?

Please refer to the package insert for step by step instructions for how to perform SHC.

<https://www.diagnosticsdirect2u.com/images/PDF/SyphilisHC.pdf>


How are results interpreted?

Results are positive, negative, or invalid.

Positive: Colored bands in the test area AND control area

Negative: 1 colored band in the control area

Invalid: No distinct color bands in either area OR 1 colored band in test area and no band in control area.



What if I find a patient with a positive SHC result in a non-clinical setting?

Draw blood for further testing in the field, if possible. Regardless, refer the patient to a local clinic for further testing and possible treatment.

What are reporting requirements for SHC results?

Positive SHC results should be reported to the health department similarly to other positive syphilis serology tests.

I'm interested in piloting the SHC in my local health jurisdiction (LHJ), is there any financial support for purchase of test kits?

Yes, please email your request to Christine Johnson and Matt Ayson at the California Department of Public Health (CDPH) STD Control Branch: christine.johnson@cdph.ca.gov and matt.ayson@cdph.ca.gov.

What if I have questions about interpretation of test results and/or treatment of patients with positive SHC results?

The CDPH STD Control Branch and the California STD-HIV Prevention Training Center (PTC) warmline clinical staff can answer questions regarding clinical management of patients with syphilis or other STDs. Clinicians should call 510-620-3400 and ask to speak to the clinician on-call. A link to current syphilis treatment recommendations can be found here:

<http://www.cdph.ca.gov/pubsforms/Guidelines/Documents/STD-Treatment-Guidelines-Color.pdf>