



Influenza and RSV Highlights



▲ **7.4%**
Influenza
positivity



▲ **5.2%**
Outpatient
ILI activity



▲ **0.7%**
Hospital
flu admissions





22 (+2)
Deaths
since
10/1/23 (new)



▲ **12.2%**
RSV
positivity

Influenza Activity Levels⁺



Geographic Area	Activity Level
California Statewide	Low
Northern Region	Low
Bay Area Region	Low
Central Region	Moderate
Upper Southern Region	Low
Lower Southern Region	Low

Key Messages

- » Influenza activity is increasing in California.
- » Now is a good time to get your flu shot to protect yourself against flu, its potentially serious complications, and reduce strain on our healthcare system.
- » Respiratory syncytial virus (RSV) activity is increasing in California.
- » Several [products](#) are available to prevent RSV infection.

In This Report

Influenza Laboratory Surveillance	3
Influenza Detections from Respiratory Laboratory Network and Clinical Sentinel Laboratories	3
Antiviral Resistance Testing	5
Influenza Outpatient, Inpatient, and Death Surveillance	6
Sentinel Provider Outpatient Visits for Influenza-like Illness	6
Influenza Admissions at Kaiser Permanente Northern California Facilities	7
Influenza Mortality Surveillance from Death Certificates	8
Laboratory-confirmed Influenza-associated Pediatric Deaths	10
Influenza-associated Outbreaks	11
California Border Region Influenza Surveillance Network	12
Syndromic Surveillance Update	12
Virologic Surveillance Update	12
Respiratory Syncytial Virus Surveillance	14
RSV Detections from Clinical Sentinel Laboratories	14
RSV Admissions at Kaiser Permanente Northern California Facilities	14
RSV Mortality Surveillance from Death Certificates	16
Laboratory-confirmed Respiratory Syncytial Virus-associated Deaths	18
Other Respiratory Viruses Surveillance	19
About This Report	20
More Information	20
Highlights Indicators	20
Influenza Activity Levels	20
California Regions	20


Influenza Laboratory Surveillance

Influenza Detections from Respiratory Laboratory Network and Clinical Sentinel Laboratories

Laboratory surveillance for influenza and other respiratory viruses involves the use of data from clinical sentinel laboratories and public health laboratories in the Respiratory Laboratory Network (RLN) located throughout California. These laboratories report the number of laboratory-confirmed influenza and other respiratory virus detections and isolations on a weekly basis.

The overall percentage of influenza detections in clinical sentinel laboratories during Week 47 was 7.4% compared to 6.4% during Week 46 ([Figure 1](#)). Additional details, including influenza typing and subtyping information from public health laboratories can be found in [Figures 1](#) and [2](#) and [Tables 1](#) and [2](#).

Figure 1. Percentage of Influenza Detections at Clinical Sentinel Laboratories, 2018–2024 Season to Date



Note: Data have been shifted so that Week 1 aligns across seasons.

Table 1. Respiratory Specimens Testing Positive for Influenza – Clinical Sentinel Laboratories, Current Week and Season to Date

	Current Week Number	Current Week Percent	Season to Date Number	Season to Date Percent
Number of Specimens Tested	18,086		84,778	
Influenza Positive	1,342	7.4	4,162	4.9
A	1,270	94.6*	3,855	92.6*
B	72	5.4*	307	7.4*

* Percentage of specimens positive for influenza

Figure 2. Number of Influenza Detections by Type and Subtype Detected in the Respiratory Laboratory Network, 2023–2024 Season to Date




Table 2. Respiratory Specimens Testing Positive for Influenza by Influenza Type and Subtype – Respiratory Laboratory Network, Current Week and Season to Date

	Current Week Number	Current Week Percent	Season to Date Number	Season to Date Percent
Influenza Positive	42		292	
A	40	95.2*	268	91.8*
A (H1)pdm09	33	82.5†	215	80.2†
A (H3)	4	10.0†	32	11.9†
A, not subtyped	3	7.5†	21	7.8†
B	2	4.8*	24	8.2*
B Victoria	0	0.0‡	13	54.2‡
B Yamagata	0	0.0‡	0	0.0‡
B, not lineage typed	2	100.0‡	11	45.8‡

† Percentage of specimens positive for influenza A

‡ Percentage of specimens positive for influenza B

Antiviral Resistance Testing

The Viral and Rickettsial Disease Laboratory (VRDL) assesses susceptibility of influenza viruses to antiviral drugs, including the neuraminidase inhibitors (oseltamivir, zanamivir, and peramivir) and the polymerase acidic (PA) endonuclease inhibitor (baloxavir), by using next-generation sequence analysis. Neuraminidase gene sequences are analyzed to detect the presence of well-known amino acid substitutions previously associated with reduced or inhibited activity to any of the three neuraminidase inhibitors. Susceptibility to baloxavir is assessed by identifying an amino acid substitution in the PA gene.

No influenza specimens have been tested for antiviral resistance by the CDPH-VRDL to date this season ([Table 3](#)).

Table 3. Number of Specimens Tested for Antiviral Resistance, 2023–2024 Season to Date

	Neuraminidase Inhibitor Resistance	PA Endonuclease Inhibitor
Influenza A (H1)pdm09	0/0	0/0
Influenza A (H3)	0/0	0/0
Influenza B	0/0	0/0


Influenza Outpatient, Inpatient, and Death Surveillance

Sentinel Provider Outpatient Visits for Influenza-like Illness

[Sentinel providers](#) (physicians, nurse practitioners, physician assistants) throughout California report on a weekly basis the number of patients seen with influenza-like illness (ILI) and the total number of patients seen for any reason. ILI is defined as any illness with fever ($\geq 100^{\circ}\text{F}$ or 37.8°C) AND cough and/or sore throat. Methods for calculating the 2023–2024 season baseline can be found on the [CDC website](#).

A total of 192 enrolled sentinel providers have reported data for Week 47. The percentage of visits for ILI during Week 47 was 5.2% compared to 4.8% during Week 46 and was above the baseline ([Figure 3](#)). Increases in ILI-related outpatient visits might also include people seeking care for other respiratory illnesses, including COVID-19.

Figure 3. Percentage of Influenza-like Illness Visits Among Patients Seen by California Sentinel Providers, 2018–2024 Season to Date




Note: Data have been shifted so that Week 1 aligns across seasons. The 2023–2024 baseline is developed by calculating the mean percentage of patient visits for ILI during non-influenza weeks for the most recent three seasons, excluding the COVID-19 pandemic, and adding two standard deviations. Additional information can be found on the CDC website.

Influenza Admissions at Kaiser Permanente Northern California Facilities

Inpatients at [Kaiser Permanente Northern California](#) facilities with an admission diagnosis including the keywords “flu,” “influenza,” or variants of the keywords are defined as influenza-related admissions. The number of influenza admissions is divided by the total number of hospital admissions occurring in the same time period to estimate the percentage of influenza admissions. Admissions for pregnancy, labor and delivery, birth, and outpatient procedures are excluded from the denominator. Influenza admission data is not comparable to previous seasons reports which included pneumonia and influenza (P&I) admissions.

The percentage of admissions for influenza in Kaiser Permanente Northern California facilities during Week 47 was 0.7% compared to 0.5% during Week 46 ([Figure 4](#)).


Figure 4. Percentage of Influenza Admissions at Kaiser Permanente Northern California Facilities, 2018-2024 Season to Date



Note: Data have been shifted so that Week 1 aligns across seasons.

To date, 66 non-intensive care unit (ICU) hospitalizations, nine ICU admissions, and no deaths have occurred among persons with influenza admission diagnoses. Most influenza admissions occurred among persons ≥ 65 years ([Figure 5](#)). Please note that influenza admissions serve as a proxy for influenza activity, but do not necessarily represent laboratory-confirmed influenza infections.

Figure 5. Age Group Distribution of Non-ICU, ICU, and Deaths Associated with Influenza Admissions at Kaiser Permanente Northern California Facilities, 2023–2024 Season to Date




Influenza Mortality Surveillance from Death Certificates

Deaths occurring in California among residents who had influenza noted in any cause of death field on the death certificate (text or coded) are defined as “influenza-coded deaths.” The percentage of influenza-coded deaths is calculated by dividing the number of influenza-coded deaths by the total number of all-cause deaths during the same period. Influenza-coded deaths are not necessarily laboratory-confirmed and are an underestimate of all influenza-associated deaths. Please note that during the 2023-2024 season, an update to the methods used to identify influenza-coded deaths resulted in some changes to data from previous seasons.


During Week 47, two influenza-coded deaths were identified. Six influenza-coded deaths were identified during Week 46. To date during the 2023–2024 winter respiratory virus season, 22 influenza-coded deaths have been identified ([Figure 6](#)). The percentage of deaths coded as influenza during Week 47 was 0.2% compared to 0.1% during Week 46 ([Figure 7](#)).

Figure 6. Number of Influenza-coded Deaths Identified from Death Certificates by Week of Death, 2023–2024 Season to Date



Note: Coding of deaths can be delayed by several weeks. Influenza-coded deaths will be included once enough information is available to identify them.


Figure 7. Percentage of Influenza-coded Deaths from Death Certificates, 2018–2024 Season to Date



Note: Data have been shifted so that Week 1 aligns across seasons.

To date, 17 (77.3%) influenza-coded deaths have been identified among persons ≥ 65 years of age during the 2023–2024 winter respiratory virus season ([Figure 8](#)).

Figure 8. Age Distribution of Influenza-coded Deaths from Death Certificates, 2018–2024 Season to Date



* Methods used to identify pediatric influenza-coded deaths on death certificates do not consider laboratory testing and thus differ from those used to identify the influenza-associated pediatric deaths presented below, which require laboratory confirmation of influenza.

[†] One death during the 2018–2019 winter respiratory virus season has unknown age and is not included in the figure.

[§] 2018–2019 season: Sept. 30, 2018–Sept. 28, 2019; mixed influenza A (H1N1)pdm09 and influenza A (H3N2) season
 2019–2020 season: Sept. 29, 2019–Sept. 26, 2020; mixed influenza B (Victoria) and influenza A (H1N1)pdm09 season
 2020–2021 season: Sept. 27, 2020–Oct. 2, 2021; influenza activity was too low to determine a predominant strain
 2021–2022 season: Oct. 3, 2021–Oct. 1, 2022; influenza A (H3N2) predominant season
 2022–2023 season: Oct. 2, 2022–Sept 30, 2023; influenza A (H3N2) predominant season

Laboratory-confirmed Influenza-associated Pediatric Deaths


Influenza-associated deaths in children <18 years of age are nationally notifiable. The weekly influenza report includes confirmed deaths formally reported to CDPH through November 25, 2023 (Week 47). Methods used to identify pediatric influenza-coded deaths on death certificates differ from those used to identify the influenza-associated pediatric deaths presented below, which require laboratory confirmation of influenza, and might not include the same individuals.

No laboratory-confirmed influenza-associated deaths in children <18 years of age were reported to CDPH during Week 47. To date, CDPH has received no reports of laboratory-confirmed influenza-associated deaths among persons <18 years of age during the 2023–2024 winter respiratory virus season.

Influenza-associated Outbreaks

No laboratory-confirmed influenza outbreaks were reported during Week 47. During Week 46, one laboratory-confirmed influenza outbreak was reported. To date, two laboratory-confirmed influenza outbreaks have been reported to CDPH during the 2023–2024 season.

Figure 9. Number of Laboratory-confirmed Influenza-associated Outbreaks by Week of First Onset, 2022–2024 Season to Date



*Earliest date associated with the outbreak was used for outbreaks without reported date of first patient's symptom onset.


California Border Region Influenza Surveillance Network

The border influenza surveillance network is comprised of outpatient sentinel provider sites whose geographical coverage extends approximately 100 kilometers (60 miles) north of the California-Baja California border and includes Imperial and San Diego Counties, as well as some parts of Riverside County.

Syndromic Surveillance Update

A total of 8 border region sentinel providers reported data during Week 47. The total number of patients screened by all sentinel sites for ILI during Week 47 was 7,595. Outpatient ILI activity was 0.8% in Week 47 ([Figure 10](#)). All influenza syndromic data summarized for the border region represent a subset of CDC influenza sentinel providers in California. Increases in ILI-related outpatient visits might also include people seeking care for other respiratory illness, including COVID-19. Methods for calculating the 2023–2024 season baseline can be found on the [CDC website](#).

Figure 10. Percentage of Influenza-like Illness Visits among Patients Seen by Sentinel Providers — California Border Region, 2018–2024 Season to Date



Note: Data have been shifted so that Week 1 aligns across seasons. The 2023–2024 baseline is developed by calculating the mean percentage of patient visits for ILI during non-influenza weeks for the most recent three seasons, excluding the COVID-19 pandemic, and adding two standard deviations. Additional information can be found on the CDC website.

Virologic Surveillance Update

The percentage of influenza detections in border region clinical sentinel laboratories during Week 47 was 9.2% ([Figure 11](#)). Additional details, including influenza typing, subtyping, and lineage typing information from border region clinical sentinel laboratories and RLN laboratories can be found in [Figure 11](#) and [Table 4](#).

Figure 11. Number of Influenza Detections by Type and Subtype Detected in RLN Laboratories and the Percentage of Specimens Testing Positive at Clinical Sentinel Laboratories – California Border Region, 2023–2024 Season to Date




Table 4. Respiratory Specimens Testing Positive for Influenza by Influenza Type, Subtype, and Lineage Type – Clinical Sentinel Laboratories and RLN, California Border Region, Current Week and Season to Date


	Current Week Number	Current Week Percent	Season to Date Number	Season to Date Percent
Clinical Sentinel Laboratories				
Number of Specimens Tested	642		4,009	
Influenza Positive	59	9.2	191	4.8
A	53	89.8*	167	87.4*
B	6	10.2*	24	12.6*
Respiratory Laboratory Network				
Influenza Positive	5		24	
A	3	60.0*	19	79.2*
A (H1)pdm09	3	100.0 [†]	16	84.2 [†]
A (H3)	0	0.0 [†]	3	15.8 [†]
A, not subtyped	0	0.0 [†]	0	0.0 [†]
B	2	40.0*	5	20.8*
B Victoria	0	0.0 [†]	0	0.0 [†]
B Yamagata	0	0.0 [†]	0	0.0 [†]
B, not lineage typed	2	100.0 [†]	5	100.0 [†]

Respiratory Syncytial Virus Surveillance

RSV Detections from Clinical Sentinel Laboratories

The overall percentage of RSV detections in clinical sentinel laboratories during Week 47 was 12.2% compared to 11.6% during Week 46 ([Figure 12](#)).

Figure 12. Percentage of RSV Detections at Clinical Sentinel Laboratories, 2018–2024 Season to Date




Note: Data have been shifted so that Week 1 aligns across seasons.

RSV Admissions at Kaiser Permanente Northern California Facilities

Inpatients at Kaiser Permanente Northern California facilities with an admission diagnosis including the keywords “RSV,” “syncytial,” “bronchiolitis,” and variants of the keywords are defined as respiratory syncytial virus (RSV)-related admissions. The number of RSV admissions is divided by the total number of hospital admissions occurring in the same period to estimate the percentage of RSV admissions. Admissions for pregnancy, labor and delivery, birth, and outpatient procedures are excluded from the denominator.

The percentage of admissions for RSV in Kaiser Permanente facilities in northern California during Week 47 was 1.4% compared to 1.0% during Week 46 ([Figure 13](#)).


Figure 13. Percentage of RSV Admissions in Kaiser Permanente Northern California Facilities, 2018–2024 Season to Date



Note: Data have been shifted so that Week 1 aligns across seasons.

To date, 158 non-intensive care unit (ICU) hospitalizations, 34 ICU admission, and two deaths have occurred among persons with RSV admission diagnoses. Most RSV admissions occurred among persons <18 years (Figure 14). Please note that RSV admissions serve as a proxy for RSV activity, but do not necessarily represent laboratory-confirmed RSV infections.

Figure 14. Age Group Distribution of Non-ICU, ICU, and Deaths Associated with RSV Admissions in Kaiser Permanente Northern California Facilities, 2023–2024 Season to Date




RSV Mortality Surveillance from Death Certificates

Deaths occurring in California among residents who had RSV noted in any cause of death field on the death certificate (text or coded) are defined as “RSV-coded deaths.” The percentage of RSV-coded deaths is calculated by dividing the number of RSV-coded deaths by the total number of all-cause deaths during the same period. RSV-coded deaths are not necessarily laboratory-confirmed and are likely to be an underestimate of all RSV-associated deaths.


During Week 47, four RSV-coded deaths were identified. Four RSV-coded deaths were identified during Week 46. To date during the 2023–2024 winter respiratory virus season, 11 RSV-coded deaths have been identified ([Figure 15](#)). The percentage of deaths coded as RSV during Week 47 was 0.16% compared to 0.09% during Week 46 ([Figure 16](#)).

Figure 15. Number of RSV-coded Deaths Identified from Death Certificates by Week of Death, 2023–2024 Season to Date



Note: Coding of deaths can be delayed by several weeks. RSV-coded deaths will be included once enough information is available to identify them.


Figure 16. Percentage of RSV-coded Deaths from Death Certificates, 2018–2024 Season to Date



Note: Data have been shifted so that Week 1 aligns across seasons.

To date, ten (90.9%) RSV-coded deaths have been identified among persons ≥ 65 years of age during the 2023–2024 winter respiratory virus season ([Figure 17](#)).

Figure 17. Age Distribution of RSV-coded Deaths from Death Certificates, 2018–2024 Season to Date



Laboratory-confirmed Respiratory Syncytial Virus-associated Deaths

Currently, as mandated under Section 2500 of the California Code of Regulations, deaths among children aged 0–4 years with laboratory-confirmed RSV are reportable to CDPH. The weekly influenza report includes confirmed deaths formally reported to CDPH through November 25, 2023 (Week 47).


No laboratory-confirmed RSV-associated deaths in children <5 years of age were reported to CDPH during Week 47. To date, CDPH has received no reports of laboratory-confirmed RSV-associated deaths among children <5 years of age during the 2023–2024 winter respiratory virus season.

[The report continues on the next page.]

Other Respiratory Viruses Surveillance

The percentage of other respiratory viruses detections in clinical sentinel laboratories can be found in [\(Figure 18\)](#) and [\(Table 5\)](#).

Figure 18. Percentage of Other Respiratory Pathogen Detections at Clinical Sentinel Laboratories, 2023-2024 Season to Date



*Coronaviruses identified include common human coronaviruses 229E, NL63, OC43, and HKU1 and do NOT include SARS-CoV-2.

Table 5. Respiratory Specimens Testing Positive for Other Respiratory Viruses – Clinical Sentinel Laboratories, Current Week and Previous Week

	Current Week Percent	Previous Week Percent
Adenovirus	2.8	3.9
Coronavirus (non SARS-CoV-2)	1.0	1.2
Enterovirus/Rhinovirus	21.7	23.2
Human Metapneumovirus	0.9	0.8
Parainfluenza types 1-4	4.4	5.8
SARS-CoV-2	6.7	5.4

About This Report

This report includes data from many sources of influenza and other respiratory virus surveillance, and it should be viewed as a preliminary “snapshot” of activity for each surveillance week. Because data are preliminary, the information may be updated in later reports as additional data are received. These data should not be considered population-based or representative of all California public health jurisdictions.

More Information

- › An accessible Excel file with data for all figures can be downloaded from the [CDPH Flu webpage](http://www.cdph.ca.gov/Programs/CID/DCDC/CDPH%20Document%20Library/Immunization/Week2023-2447_DataTables.xlsx) (www.cdph.ca.gov/Programs/CID/DCDC/CDPH%20Document%20Library/Immunization/Week2023-2447_DataTables.xlsx).
- › For questions regarding influenza surveillance and reporting in California, please email InfluenzaSurveillance@cdph.ca.gov.
- › To obtain additional information regarding influenza, please visit the [CDPH Influenza website](http://www.cdph.ca.gov/Programs/CID/DCDC/Pages/Immunization/Influenza.aspx) (www.cdph.ca.gov/Programs/CID/DCDC/Pages/Immunization/Influenza.aspx).
- › For information about national influenza activity, please visit the U.S. Centers for Disease Control and Prevention’s [FluView](http://www.cdc.gov/flu/weekly/index.htm) (www.cdc.gov/flu/weekly/index.htm) and [FluView Interactive](http://www.cdc.gov/flu/weekly/fluviewinteractive.htm) (www.cdc.gov/flu/weekly/fluviewinteractive.htm) websites.
- › For information about COVID-19 in California, please visit the [California COVID-19 website](http://www.covid19.ca.gov) (www.covid19.ca.gov).

Highlights Indicators

Triangle symbols are used to indicate direction of change between the previous week and the current week for laboratory flu positivity, outpatient ILI activity, and hospital flu admissions: Increase (▲), decrease (▼), no change (►).

[Back to Top of Report](#)

Influenza Activity Levels⁺

Minimal: The percentage of specimens positive for influenza is <2%.

Low: The percentage of specimens positive for influenza is between 2% and <10%.

Moderate: The percentage of specimens positive for influenza is between 10% and <20%.

High: The percentage of specimens positive for influenza is between 20% and <40%.

Very High: The percentage of specimens positive for influenza is ≥40%.

Insufficient Data: Total number of specimens tested is below the threshold for activity level determination.

California Regions

Northern: Alpine, Amador, Butte, Colusa, Del Norte, El Dorado, Glenn, Humboldt, Lake, Lassen, Mendocino, Modoc, Nevada, Placer, Plumas, Sacramento, Shasta, Sierra, Siskiyou, Sutter, Tehama, Trinity, Yolo, and Yuba counties

Bay Area: Alameda, Contra Costa, Marin, Napa, San Francisco, San Mateo, Santa Clara, Santa Cruz, Solano, and Sonoma counties

Central Valley: Calaveras, Fresno, Inyo, Kings, Madera, Mariposa, Merced, Mono, Monterey, San Benito, San Joaquin, Stanislaus, Tulare, and Tuolumne counties

Upper Southern: Kern, Los Angeles, San Luis Obispo, Santa Barbara, and Ventura counties

Lower Southern: Imperial, Orange, Riverside, San Bernardino, and San Diego counties

⁺ Influenza activity levels are derived from the percentage of specimens from clinical sentinel laboratories that tested positive for influenza.